## SUPPLEMENTAL LABELING

TOLFENPYRAD GROUP 21A INSECTICIDE GROUP 39 FUNGICIDE



EPA Reg. No. 71711-31

For Use On: Brassica Head and Stem Vegetable (Crop Group 5-16)

broccoli; Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; cultivars, varieties, and hybrids of these

commodities

Use Pattern: Aerial Application

This supplemental labeling expires 07/15/2025 and must not be used or distributed after this date.

## **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for TORAC<sup>®</sup> Insecticide before applying. Use of TORAC Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for TORAC Insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

## APPLICATION RATE CHART FOR TORAC INSECTICIDE

Brassica Head and Stem Vegetable (Crop Group 5-16)	
Rate/Acre	Use Directions
to 21.0 fl oz/acre to 0.21 lb ai/acre)	Aerial Application: Use a minimum of 5 gallons of water per acre.
to 21.0 fl oz/acre to 0.21 lb ai/acre)	<ul> <li>Ground Application: Use a minimum of 20 gallons of water per acre.</li> <li>USE RESTRICTIONS</li> <li>Do not apply more than 42.0 fluid ounces (0.42 lb ai) per acre per crop cycle.</li> </ul>
1.0 fl oz/acre .21 lb ai/acre)	<ul> <li>per crop cycle.</li> <li>Do not make more than 2 applications per crop cycle.</li> <li>Allow at least 14 days between applications.</li> <li>Do not make more than 4 applications per year.</li> <li>Preharvest Interval (PHI): 1 day</li> </ul>

©2022 Nichino America, Inc. Torac is a registered trademark of Nichino America, Inc. The Fujiichi-mark is a trademark of Nihon Nohyaku Co., Ltd.