



Section 24(c) Special Local Need Label

FLUAZIFOP-P-BUTYL

GROUP

1A

HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Fusilade® DX Herbicide
For Quackgrass Control in Alfalfa Grown for Seed

EPA Reg. No. 100-1070
EPA SLN No. OR-040006

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended.

Active Ingredient:

Fluazifop-P-butyl

Butyl (R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate*..... 24.5%

Other Ingredients:..... 75.5%

Total:..... 100.0%

Fusilade DX Herbicide formulation is an emulsifiable concentrate (EC).

*Fusilade DX Herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon
Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Environmental Hazards: Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product is toxic to fish and aquatic invertebrates. This product is toxic plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift and do not apply when weather conditions favor drift from target areas. Do not apply to areas where runoff into water bodies is expected. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Groundwater Advisory: Fluazifop-P-butyl has properties and characteristics associated with chemicals detected in groundwater and is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, including ponds, streams, and springs will reduce the potential loading of fluazifop-P-butyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Section 24(c) label and the main EPA-registered Fusilade DX Herbicide label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this 24(c) label and on the main EPA-registered Fusilade DX Herbicide product label.
- Failure to follow the Directions for Use and Precautions on this label may result in poor pest control, crop injury, or illegal residues.

Alfalfa (Grown for Seed): For control of quackgrass (*Agropyron repens*) in alfalfa grown for seed in Oregon, make two applications of Fusilade DX Herbicide following the application directions on the EPA-registered label.

WEED	APPLICATION	HEIGHT	RATE
Quackgrass	1st 2nd	6 to 10 inches up to 10 inches	12 fl oz/A 8 fl oz/A

Make the second application 2-3 weeks after the first, but before the quackgrass exceeds 10 inches in height. Always use crop oil concentrate at 1% v/v.

Restrictions/Precautions:

- Do not apply this product through any type of irrigation system.

SPECIAL CROP USE RESTRICTIONS

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food

or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

©2022 Syngenta

Fusilade® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: OR1070062EA1122