

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

BELAY INSECTICIDE

EPA Reg. No. 59639-150

BELAY® INSECTICIDE FOR USE ON LEAF-FOOTED BUG IN TREE NUTS AND SPOTTED WING DROSOPHILA ON FIG, PEACH, POMEGRANATE

This FIFRA Section 2(ee) recommendation expires December 31, 2025.

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Belay* Insecticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

RECOMMENDATIONS FOR THE CONTROL OF LEAFFOOTED BUG AND SPOTTED WING DROSOPHILA IN FIG, PEACH, POMEGRANATE AND TREE NUTS WITH FOLIAR APPLICATIONS

Pests	Crops	<i>Belay</i> Insecticide Application Rates	Special Instructions
Leaf-footed Bug	Fig Peach Pomegranate Tree Nuts	4 to 6 fl oz/A (0.067 to 0.1 lb ai/A)	Apply <i>Belay</i> Insecticide when target pest threshold populations are observed. Do not wait until population above threshold has been established. The amount of <i>Belay</i> Insecticide per acre will depend on tree size and pest pressure. Choose lower rates for light infestation and/or small trees and the higher rates for heavy infestations and/or larger trees.
Spotted Wing Drosophila	Fig Peach Pomegranate	4 to 6 fl oz/A (0.067 to 0.1 lb ai/A)	Apply by ground on mature trees using 100 to 400 gallons per acre as a full coverage spray. Non-bearing trees and small size trees can use below 100 gallons per acre. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.
<ul style="list-style-type: none">• Do not apply more than 12 fl oz (0.2 lb ai) of <i>Belay</i> Insecticide per acre per season.• Do not apply within 21 days of harvest on peach or tree nuts.• Do not apply within 7 days of harvest on fig or pomegranate.• Do not feed or allow livestock to graze on cover crops from treated orchards.• <i>Belay</i> Insecticide must not be applied during bloom and/or if bees are foraging.			

Refer to *Belay* Insecticide label for additional insects controlled.

Please contact Valent U.S.A. LLC at 1-800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2021 Valent U.S.A. LLC

Belay is a registered trademark of Sumitomo Chemical Company, Ltd.

Registrant: **Valent U.S.A. LLC**
P.O. Box 5075
San Ramon, CA 94583