

## FIFRA SECTION 24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF LOUISIANA



For the Control of Red Rot (*Glomerella tucumanensis*, *Colletotrichum falcatum*) and  
Pineapple Disease (*Ceratostyis paradoxa*) in Sugarcane

EPA Reg. No. 100-1324-5905  
EPA SLN No. LA20-0004

This label expires and must not be distributed or used in accordance with this SLN registration after  
May 20, 2025

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

### Restrictions:

- Avaris 2XS may be applied **either** as an In-Furrow Application, **or** a Conveyor Belt Treatment.
- Regardless of which application method used, DO NOT apply more than 21 fl oz/A of Quilt Xcel Fungicide at planting.

### In-Furrow Application

Apply Avaris 2XS to sugarcane seed pieces as an in-furrow spray at planting at the rate of 1.4 – 3.0 fl oz per 1000 row feet using sufficient water to ensure coverage. Mount the spray nozzle so the spray is directed into the furrow just before the seed cane is covered. Use the higher rate when the weather conditions are expected to be conducive for disease development.

### Conveyor Belt Treatment (Attached to Planting Equipment)

Treat seed pieces with Avaris 2XS spray solution at the rate of 1.4 – 3.0 fl oz per 1000 row feet, using in-line directed spray sufficient to wet cut ends.

### 24(c) Registrant:

Helena Agri-Enterprises, LLC  
225 Schilling Boulevard, Suite 300  
Collierville, TN 38017