

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



AGROXONE® 750 HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA present as the dimethylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For control of broadleaf weeds in Cereals, Linseed, Pastures, Rice, Sugar Cane and Turf as per the Directions for Use

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

CONTENTS: 20L, 110L, 1000L

APVMA Approval No: 60733/59739



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

© Agroxone is a registered trademark of A H Marks & Co UK

SIPCAM AGROXONE 750 HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes and will irritate the skin. When opening container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126). If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcarn's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcarn's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcarn or the place in which you purchased this product for a full refund.

Batch No:

Date of Manufacture:



This product is NOT classified as a Dangerous Good under the Australian code for the transport of dangerous goods by road and rail.

For specialist advice in an emergency only, call 1800 033 111, all hours Australia wide

G.H.S. STATEMENTS

PREVENTION P262: Do not get in eyes, on skin, or on clothing.

P270: Do not eat, drink or smoke when using this product.

RESPONSE P370+P378: Not combustible. Use extinguishing media suited to burning materials.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



AGROXONE® 750 HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA present as the dimethylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For control of broadleaf weeds in Cereals, Linseed, Pastures, Rice, Sugar Cane and Turf as per the Directions for Use

THIS LEAFLET IS PART OF THE LABEL

APVMA Approval No: 60733/59739



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

© Agroxone is a registered trademark of A H Marks & Co UK

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

FIELD CROPS

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Barley, Oats, Triticale, Wheat	Refer to Weeds Table	SA, Tas only	460 mL-1.45 L	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.
		Qld only		Apply from early tillering (4-5 fully emerged main stem leaves plus 1 or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).
		NSW, ACT only		Apply 460 mL rate after crop has reached 5 leaf stage and weeds marked # in Weeds Table have 3-6 leaves. Apply higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.
		Vic only	460-960 mL	Apply 460 mL rate after crop has reached 2-3 leaf stage and weeds marked # in the Weeds Table have 2-4 leaves. Apply higher rate after crop plants have 5 leaves to fully tillered.
		WA only	600 mL-1.35 L	Apply to crop from 5 leaf to flag leaf emergence (Zadoks 15-37).
	Skeleton Weed	SA only	660 mL plus 500 mL clopyralid (300 g/L)	Apply to rosettes from 5-15 cm diameter when the crop is between the five leaf and late tillering growth stages. It is essential that all the Skeleton Weed is emerged and a minimum of 5 cm diameter, which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.
Soldier Thistle		460 mL plus 200 mL clopyralid (300 g/L)	Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.	
Barley, Cereal Rye, Triticale, Wheat	Capeweed, Doublegee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	270 mL plus 350 mL diuron (500 g/L) or 330 mL plus 400 mL diuron (500 g/L)	Use the lower rate when crop is at the 3-4 leaf stage (Zadoks 13-14). Use the higher rate when crop is at the 4-5 leaf stage (Zadoks 14-15). Apply by aircraft or boom with 20-100 L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics. TANK MIXES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products. (Refer to Diuron label for complete list of restraints and directions for use)

DIRECTIONS FOR USE (continued)

FIELD CROPS

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Barley, Cereal Rye, Triticale, Wheat	Amsinckia (Yellow Burr), Capeweed, Charlock, Deadnettle, Doublegee (Spine Emex, Three Cornered Jack), Erodium sp, Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip	NSW, ACT, Vic only	330 mL plus 500 mL diuron (500 g/L)	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW, ACT, SA only) and 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT apply to undersown Medic or Lucerne. DO NOT use on Kamilarol, Kite, Olympic or Shortim Wheat. TANK MIXES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products. (Refer to Diuron label for complete list of restraints and directions for use)
	Capeweed, Mustard, Three Cornered Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel	SA only	230 mL plus 500 mL diuron (500 g/L)	DO NOT use on very light sandy soils. TANK MIXES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products. (Refer to Diuron label for complete list of restraints and directions for use)
Flax	Refer to Weeds Table	Vic, Tas only	325 mL	Apply to crop when 8-20 cm high.
Linseed		Qld only	460 mL-1.45 L	Apply when crop is 10-15 cm tall with at least 170 L/ha of water. DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft.
		NSW, ACT, Vic, SA, Tas only	460-730 mL	
Rice	Buddah (Butter) Pea, Dirty Dora, Sedges, Starfruit	NSW only	485-970 mL	Apply by aircraft at early-mid tillering stage of rice, but not later than panicle initiation. Use 485 mL/ha at 35 days after aerial sowing (approximately 2 L/rice plant). Use 970 L/ha at 45 days or more after cereal sowing (3 tillers or more/rice plant). The crop should be sprayed before flowering of the weeds. Water management: Before herbicide application, lower water level in the bays to at least 2/3 of the weed growth for direct contact with SIPCAM AGROXONE 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying.

DIRECTIONS FOR USE (continued)

FIELD CROPS

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	Qld only	930 mL	Post emergent: Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties.
	As above plus Fat Hen, Noogoora Burr		1.45 L	

PASTURES, SEED CROPS AND TURF

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Grass Pastures	Refer to Weeds Table	All States	460 mL-2.7 L	Apply to established grass based pastures. Damage may occur to Legumes if present. See Legume Tolerance under General Instructions. Note: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying.
	Horehound (seedling and established)	NSW, ACT only	2 L	Apply in autumn when Horehound plants are at the seedling stage. If Sub-Clover or Medic plants are present they will be damaged by this application.
	Solder Thistle	SA only	460 mL-1.45 L	Use low rate on seedling with no more than 2 true leaves. Use 100 L water for best results. DO NOT exceed 575 mL if Clover is present.
Grass Seed Crops	Refer to Weeds Table	Qld, SA, Tas, Vic, WA only	460 mL-2.7 L	Apply to established grass seed crops.

DIRECTIONS FOR USE (continued)

PASTURES, SEED CROPS AND TURF

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Pastures Spray/graze technique	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, Vic only	230-930 mL	<p>Spray to actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep, as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. See Legume Tolerance under General Instructions.</p> <p>Precaution: An increased quantity of poisonous plants may be eaten by stock using spray/graze, eg Caltrop, Capeweed, Paterson's Curse and Variegated Thistle, and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended period of grazing. Avoid grazing with young or breeding stock.</p> <p>DO NOT graze horses or pigs on Paterson's Curse. Observe withholding period of at least 7 days after spraying.</p>
		Tas only	460 mL	
	Refer to Weeds Table	WA only	610 mL-1.25 L	
	Capeweed, Dandelion, Seedling Dock, Erodium, Geranium, Mustard, Annual Thistles, Three Cornered Jack, Turnip Weed, Wild Radish, Wild Turnip	SA only	330 mL	
Soldier Thistle	395 mL			
Sub-Clover Seed Crops	Refer to Weeds Table	Vic, SA only	460-960 mL	DO NOT exceed rate specified. Spray in autumn and early winter.
Turf		All States	330 mL or 2.8 mL/100 m ²	Newly laid: Delay until well established or at least 2 months old. Apply in high volume to actively growing weeds. See Legume Tolerance under General Instructions. Buffalo Grass is more easily damaged than other grasses.
	930 mL-1.8 L or 10-19 mL/100 m ²		Established: Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and Strawberry Clover. See Legume Tolerance under General Instructions. Buffalo Grass is more easily damaged than other grasses.	

DIRECTIONS FOR USE (continued)

SPOT SPRAYING

Crop / Situation	Weeds	State	Rate	Mixing Rates / Comments
High volume spraying	Refer to Weeds Table	All States	-	Add 1/10 th of rate in Weeds Table to 150 L of water. Each 150 L of mix will cover 1,000 m ² (1/10 th ha). Eg If rate in Weeds Table is 1.4 L, use 140 mL/150 L water.
Knapsack application				Add 1/100 th of rate in Weeds Table to 10 L of water. Each 10 L of mix will cover 100 m ² (1/100 th ha). Eg If rate in Weeds Table is 1 L, use 10 mL/10 L water.

NON-CROP AREAS, FALLOW LAND, INDUSTRIAL AND COMMERCIAL AREAS, RIGHTS-OF-WAY

Crop / Situation	Weeds	State	Rate/ha	Critical Comments
Non-crop areas, Fallow land, Industrial and Commercial areas, Rights-of-Way	Refer to Weeds Table	All States	Refer to Weeds Table	Apply at rates shown in Weeds Table when weeds only are present, and non-crop or pasture is involved.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

CEREALS, PASTURE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

WEEDS TABLE

Where weeds are to be sprayed in a crop or pasture situation, use only the rate given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however, some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weeds Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

Weeds	Application Rate/ha						Critical Comments
	Vic	NSW, ACT	SA	Tas	Qld	WA	
Ball Mustard	-	-	1.15 L	-	-	-	Apply at rosette stage.
Bathurst Burr	1.35 L	660 mL-1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	Spray young seedlings only.
Black Bindweed	-	970 mL-1.35 L	-	-	930 mL-1.35 L	-	Apply at young stage prior to flowering
California Burr		660 mL-1.35 L		1.45 L			Spray young seedlings only.
Cape Tulip		-		-		1.35 L	Apply just prior to flowering. Gives mild suppression only.

DIRECTIONS FOR USE (continued)

WEEDS TABLE

Weeds	Application Rate/ha						Critical Comments
	Vic	NSW, ACT	SA	Tas	Qld	WA	
Capeweed	-	1.45 L	1.45 L	1.45 L	1.45 L	-	Apply at seedling stage.
Carrot Weed	1.35 L	-	-	-	1.35 L	-	
Charlock	460 mL	660 mL	460 mL	660 mL- 1.45 L	460 mL	-	Spray up to rosette stage.
Cutleaf Mignonette	-	-	1.45 L	-	-	-	Apply at young rosette stage.
Dandelion	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	
Deadnettle	-	1.45 L	-	-	1.45 L	-	Moderately susceptible. Apply at seedling stage.
Docks	960 mL	-	1.45 L	1.45 L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	1.35 L	660 mL- 1.35 L	-	1.35 L	1.35 L	-	Spray up to rosette stage.
Fennel	1.8 L	1.8 L	1.8 L	1.8 L	1.8 L	-	Apply at young stage.
Field Bindweed	1.35 L	-	1.35 L	1.35 L	1.35 L	1.35 L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	1.45 L	-	-	Apply at young stage.
Flatweed or Catsear	-	-	-	-	-	930 mL	Apply when weeds are 10-15 cm high.
Fumitory, Red	930 mL	930 mL	930 mL	-	930 mL	-	Apply at young stage prior to flowering.
Hoary Cress	3 L	-	1.45 L	-	3 L	-	Apply before flowering.
Horehound	-	2 L	-	-	-	-	Apply at rosette stage.
Hedge Mustard	730 mL#	-	460 mL	460 mL	460 mL	825 mL	Apply at rosette stage.
Hexham Scent or Melilotus	960 mL	800 mL	930 mL	-	930 mL	-	Apply at young seedling stage.
Lesser Swinecress	-	-	-	1.45 L	-	-	Apply at young stage.
Lincoln Weed	-	-	460 mL	-	-	-	Apply at rosette stage.
London Rocket	-	-	-	-	-	825 mL	-
Lupins	-	460-960 mL	-	-	-	-	Spray up to 10 cm high.
Mintweed	1.35 L	1.35 L	-	-	1.35 L	-	Spray young seedlings only.
Mustards	730 mL#	660 mL#	730 mL	730 mL	-	825 mL	Spray up to rosette stage.
Noogoora Burr	1.35 mL	660mL- 1.35 L	1.35 L	-	1.35 L	1.35 L	Spray young seedlings only.
Opium Poppy	-	-	-	1.45 L	-	-	Apply at young stage.
Paterson's Curse	-	660-960 mL	-	-	2.7 L	-	Apply at early rosette stage.
Pimpernel	-	-	460 mL	-	-	-	Apply at rosette stage.
Plantains	-	1.35 L	-	1.35 L	1.35 L	-	Apply at young stage.

DIRECTIONS FOR USE (continued)

WEEDS TABLE

Weeds	Application Rate/ha						Critical Comments	
	Vic	NSW, ACT	SA	Tas	Qld	WA		
Rapeseed	-	660-960 mL	-	-	-	-	Spray up to rosette stage.	
Rough Poppy		460-960 mL						
Safflower		460-660 mL						
Shepherd's Purse		-		1.45 L				Apply at rosette stage.
Skeleton Weed		960mL – 1.35 L		-				Spray rosettes before aerial growth commences.
Sorrel		-	1.45 L					Apply up to rosette stage.
Stinkwort	1.35 L	660-960 mL	1.35 L			1.35 L	Apply when weeds are 10-15 cm high.	
Sunflower seedlings	-	690 mL-1 L	-			-	Spray up to rosette stage.	
Thistles								
Artichoke	1.05 L	-	1.45 L		1.35 L	460 mL	Apply at rosette stage.	
Nodding	-	1.65 L	-		-	-	Spray early rosettes. Repeat spray required in following spring or autumn.	
Saffron	960 mL-1.65 L	660 mL-1.35 L	930 mL-1.65 L	930 mL-1.65 L	930 mL-1.65 L	930 mL-1.65 L	Spray up to rosette stage. WA: September .	
Scotch or Cotton	1.45 L	-	460-930 mL	1.45 L	-	-	Spray young seedlings only.	
Slender	2 L	660 mL-1.35 L	660 mL	2 L		2 L		
Soldier	730 mL	-	730 mL	-		-	Spray young seedlings. Use higher rate for larger weeds.	
Spear	-	960 mL-1.35 L	-	1.45 L			Apply at rosette stage.	
Star		-	730-930 mL	-			Apply at seeding stage. Use higher rate for larger weeds.	
Thistles								
Variegated	960 mL	660 mL-1.35 L	460-930 mL	960 mL	960 mL	960 mL	Spray young seedlings only. SA: Use 1.35 L/ha for well developed weeds.	
Turnip Weed	730 mL#	660 mL#	660 mL	-	-	-	Spray up to rosette stage.	
Wild Artichoke	-	-	1.45 L				Apply at rosette stage.	
Wild Radish	930 mL	660 mL#	930 mL	1.35 L	460-730 mL	800 mL	Spray up to rosette stage.	
Wild Turnip	730 mL		460 mL			600 mL		

Refer to Critical Comments for Barley, Oats, Triticale, Wheat in the Field Crops table

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application. DO NOT spray crops or weeds outside the stages indicated in the Critical Comments, as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

LEGUME TOLERANCE

The use of MCPA is preferred to 2,4-D amine (amicide) or ester (estercide) in situations where Legumes are undersown in the crop.

Note: The tolerance of pasture Legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with Lucerne as damage may result.

MIXING

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

APPLICATION

Boom spray: Use 30-120 L water/ha. Use minimum of 170 L water/ha for Linseed.

Aerial spray: Use 10-90 L water/ha.

EQUIPMENT

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY

This product can be tank mixed with clopyralid, dicamba, diquat, diuron, metsulfuron-methyl, paraquat, picloram, sethoxydim and terbutyrn. The effectiveness of dimethoate, oxydemeton methyl and phosmet and may be reduced by alkaline hydrolysis if tank mixed with the product. For these situations the addition of a surfactant (350 g/L soyal phospholipids + 350 g/L propionic acid) is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

RESISTANT WEEDS WARNING

GROUP I HERBICIDE

SIPCAM AGROXONE 750 Herbicide is a member of the phenoxy group of herbicides. SIPCAM AGROXONE 750 Herbicide has the disruptor of plant cell growth mode of action. For weed resistance management SIPCAM AGROXONE 750 Herbicide is a Group I herbicide. Some naturally occurring weed biotypes resistant to SIPCAM AGROXONE 750 Herbicide and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SIPCAM AGROXONE 750 Herbicide or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcarn Pacific Australia Pty Limited accepts no liability for any losses that may result from the failure of SIPCAM AGROXONE 750 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT allow spray drift and vapour movement onto susceptible crops such as cotton, fruit trees, legume crops, lucerne, oilseed crops, pastures, tomatoes, vegetables, vines or other sensitive crops and trees (eg Kurrajongs, Belahs, Eucalypts).

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

RE-ENTRY PERIOD

Hand weeding should not be performed for 7 days after spraying application unless workers wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit

specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes and will irritate the skin. When opening container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126). If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.