

FIFRA Section 24(c) Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Curtail[®] M

EPA Reg. No. 62719-86

24(c) Special Local Need Registration SLN ND-090003

Control of Broadleaf Weeds in Flax

(For Distribution and Use Only in the State of North Dakota)

Expiration Date: December 31, 2020

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Curtail[®] M herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Curtail M according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Curtail M herbicide.

Directions for Use

Refer to product label for a complete list of weeds controlled, General Use Precautions, Mixing and Application instructions.

Apply 1 1/3 to 1 3/4 pints per acre of Curtail M when flax is 2 to 6 inches tall and target weeds are actively growing. The higher rate may be used when the condition of the weeds and/or crop at the time of treatment may prevent optimum control. Follow the application directions for ground or aerial application on the product label for Curtail M, including advisories for avoiding spray drift. To control or suppress weeds listed on the label, apply after maximum emergence of the target weeds but before they exceed 3 inches in height or diameter (for rosettes). To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged from the soil (plants 4 to 6 inches in height) up to bud stage.

Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds that have emerged at the time of application will be affected.** If foliage is wet at the time of application, control may be decreased. Curtail M is rainfast within 6 hours after application.

Tank Mixtures: Curtail M may be used in combination with other herbicides that are labeled for flax. Refer to the label of the tank mix partner for lists of other weeds controlled, rates of application, and use precautions.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 72 days of harvest.
- Do not apply after flax has begun bolting as crop injury may occur if applied during the bloom period.
- Do not apply more than 1 3/4 pint per acre or make more than 1 application per crop season.
- Carefully follow rotational crop restrictions and other use precautions and limitations on the product label for Curtail M.

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R123-003

Accepted: 12/03/15

Replaces R123-002