



Section 2(ee) Recommendation

Date: August 21, 2012

Product: Daconil Action™
EPA Reg. No. 100-1364

Use: Daconil Action for the suppression of Pythium Blight and Bacterial Wilt on turfgrass

State: CA

DIRECTIONS FOR USE

Daconil Action fungicide couples a multiple-site contact fungicide with a systemic compound used for control and suppression of disease through induction of host plant resistance. Daconil Action exhibits an additional unique mode of action, beyond chlorothalonil alone, which mimics the natural systemic acquired resistance (SAR) response found in most plant species. This SAR response does not involve direct activity against the target pathogens.

This recommendation applies to both Group A Turf and Group B Turf as listed on the federal label.

Daconil Action provides suppression of the following diseases on turf.

Target Disease	Use Rate (fl. oz. product per 1000 sq. ft.)	Use Rate (Pints Product/Acre	Application Interval (days)	Instructions
Pythium Blight	2.0 2.0 – 3.5	5.4 5.4 -9.6	7-10 7 to 14	For optimum suppression of Pythium blight, should be applied prior to the appearance of disease. Follow all directions for use, restrictions and precautions as given on the federal label.
Bacterial Wilt (<i>Xanthomonas</i> spp). (<i>Acidovorax</i> spp.)	3.5	9.6	14	For optimum suppression of Bacterial Wilt, should be applied prior to the appearance of disease. Follow all directions for use, restrictions and precautions as given on the federal label.

The user must refer to the federally approved label and read and follow all directions for use, restrictions, and precautions. The user should have this recommendation in their possession at the time of use. This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the EPA.

Daconil Action™ and the Syngenta logo are trademarks of a Syngenta Group Company.

©2012 Syngenta Crop Protection, LLC,
410 Swing Road, Greensboro, NC 27409

CA1364163AC0812