

Maxi-Boost®

Guaranteed Analysis:

Magnesium (Mg)	1.05%	Manganese (Mn)	0.25%
Sulphur (S)	1.38%	0.25% Chelated Manganese	
Boron (B)	0.25%	Molybdenum (Mo)	0.004%
Copper (Cu)	0.25%	Zinc (Zn)	0.14%
0.25% Chelated Copper		0.14% Chelated Zinc	
Iron (Fe)	0.5%		
0.5% Chelated Iron			

Derived from iron-copper-zinc-manganese EDTA, boric acid, sodium molybdate, and magnesium sulphate

For use as a fertilizer in fertigation applications to promote root growth and supplement nutrient levels on agricultural crops.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Directions for Use (Use only as Directed):

- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality. Not compatible with aluminum equipment.
- Maxi-Boost® is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

Crop	Fertigation Rates*
Forage Crops: Alfalfa, Clover, Hay Crops Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane Legumes: Dry bean, Lentil, Peanut, Pea, Soybean Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince Other Fruit: Citrus (all), Fig, Kiwi, Melons Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut Grapes: Currant, Raisin, Table, Wine Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	<p>The number and frequency of applications determine application rates.</p> <p>Apply no more than 1 quart/acre/application.</p> <p>Apply no more than 6 quarts/season.</p> <p>Always apply toward the end of an irrigation cycle.</p>

*Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials after the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: The use of the fertilizing material on any crops other than those recommended may result in serious injury to the crops.

CAUTION: The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

Caution: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush skin or eyes with water for at least 15 minutes.

A True Solution containing Micronutrients and Seaweed

Net Contents:
2.5 Gal. (9.46 L)

Net Weight:
25.39 lbs. (11.54 kg)

Net Weight/Gallon:
10.16 lbs. (4.62 kg)

Batch No.: XXXXXX
Rev: 091214