

# PhitaK 0-46\*-11

## Guaranteed Analysis:

|  |     |
|--|-----|
| Total Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) | 46% |
| Soluble Potash (K <sub>2</sub> O)                      | 11% |

Derived from potassium phosphite

**For use as a fertilizer in foliar and fertigation applications to supplement phosphorous and potassium levels on agricultural crops.**

\*Phosphorous acid products are for use as a supplemental fertilizer treatment.

\*Upon foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or will be used directly by plants, as phosphite ions.

\*As a soil application to annual crops, a lesser response from the initial crop, with a corresponding superior response for succeeding crops, may be observed. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

## Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- PhitaK is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- **Do not freeze. Store above 50°F(10°C)**
- **Consult a NutriAg representative for application timing and necessity of its use.**

## Suggested Application Rates:

| Crop  | Foliar Rates*                  | Fertigation Rates*  |
|---|--------------------------------|---|
| <b>Forage Crops:</b> Alfalfa, Clover, Hay Crops<br><b>Field Crops:</b> Cereals, Corn, Cotton, Sorghum, Sugar cane<br><b>Legumes:</b> Dry bean, Lentil, Peanut, Pea, Soybean<br><b>Root Vegetables (including, but not limited to):</b> Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip<br><b>Leafy Vegetables (including but not limited to):</b> Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach<br><b>Other Vegetables:</b> Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato<br><b>Berry Crops:</b> Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry<br><b>Tree Fruit (including but not limited to):</b> Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince<br><b>Other Fruit:</b> Citrus (all), Fig, Kiwi, Melons<br><b>Nut Trees (including but not limited to):</b> Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut<br><b>Grapes:</b> Currant, Raisin, Table, Wine<br><b>Other Crops:</b> Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower | 1-4 pints/acre/<br>application | The number and frequency of applications determine application rates.<br><br>Apply no more than 3-4 quarts/acre/ application.<br><br>Apply no more than 6 quarts/season.<br><br>Always apply toward the end of an irrigation cycle. |

\*Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

**Warranty Statement:** Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

**Caution: KEEP OUT OF REACH OF CHILDREN**

## A Dual-Action True Solution for Phosphorous and Potassium Nutrition



### Danger

Causes serious eye damage. Contains 34-37% of ingredients of unknown toxicity (inhalation). Contains 27-29% of ingredients of unknown toxicity (dermal).

Wear eye protection/face protection.

If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a poison center/doctor

Dispose of contents/container in accordance with local, regional, national and international regulations.

## Net Contents:

2x2.5 Gal. (2x9.46 L)

## Net Weight:

62.30 lbs. (28.26kg)

## Net Weight/Gallon:

12.46 lbs. (5.65 kg)

## Batch No.: 260115

Rev: 300914