

Guaranteed Analysis:

Boron (B) 0.5%
Zinc (Zn) 10.2%

A Zinc and Boron True Solution with PAC*Chelation Technology

Derived from zinc oxide and boric acid

For use as a fertilizer in foliar, fertigation and soil applications to supplement zinc and boron levels on agricultural crops. *PAC refers to NutriAq's "PolyAldoCarbosate" Technology

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground sprays minimum of 5 gallons of water/acre: Dilute sprays minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- ZincMax is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

Сгор	Foliar Rates**	Fertigation Rates**	Soil Rates**
Forage Crops: Alfalfa, Clover, Hay Crops		-	This product is
Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane		The number and	compatible with
Legumes: Dry bean, Lentil, Peanut, Pea, Soybean		frequency of	many liquid
Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip,		applications determine	fertilizers.
Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip		application rates.	
Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro,			Fertilizers vary in
Cole Crops, Lettuce, Parsley, Spinach		Apply no more than	analysis and
Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra,	1-4 pints/acre/	2 quarts/acre/	quality, and
Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato		application.	should be jar
Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry	application		tested with this
Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry,		Apply no more than	product before
Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince		8 quarts/season.	use.
Other Fruit: Citrus (all), Fig, Kiwi, Melons		•	
Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan,		Always apply toward	Apply up to
Pistachio, Walnut		the end of an irrigation	2 quarts/acre at
Grapes: Currant, Raisin, Table, Wine		cycle.	planting or as a
Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower		-	side dress.

^{**}Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fif for purposes stated on such label only who used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

Caution: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush skin or eyes with water for at least 15 minutes.

Net Contents:

2.5 Gal (9.46 L)

Net Weight:

26.08 lbs. (11.83 kg)

Net Weight/Gallon: 10.43 lbs. (4.73 kg)

Batch No.: 161111