

QUALI-PRO[®]

Glyphosate T&O

Herbicide

ACTIVE INGREDIENT:	% BY WT.
*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt	41.0%
OTHER INGREDIENTS:	59.0%
TOTAL	100.0%

*This product contains 480 grams per liter or four pounds per U.S. gallon of glyphosate—the active ingredient—in the form of its isopropylamine salt. This equals 356 grams per liter, or three pounds per U.S. gallon of the acid, glyphosate.

EPA Reg. No. 66222-208

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Immediately hold eyelids open and flush with plenty of water for at least 15 minutes.• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.• Get medical attention.
IF INHALED:	<ul style="list-style-type: none">• Remove individual to fresh air.• If not breathing, give artificial respiration, preferably mouth-to-mouth.• Get medical attention.
IF SWALLOWED:	<ul style="list-style-type: none">• This product will cause gastrointestinal tract irritation.• Immediately dilute by swallowing water or milk.• Get medical attention.• Never give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Prosar at 1-877-250-9291 for emergency medical treatment information.	

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING! CAUSE SUBSTANTIAL BUT TEMPORARY EYE INJURY. HARMFUL IF SWALLOWED OR INHALED. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

When using spray solutions of this product, mix, store, and apply only in stainless steel, fiberglass, aluminum, plastic, or plastic-lined steel containers.

DO NOT USE GALVANIZED STEEL OR UNLINED STEEL (EXCEPT FOR STAINLESS STEEL) CONTAINERS OR SPRAY TANKS WHEN SPRAYING, MIXING, STORING, OR APPLYING THIS PRODUCT. Quali-Pro Glyphosate T&O, or spray solutions with this product, react with these containers to produce hydrogen gas, a possibly very highly-combustible gas mixture that could explode or flash, thereby causing serious personal injury if ignited by spark, welder's torch, lighted cigarette, open flame, or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

GENERAL INFORMATION

DO NOT ALLOW CONTACT WITH EXPOSED NON-WOODY ROOTS, FRUIT, FOLIAGE OR GREEN STEMS OF DESIRABLE TREES AND PLANTS, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

Use only in accordance with instructions on this label. Read the entire label before use, including the **LIMITED WARRANTY, TERMS OF SALE, AND LIMITATIONS OF LIABILITY** section. If the terms in this section are not acceptable, immediately return unopened product.

QUALI-PRO GLYPHOSATE T&O IS AN END-USE PRODUCT AND IS NOT REGISTERED FOR REFORMULATION.

DO NOT APPLY QUALI-PRO GLYPHOSATE T&O BY AIR, EXCEPT UNDER SPECIFIC CONDITIONS LISTED WITHIN THIS LABEL.

Quali-Pro Glyphosate T&O is a water-soluble liquid that readily mixes with water for application as a foliar spray to control or destroy most herbaceous plants. This product can be applied by using most standard industrial or field-type sprayers after it is diluted and thoroughly mixed with water.

Quali-Pro Glyphosate T&O works by moving through the plant from where it contacts the foliage to and into the root system. Within 2 to 4 days, visible effects on most annual weeds are apparent; however, effects may not be visible on most perennial weeds for 7 days or more. Effectiveness and visual effects of this product may be slower if very cool or cloudy weather follows treatment. A gradual wilting and yellowing of the plant will be visible, which then advances to total browning of all above-ground growth and deterioration of underground plant parts.

Unless stated otherwise within this label, application should be delayed until vegetation has emerged and has reached the specific stages described for control of vegetation under the **WEEDS CONTROLLED** sections of this label. Unemerged plants that arise from unattached underground rhizomes or root stocks of perennials are not affected by Quali-Pro Glyphosate T&O, and will keep growing. Because of this, treatment should be made at late growth stages (approaching maturity) for best control of the majority of perennial weeds.

Always apply the higher rate of this product per acre within recommended ranges when weed growth is heavy or dense, or when weeds are growing in a noncultivated, nondisturbed area.

Reduced weed control may result if weeds are treated when drought stress, insect damage, or disease is present. Additionally, if weeds are heavily covered with dust, reduced weed control will result.

Effectiveness may be reduced if applications are made to annual or perennial weeds that have been grazed, mowed or otherwise cut, unless they have been permitted to regrow to their recommended stages for treatment.

If rainfall or irrigation occurs within 6 hours of application, reduced effectiveness may result. Heavy irrigation or rainfall within 2 hours of application may wash off the chemical from foliage. In this case, a repeat treatment may be required.

Quali-Pro Glyphosate T&O does not provide residual weed control. Use a label-approved herbicide program if subsequent residual weed control is desired. Users must read and carefully observe all cautionary statements and all other information on labels of all herbicides used.

NOTE: Buyer and users are responsible for all losses or damage resulting from the use and/or handling of mixtures of Quali-Pro Glyphosate T&O with herbicides or other materials that are not specifically recommended on this product label. Reduced performance may result if users mix Quali-Pro Glyphosate T&O with other herbicides or other products not recommended on this label.

Best results are obtained with uniform and complete spray coverage. However, do not spray weed foliage to the extent of runoff.

DOMESTIC ANIMALS: Quali-Pro Glyphosate T&O is considered to be relatively nontoxic to dogs and other domestic animals. Note,

however, that ingestion of this product, or of large amounts of vegetation that has been freshly sprayed with this product, may cause temporary gastrointestinal irritation, such as diarrhea, colic, vomiting, etc. If these symptoms are observed, the animal should be given plenty of fluids in order to prevent dehydration. A veterinarian should be contacted in the event symptoms persist for more than 24 hours.

NOTE: Using this product in a manner not consistent with the product label may result in personal injury, injury to animals or crops, or in other unintended consequences.

MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT WITH THE CAPACITY TO DELIVER THE DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. DIRECT HAND-GUN APPLICATIONS PROPERLY TO AVOID SPRAYING DESIRABLE PLANTS. **NOTE:** POOR RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

For mixing, additives, and application instructions for Industrial, Turf, and Ornamental uses, see the section titled **NONCROP USES**.

MIXING

Quali-Pro Glyphosate T&O mixes readily with water. Spray solutions of this product should be mixed in accordance with the following: Fill the mixing/spray tank with the correct amount of water. Then add the recommended amount of Quali-Pro Glyphosate T&O (per the **DIRECTIONS FOR USE** and **WEEDS CONTROLLED** sections) near the end of the filling process. Mix well. Be careful to avoid back siphoning. If required by state or local regulations, use approved anti-back-siphoning devices. When mixing and applying this product, foaming of the spray solution can occur. To minimize or prevent foam, do the following: Do not use mechanical agitators; terminate by-pass and return lines at tank bottom, and, if necessary, use an approved defoaming or anti-foam agent.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of Quali-Pro Glyphosate T&O with water carriers by first mixing small, proportional quantities in advance.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Mix labeled tank mixtures of Quali-Pro Glyphosate T&O with water as described below:

1. Place a wetting basket or a 20- to 35-mesh screen over filling port.
2. Fill through the screen, adding one half the total amount of water,

then start agitation.

3. If using ammonium sulfate, add it slowly through the screen into the tank. Continue agitation. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before the addition of other products.
4. If using a wettable powder, first make a slurry with the water carrier, and SLOWLY add it through the screen into tank. Continue agitating.
5. If using a flowable formulation, premix one part flowable with one part water. SLOWLY add diluted mixture through screen into tank. Continue agitating.
6. If using an emulsifiable concentrate formulation, premix two parts water with one part emulsifiable concentrate. Then slowly add diluted mixture through the screen into tank. Continue agitating.
7. Continue filling spray tank with water and add required amount of Quali-Pro Glyphosate T&O when approaching the end of the filling process.
8. If nonionic surfactant is recommended, add it to the spray tank before ending the filling process.
9. Individual formulations are added to the spray tank in the following order: Wettable powder; flowable; emulsifiable concentrate; drift control additive; water-soluble liquid (e.g., Quali-Pro Glyphosate T&O) followed by surfactant.

Maintain agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation must be used to resuspend the mixture before spraying is resumed.

To minimize foaming, keep the bypass line on or near the bottom of the tank. The screen size used in line or nozzle strainers should be no finer than 50 mesh. Avoid spraying a fine mist by carefully selecting the proper nozzle. For best results when using conventional ground application equipment, flat fan nozzles should be used.

Clean sprayer and parts immediately when finished using this product by thoroughly flushing with water.

ADDITIVES

SURFACTANTS

Use only nonionic surfactants approved for use with herbicides. Do not reduce rates of Quali-Pro Glyphosate T&O when adding surfactant. Use 0.5-percent surfactant concentration (which is 2 quarts per 100 gallons of spray solution) when using those surfactants containing a minimum of 70 percent active ingredient, or a 1-percent surfactant concentration (which is 4 quarts per 100 gallons of spray solution) for those surfactants that contain less than 70 percent active ingredient. Read and carefully observe cautionary statements and all other information that is listed on the surfactant label.

AMMONIUM SULFATE

Adding 1 to 2 percent dry ammonium sulfate by weight (8.5 to 17 pounds per 100 gallons of water) may improve the performance of Quali-Pro Glyphosate T&O, particularly under hard water conditions, drought conditions, or when tank mixing with certain residual herbicides, on perennial and annual weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Dry

ammonium sulfate must be completely dissolved in the spray tank before adding herbicides or surfactants. Ensure the spray system is thoroughly rinsed with clean water after use to reduce corrosion. Adding the same amount of dry ammonium sulfate may also increase the performance of Quali-Pro Glyphosate T&O plus 2,4-D Banvel™ or residual herbicide tank mixtures on perennial and annual weeds. If environmental stress is present, performance improvement may be more apparent.

Please note that nozzle tip plugging may result if low quality ammonium sulfate (that contains material that cannot readily dissolve) is used. To determine the quality of dry ammonium sulfate, conduct a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water, then agitate for 1 minute. If the test leaves some sediment undissolved, predissolve the ammonium sulfate in water, then filter prior to adding it to the spray tank. If adding ammonium sulfate directly to the spray tank, add it slowly with agitation. Adding it too quickly could block the outlet line. Ammonium sulfate must be completely dissolved in the spray tank before adding surfactant or herbicides. Users should thoroughly rinse the spray system with clean water when spraying is completed to reduce corrosion.

NOTE: Using ammonium sulfate as an additive does not preclude any requirements for additional surfactant. Do not use herbicide rates that are lower than those recommended on this label.

COLORANTS OR DYES

Colorants or marking dyes that are agriculturally approved may be added to Quali-Pro Glyphosate T&O. However, dyes or colorants used in spray solutions of Quali-Pro Glyphosate T&O may reduce product effectiveness, especially when used at lower rates or dilutions. Follow all manufacturer's recommendations when using colorants or dyes.

DRIFT CONTROL ADDITIVES

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars, and Controlled Droplet Applicator (CDA) equipment. When using a drift control additive, read and carefully observe all cautionary statements and all other information appearing on the additive label. Using drift control additives can affect spray coverage which may result in reduced performance.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

The following application equipment may be used when applying this product.

By Air: Helicopter and Fixed Wing.

By Broadcast Spray:

Controlled-Droplet Applicator (CDA)—Boom-mounted or hand-held applicators that produce spray with a narrow range of droplet sizes.

Hand-Held and High-Volume Spray Equipment—Pump up pressure sprayers, handguns, mistblowers*, hand-wands, knapsack and backpack sprayers, lances and

other hand-held and motorized spray equipment used to direct the spray directly onto the weed foliage.

Ground Broadcast Spray—Boom or boomless systems, pull-type sprayer, pick-up sprayers, floaters, spray coupes, and other ground broadcast equipment.

Injection Systems—Aerial or ground injection sprayers.

- * This product is not registered in Arizona or California for application with mistblowers.

Selective Equipment—Wiper applicators, shielded sprayers, and recirculating sprayers.

Review the **Selective Equipment** section of the discussion below for rates of application and specific instructions.

SPRAY DRIFT MANAGEMENT

IMPORTANT: AVOID SPRAY DRIFT. EXERCISE EXTREME CAUTION WHEN APPLYING QUALI-PRO GLYPHOSATE T&O TO AVOID INJURY TO DESIRABLE CROPS AND PLANTS.

The herbicide solution must not be permitted to drip, splash, mist, or drift onto desirable vegetation because very small quantities of this product can severely damage or destroy crops, plants, or other planted areas that were not intended for treatment.

Avoiding spray drift is the responsibility of the applicator. The potential for spray drift is determined by the interaction of many equipment- and weather-related factors. All applicators and growers must consider all of these factors when making decisions regarding product application.

AERIAL EQUIPMENT

Unless otherwise specified on this label, use Quali-Pro Glyphosate T&O in 3 to 15 gallons of water per acre. For specific Quali-Pro Glyphosate T&O rates, review the **WEEDS CONTROLLED** sections of this label. Do not exceed 1 quart Quali-Pro Glyphosate T&O per acre unless otherwise specified. **Aerial applications of Quali-Pro Glyphosate T&O may be made in preharvest applications, fallow and reduced tillage systems, and annual cropping conventional tillage systems.** For recommended volumes and application rates, review the individual use area sections of this label. **FOR AERIAL APPLICATION IN ARKANSAS, CALIFORNIA, OR FRESNO COUNTY, CALIFORNIA, REFER TO THE SUPPLEMENTAL USES SECTION AT THE END OF THIS LABEL FOR SPECIFIC RESTRICTIONS, INSTRUCTIONS, AND REQUIREMENTS.**

Quali-Pro Glyphosate T&O plus dicamba tank mixtures may not be applied by air in California.

AERIAL SPRAY DRIFT MANAGEMENT

To avoid off-target drift movement, the following drift management requirements must be followed during aerial applications.

1. The distance of the outermost nozzles on the boom must not be more than 3/4 the length of the rotor or the wingspan.
2. Nozzles must always be pointed backward parallel with the air stream and must never be pointed downward at more than 45

degrees. Some states have more stringent regulations that must be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the **Wind, Temperature and Humidity**, and **Temperature Inversion** sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom Length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application Height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment

to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water.

Drift-control additives may be used. When a drift-control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application – To avoid streaked, uneven, or overlapped application, use appropriate marking devices.

After each day of spraying, aircraft must be thoroughly washed—especially the landing gear—to remove product residues that have accumulated from spraying or from spills. UNCOATED STEEL SURFACES ARE SUSCEPTIBLE TO CORROSION FROM PROLONGED EXPOSURE TO THIS PRODUCT. FAILURE OF THE PART DUE TO CORROSION IS ALSO POSSIBLE. LANDING GEAR ARE MOST SUSCEPTIBLE. The application and maintenance of an organic coating of paint that meets aerospace specification MIL-C-38413 may help in preventing corrosion.

NOTE: Aerial application of Quali-Pro Glyphosate T&O plus Oust®, 2,4-D, or Banvel tank mixtures are not permitted in California. For tank mixes in all other states, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

BROADCAST EQUIPMENT

For control of those perennial or annual weeds listed on this label, using broadcast equipment. Unless otherwise specified on this label, use the recommended rates of Quali-Pro Glyphosate T&O in 3 to 40 gallons of water per acre. Review the **WEEDS CONTROLLED** sections of this label for specific rate information. As weed density increases, spray volume should increase (within

the recommended range) to ensure adequate coverage. Avoid spraying a fine mist by carefully selecting the proper nozzle. Use flat fan nozzles for best results when using ground application equipment. Apply with an even distribution of spray droplets.

INJECTION SYSTEMS

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically recommended.

CONTROLLED DROPLET APPLICATION (CDA)

When applying Quali-Pro Glyphosate T&O by vehicle-mounted CDA equipment, the rate of this product per acre must not be less than the amount recommended on this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, use 3 to 15 gallons of water per acre.

To control labeled annual weeds with hand-held CDA equipment, apply a 20-percent solution of Quali-Pro Glyphosate T&O at a flow rate of 2 fluid ounces per minute and a walking speed of 2.2 feet/second (equivalent to 1 quart per acre). To control labeled perennial weeds, apply a 20- to 40-percent solution of Quali-Pro Glyphosate T&O at a flow rate of 2 fluid ounces per minute with a walking speed of 1.1 feet/second (equivalent to 2 to 4 quarts per acre).

The spray pattern of CDA equipment is not easily visible. Exercise extreme care to avoid spray or drift contacting any foliage or green tissue of desirable vegetation. Severe damage or destruction may result.

HAND-HELD and HIGH-VOLUME EQUIPMENT

Use coarse sprays only.

Clean water should be used when mixing this product; then apply to the foliage of vegetation to be controlled. For those applications that are to be made on a spray-to-wet basis, ensure uniform and complete spray coverage. Do not spray product to the point of runoff.

To control annual weeds that are listed on this label, spray a 0.5-percent solution of Quali-Pro Glyphosate T&O plus nonionic surfactant to weeds that are smaller than 6 inches in height or runner length. This product should be applied before seedhead formation in grass or before bud formation in broadleaf weeds. Do not till or mow for at least three days after application, unless otherwise specified.

Unless otherwise specified, for annual weeds over 6 inches tall, or if additional surfactant is not to be used, use a 1-percent solution. On harder to control perennials (such as dock, field bindweed, hemp dogbane, bermudagrass, milkweed, and Canada thistle), best results are obtained by using a 2-percent solution.

If using application methods that result in less than total coverage, a 5-percent solution should be used for perennial and annual weeds, and use a 5- to 10-percent solution for trees and woody brush.

To prepare the desired volume of spray solution, mix the proper

amount of Quali-Pro Glyphosate T&O in water in accordance with the table below.

Spray Solution

AMOUNT OF QUALI-PRO GLYPHOSATE T&O						
DESIRED VOLUME	1/2%	1 %	1 1/2 %	2 %	5%	10%
1 gallon	2/3 oz	1-1/3 oz	2 oz	2-2/3 oz	6-1/2 oz	13 oz
25 gallons	1 pt	1 qt	1-1/2 qt	2 qt	5 qt	10 qt
100 gallons	2 qt	1 gal	1-1/2 gal	2 gal	5 gal	10 gal

Note: 2 tablespoons = 1 fluid ounce

When using knapsack sprayers, mix the recommended amount of Quali-Pro Glyphosate T&O with water in a larger container. Then fill sprayer with the mixed solution.

SELECTIVE EQUIPMENT

Quali-Pro Glyphosate T&O may be applied through a recirculating spray system, a hooded sprayer, a sponge bar, a wiper applicator, or a shielded applicator after dilution and thorough mixing with water. It may be applied to the weeds listed on this label that are growing in any noncrop site specified on this label, but only when specifically recommended in cropping systems.

A recirculating spray system works by directing the spray solution onto those weeds growing above desirable vegetation. Any spray solution not intercepted by the weeds is then collected and returned to the sprayer for reuse.

A shielded applicator works by directing the herbicide solution onto weeds, while at the same time shielding desirable vegetation from the herbicide.

A wiper applicator works by rubbing weeds with an absorbent material that contains the herbicide solution.

IMPORTANT: DO NOT LET THIS PRODUCT OR SPRAY MIXTURE COME INTO CONTACT WITH DESIRABLE VEGETATION.

If this product comes into contact with desirable vegetation, severe damage or destruction may result. Adjust applicators that are used above desired vegetation to ensure the lowest spray stream or wiper contact point is a minimum of 2 inches above desirable vegetation. Discoloration, stunting, or plant destruction may result if mist, foam, droplets, or splatter of the herbicide solution settles on desirable vegetation.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps,

severe infestations, or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

SHIELDED APPLICATORS

When this product is applied in accordance with label instructions for shielded applicators, Quali-Pro Glyphosate T&O will control weeds listed in the **WEEDS CONTROLLED** sections of this label.

The following calculation should be used to convert from a broadcast rate per acre to a band rate per acre:

Band width in inches	X	Herbicide Broadcast RATE per acre	=	Herbicide Band RATE per acre
Row width in inches				
Band width in inches	X	Broadcast VOLUME of solution per acre	=	Band VOLUME of solution per acre
Row width in inches				

Nozzles that provide uniform coverage within the area treated should be utilized. To protect desirable vegetation, make sure shields on shielded sprayers are adjusted properly. **EXERCISE EXTREME CAUTION TO AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION.**

See the **WEEDS CONTROLLED** sections of this label for specific rates of application and instructions for control of perennial and annual weeds.

WIPER APPLICATORS AND SPONGE BARS

Wiper applicators are meant to physically wipe an appropriate amount of Quali-Pro Glyphosate T&O onto weeds.

Wiper applicator equipment must be designed, operated, and maintained to ensure the herbicide solution does not contact desirable vegetation. This equipment should be operated at ground speeds of 5 mph or less. Improved performance may be attained by reducing speed in areas having heavy weed infestations; this ensures adequate wiper saturation. Better results are possible if 2 applications are made in opposite directions.

Avoid dripping or leaking herbicide onto desirable vegetation. To ensure adequate contact with weed surfaces, adjust the height of the applicator. Wiping surfaces should be kept clean. If applying on sloping ground, this product may migrate, thereby causing dripping on the lower end and wick drying on the upper end of the wiper applicator.

If weeds are wet, Do not use wiper equipment.

When mixing the herbicide solution, mix only enough solution to be used during a 1-day period; reduced effectiveness may result from using leftover solution amounts. Wiper parts should be cleaned

immediately by thoroughly flushing with water when application is completed.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators – Mix 1 gallon of this product in 2 gallons of water to prepare a 33-percent solution. Apply this solution to weeds listed in this **WIPER APPLICATORS AND SPONGE BARS** section. For Industrial, Turf, and Ornamental uses, solutions ranging from 33 to 75 percent of Quali-Pro Glyphosate T&O in water may be used.

For Porous-Plastic Applicators: Solutions varying from 33 to 100 percent of Quali-Pro Glyphosate T&O in water may be used.

For Panel Applicators and Pressure-feed Systems: Solutions ranging from 33 to 100 percent of this product in water may be used.

Quali-Pro Glyphosate T&O CONTROLS the weeds listed below when applied using **WIPER APPLICATORS AND SPONGE BARS** as recommended under this section.

ANNUAL GRASSES

Corn
Zea mays

Panicum, Texas
Panicum texanum

Rye, common
Secale cereale

Shattercane
Sorghum bicolor

ANNUAL BROADLEAVES

Sicklepod
Cassia obtusifolia

Spanishneedles
Bidens bipinnata

Starbur, bristly
Acanthospermum hispidum

Quali-Pro Glyphosate T&O SUPPRESSES the weeds listed below when applied using **WIPER APPLICATORS AND SPONGE BARS** as recommended under the conditions described in this section.

ANNUAL BROADLEAVES

Beggarweed, Florida
Desmodium tortuosum

Dogfennel
Eupatorium capilliflorum

Pigweed, redroot
Amaranthus retroflexus
Ragweed, common
Ambrosia artemisiifolia

Ragweed, giant
Ambrosia trifida

Sunflower
Helianthus annuus

Thistle, musk
Carduus nutans

Velvetleaf
Abutilon theophrasti

PERENNIAL GRASSES

Bermudagrass
Cynodon dactylon

Guineagrass
Panicum maximum

Johnsongrass
Sorghum halepense

Smutgrass
Sporobolus poiretii

Vaseygrass
Paspalum urvillei

PERENNIAL BROADLEAVES

Dogbane, hemp
Apocynum cannabinum

Milkweed
Asclepias syriaca

Nightshade, silverleaf
Solanum elaeagnifolium

Thistle, Canada
Cirsium arvense

WEEDS CONTROLLED

This product controls many annual and perennial grasses and broadleaf weeds. For Industrial, Turf, and Ornamental use applications, see the section titled **NONCROP USES** for specific weeds controlled.

ANNUAL WEEDS

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage, unless otherwise specified.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.
- Apply this product before seedhead formation to prevent seed production.
- Because Quali-Pro Glyphosate T&O does NOT provide residual control, delay application until maximum weed emergence. Users may have to repeat treatments to control weeds that germinate later.

LOW-VOLUME BROADCAST APPLICATION (LOW-RATE TECHNOLOGY)

Quali-Pro Glyphosate T&O CONTROLS the weeds listed below when applied as follows:

1. Recommended water carrier volumes, unless otherwise specified: 3 to 10 gallons per acre for ground applications; and 3 to 5 gallons per acre when applying by air. (For approved sites, review the section of this label titled **APPLICATION EQUIPMENT AND TECHNIQUES**.)
2. A nonionic surfactant may be added at 0.5 to 1 percent by total spray volume. When using surfactants containing a minimum of 70 percent active ingredient, use 0.5-percent surfactant concentration. When using surfactants containing less than 70 percent active ingredient, use a 1-percent surfactant concentration.

NOTE

- Improved performance on annual weeds may be attained with the addition of 2 percent dry ammonium sulfate by weight, or 17 pounds per 100 gallons of water. Review the sections of this label titled **MIXING ADDITIVES** and **APPLICATION INSTRUCTIONS**.
- Unless otherwise specified, Do not tank mix with soil residual herbicides when using these rates.
- If weeds have been grazed, cut, or mowed, allow regrowth to occur prior to applying this product.
- For control of additional broadleaf weeds, review the section of this label titled **TANK MIXTURES**.

WEED SPECIES	MAXIMUM HEIGHT / LENGTH	RATE PER ACRE* (Fluid Ounces)
Foxtail <i>Setaria</i> spp.	12"	8 oz.
Barleygrass <i>Echinochloa crus-galli</i>	6" (0 to 4") (4 to 6")	12 oz. (16 oz.) (24 oz.)
Bluegrass, annual <i>Poa annua</i>		
Brome, downy** <i>Bromus tectorum</i>		

WEED SPECIES	MAXIMUM HEIGHT / LENGTH	RATE PER ACRE* (Fluid Ounces)
Mustard, blue <i>Chorispora tenella</i>	6"	12 oz.
Mustard, tansy <i>Descurainia pinnata</i>		
Mustard, tumble <i>Sisymbrium altissimum</i>		
Mustard, wild <i>Sinapis arvensis</i>		
Spurry, umbrella <i>Holosteum umbellatum</i>		
Barley <i>Hordeum vulgare</i>	12"	
Rye, common <i>Secale cereale</i>		
Sandbur, field <i>Cenchrus</i> spp.		
Shattercane <i>Sorghum bicolor</i>		
Stinkgrass <i>Eragrostis cilianensis</i>		
Wheat <i>Triticum aestivum</i>	18"	
Morningglory <i>Ipomoea</i> spp.	2"	16 oz.
Sicklepod <i>Cassia obtusifolia</i>		
Bluegrass, bulbous <i>Poa bulbosa</i>	6"	
Cheat <i>Bromus secalinus</i>		
Chickweed, common <i>Stellaria media</i>		
Chickweed, mouseear <i>Cerastium vulgatum</i>		

WEED SPECIES	MAXIMUM HEIGHT / LENGTH	RATE PER ACRE* (Fluid Ounces)
Corn <i>Zea mays</i>	6"	16 oz.
Goatgrass, jointed <i>Aegilops cylindrica</i>		
Groundsel, common <i>Senecio vulgaris</i>		
Henbit <i>Lamium amplexicaule</i>		
Horseweed/Marestail*** <i>Conyza canadensis</i>		
Lambsquarters, common <i>Chenopodium album</i>		
Pennycress, field Fanweed <i>Thlaspi arvense</i>		
Rocket, London <i>Sisymbrium irio</i>		
Ryegrass, Italian <i>Lolium multiflorum</i>		
Shepherd's Purse <i>Capsella bursa-pastoris</i>		
Spurge, annual <i>Euphorbia</i> spp.		
Buttercup <i>Ranunculus</i> spp.	12"	16 oz.
Cocklebur <i>Xanthium strumarium</i>		
Crabgrass <i>Digitaria</i> spp.		
Dwarf dandelion <i>Krigia cespitosa</i>		
Falseflax, smallseed <i>Camelina microcarpa</i>		
Foxtail, Carolina <i>Alopecurus carolinianus</i>		

WEED SPECIES	MAXIMUM HEIGHT / LENGTH	RATE PER ACRE* (Fluid Ounces)
Johnsongrass, seedling <i>Sorghum halepense</i>	12"	16 oz.
Oats, wild <i>Avena fatua</i>		
Panicum, fall <i>Panicum dichotomiflorum</i>		
Panicum, Texas <i>Panicum texanum</i>		
Pigweed, redroot <i>Amaranthus retroflexus</i>		
Pigweed, smooth <i>Amaranthus hybridus</i>		
Witchgrass <i>Panicum capillare</i>		
Sicklepod <i>Cassia obtusifolia</i>	3 to 4"	24 oz.
Signalgrass, broadleaf <i>Brachiaria platyphylla</i>	4"	
Horseweed/Marestail*** <i>Conyza canadensis</i>	7 to 12"	
Lambsquarters, common <i>Chenopodium album</i>		
Spurge, annual <i>Euphorbia</i> spp.		
Rice, red <i>Oryza sativa</i>	4"	32 oz.
Teaweed <i>Sida spinosa</i>		
Sprangletop <i>Leptochloa</i> spp.	6"	
Geranium, Carolina <i>Geranium carolinianum</i>	12"	
Goosegrass <i>Eleusine indica</i>		

WEED SPECIES	MAXIMUM HEIGHT / LENGTH	RATE PER ACRE* (Fluid Ounces)
Primrose, cutleaf evening <i>Oenothera laciniate</i>	12"	32 oz.
Pusley, Florida <i>Richardia scabra</i>		
Sicklepod <i>Cassia obtusifolia</i>	5 to 12"	
Spanishneedles <i>Bidens bipinnata</i>		
Filaree <i>Erodium</i> spp.	12"	48 oz.
Sprangletop <i>Leptochloa</i> spp.		

*Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana, and Texas for preplant treatments.

* For those weeds with recommended rates of less than 32 fluid ounces per acre, this product may be used at rates up to 32 fluid ounces per acre where heavy weed densities exist.

** Use 16 fluid ounces per acre for control in no-till systems.

*** Resistance to some biotypes has been observed. If users encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

● TANK MIXTURES

● QUALI-PRO GLYPHOSATE T&O plus BANVEL plus NONIONIC SURFACTANT

● QUALI-PRO GLYPHOSATE T&O plus 2,4-D plus NONIONIC SURFACTANT

IN CALIFORNIA, DO NOT APPLY BANVEL OR 2,4-D TANK MIXTURES BY AIR.

These tank mixtures are recommended for use in fallow and reduced tillage areas only. Follow use directions as given in the **LOW-VOLUME BROADCAST APPLICATION** section.

For tank mixture instructions relating to Industrial, Turf, and Ornamental applications, see the section titled **NONCROP USES**.

A mixture of Quali-Pro Glyphosate T&O plus 2,4-D or Banvel can be used to control the broadleaf weeds and annual grasses listed for Quali-Pro Glyphosate T&O alone at the indicated heights (except applications at 8 fluid ounces per acre), plus the broadleaf weeds listed below. For weeds listed at 8 fluid ounces per acre of Quali-Pro Glyphosate T&O alone, 12 fluid ounces should be used in these tank mixtures.

NOTE: For all products used in tank mixtures, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements. Also, review each product's label for crop rotation restrictions and cautionary statements. If Banvel is mixed with Quali-Pro Glyphosate T&O, short-term residual control of selected weed species may result. However, some crop injury is possible if Banvel is applied within 45 days of planting.

To control dense populations of the annual broadleaf weeds listed below (when weeds are less than the indicated height), apply 12 to 16 fluid ounces of Quali-Pro Glyphosate T&O plus 0.25 pound active ingredient of Banvel, or 0.5 pound active ingredient of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre.

Cocklebur (12")
Xanthium strumarium

Kochia* (6")
Kochia scoparia

Lambsquarters (12")
Chenopodium album

Lettuce, prickly (6")
Lactuca serriola

Marestail/Horseweed** (6")
Conyza canadensis

Morningglory (6")
Ipomoea spp.

Pigweed, redroot (12")
Amaranthus retroflexus

Pigweed, smooth (12")
Amaranthus hybridus

Thistle, Russian (12")
Salsola kali

* Kochia is controlled with a Banvel tank mixture only.

** Resistance to some biotypes has been observed. If users encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

To control the annual broadleaf weeds listed below (when less than 6 inches in height), apply 16 fluid ounces of Quali-Pro Glyphosate T&O plus 0.5 pound active ingredient of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre.

Ragweed, common
Ambrosia artemisiifolia

Ragweed, giant
Ambrosia trifida

Smartweed, Pennsylvania
Polygonum pennsylvanicum

Velvetleaf
Abutilon theophrasti

HIGH-VOLUME BROADCAST APPLICATIONS

Quali-Pro Glyphosate T&O will control weeds listed below when applied as directed using water carrier volumes of 10 to 40 gallons per acre for ground applications.

Unless otherwise specified, apply 1 to 1.5 quarts of Quali-Pro Glyphosate T&O per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume. 1 quart Quali-Pro Glyphosate T&O per acre should be used if weeds do not exceed 6 inches tall, and 1.5 quarts per acre should be used if weeds are more than 6 inches tall. Before application, allow sufficient time for new growth to reach recommended stages if weeds have been cut, grazed, or mowed. These rates will also control weeds mentioned in the **LOW VOLUME BROADCAST APPLICATION** section of this label.

WEED SPECIES

Balsamapple*
Momordica charantia

Bassia, fivehook
Bassia hyssopifolia

Brome
Bromus spp.

Fiddleneck
Amsinckia spp.

Fleabane, hairy
Conyza bonariensis

Fleabane
Erigeron spp.

Kochia
Kochia scoparia

Lettuce, prickly
Lactuca serriola

Panicum
Panicum spp.

Ragweed, common
Ambrosia artemisiifolia

Ragweed, giant
Ambrosia trifida

Smartweed, Pennsylvania
Polygonum pennsylvanicum

Sowthistle, annual
Sonchus oleraceus

Sunflower
Helianthus annuus

Thistle, Russian
Salsola kali

Velvetleaf
Abutilon theophrasti

*Apply with hand-held equipment only.

PERENNIAL WEEDS

For controlling or destroying most perennial weeds, apply Quali-Pro Glyphosate T&O as follows.

For application instructions and perennial weeds controlled for Industrial, Turf, and Ornamental uses, see the section titled **NONCROP USES**.

NOTE: In the event weeds have been tilled or mowed, do not treat plants until they have resumed active growth AND have reached the recommended stages.

It may be necessary to repeat treatments on those weeds that regenerate from underground parts or from seed. These repeat treatments must be applied before crops emerge.

Improved performance on perennial weeds may be attained with the addition of 1 to 2 percent dry ammonium sulfate by weight, or 8.5 to 17 pounds per 100 gallons of water. The increased effectiveness may be most apparent if environmental stress is present. Review the sections of this label titled **MIXING ADDITIVES** and **APPLICATION INSTRUCTIONS**.

Quali-Pro Glyphosate T&O will **CONTROL** the **PERENNIAL WEEDS** listed below if applied as directed under the conditions described.

Alfalfa
Medicago sativa

Alligatorweed*
Alternanthera philoxeroides

Anise (fennel)
Foeniculum vulgare

Artichoke, Jerusalem
Helianthus tuberosus

Bahiagrass
Paspalum notatum

Bentgrass
Agrostis spp.

Bermudagrass
Cynodon dactylon

Bermudagrass, water (knotgrass)
Paspalum distichum

Bindweed, field
Convolvulus arvensis

Bluegrass, Kentucky
Poa spp.

Blueweed, Texas
Helianthus ciliaris

Brackenfern
Pteridium aquilinum

Bromegrass, smooth
Bromus inermis

Bursage, woollyleaf
Franseria tomentosa

Canarygrass, reed
Phalaris arundinacea

Cattail
Typha spp.

Clover, red
Trifolium pratense

Clover, white
Trifolium repens

Cogongrass
Imperata cylindrica

Dallisgrass
Paspalum dilatatum

Dandelion
Taraxacum officinale

Dock, curly
Rumex crispus

Dogbane, hemp
Apocynum cannabinum

Fescues
Festuca spp.

Fescue, tall
Festuca arundinacea

Guineagrass
Panicum maximum

Horsenettle
Solanum carolinense

Horseradish
Armoracia rusticana

Ice plant
Mesembryanthemum crystallinum

Johnsongrass
Sorghum halepense

Kikuyugrass
Pennisetum clandestinum

Knapweed
Centaurea repens

Lantana
Lantana camara

Lespedeza
Lespedeza spp.

Milkweed
Asclepias spp.

Muhly, wirestem
Muhlenbergia frondosa

Mullein, common
Verbascum thapsus

Napiergrass
Pennisetum purpureum

Nightshade, silverleaf
Solanum elaeagnifolium

Nutsedge; purple, yellow
Cyperus rotundus
Cyperus esculentus

Orchardgrass
Dactylis glomerata

Pampasgrass
Cortaderia spp.

Paragrass
Brachiaria mutica

Phragmites*
Phragmites spp.

Poison hemlock
Conium maculatum

Quackgrass
Agropyron repens

Redvine*
Brunnichia ovata

Reed, giant
Arundo donax

Ryegrass, perennial
Lolium perenne

Smartweed, swamp
Polygonum cockineum

Spurge, leafy*
Euphorbia esula

Starthistle, yellow
Centaurea solstitialis

Sweet potato, wild*
Ipomoea pandurata

Thistle, Canada
Cirsium arvense

Thistle, artichoke
Cynara cardunculus

Timothy
Phleum pratense

Torpedograss*
Panicum repens

Trumpet creeper*
Campsis radicans

Vaseygrass
Paspalum urvillei

Velvetgrass
Holcus spp.

Wheatgrass, western
Agropyron smithii

*Partial Control

Quali-Pro Glyphosate T&O is NOT registered for use on water bermudagrass in California.

For specific application instructions and labeled uses, review the **DIRECTIONS FOR USE** and **MIXING, ADDITIVES, and APPLICATION INSTRUCTIONS** sections of this label.

Alfalfa: Use 1 quart of Quali-Pro Glyphosate T&O per acre, plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply in the fall after the last hay cutting takes place. Alfalfa should be allowed to regrow to a minimum height of 6 to 8 inches prior to application. Follow applications with deep tillage a minimum of 7 days after treatment, but before soil freeze-up.

Alligatorweed: Use 4 quarts of Quali-Pro Glyphosate T&O per acre, or a 1.5-percent solution with hand-held equipment to obtain partial control. Applications should be made when the majority of plants are in bloom. Additional applications will be necessary to maintain control.

Anise (fennel); Poison Hemlock: Use a 1- to 2-percent solution of Quali-Pro Glyphosate T&O as a spray-to-wet treatment. Plants should be treated at the bud to full-bloom stage of growth to obtain best results. In order to control plants arising from seeds, repeat applications may be needed in succeeding years.

Bentgrass: Use Quali-Pro Glyphosate T&O for suppression in grass seed production areas. For use by ground applications only. Apply 1.5 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Prior to applying in the fall, ensure the entire crown area has resumed growing. Plants should be actively growing, and have a minimum of 3 inches of growth. Avoid tillage prior to application. For optimum results, tillage 7 to 10 days after application is recommended. Ineffective control of bentgrass may result if tillage is not used after treatment.

Bermudagrass: Use 5 quarts of Quali-Pro Glyphosate T&O per acre for adequate control; or, 3 quarts per acre will provide partial control. For best results, treat bermudagrass when it is actively growing, and when seedheads are present. A repeat application may be necessary to maintain control. Allow a minimum of 7 days after treatment before tillage.

Bermudagrass, water (knotgrass): Use 1.5 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Water bermudagrass should be actively growing—and 12 to 18 inches in length—for effective application. Wait a minimum of 7 days before flooding the field, flushing, or tillage.

For fall treatment only: Use 1 quart of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Till fallow fields prior to treatment. Apply prior to frost on plants that are actively growing and 12 to 18 inches long. Wait a minimum of 7 days before tillage.

Bindweed, field: For control of field bindweed, apply 3 to 4 quarts of

Quali-Pro Glyphosate T&O per acre east of the Mississippi River, and 4 to 5 quarts per acre west of the Mississippi River. Treatment should be applied when weeds are actively growing and are at or beyond full bloom. If the weed is under drought stress, do not treat—good soil moisture is needed for active growth. Apply in late summer or fall for optimum results, but fall applications must be made before a killing frost. Wait a minimum of 7 days after application before tillage.

For control using ground application equipment only, apply 2 quarts of Quali-Pro Glyphosate T&O plus 0.5 pound active ingredient of Banvel in 10 to 20 gallons of water per acre.

The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.

To suppress field bindweed on irrigated agricultural land (using ground equipment only), mix and apply 1 to 2 quarts of Quali-Pro Glyphosate T&O plus 1 pound of 2,4-D active ingredient in 10 to 20 gallons of water per acre. Apply after harvest or in fall fallow ground when most of the runners are a minimum of 12 inches long and are actively growing. Use of at least one irrigation will promote active field bindweed growth.

For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

In California only: Use 1 to 5 quarts of Quali-Pro Glyphosate T&O per acre. Depending on local conditions, the rate needed to suppress or control weeds will vary within this range.

To suppress field bindweed on irrigated land where annual tillage is performed, apply 1 quart of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Bindweed should be a minimum of 12 inches long before application, with maximum weed emergence and runner growth. Treatments should not be made when weeds are under drought stress; good soil moisture is needed for active growth. Allow a minimum of 3 days after application before tillage.

Bluegrass, Kentucky; Bromegrass, smooth; Orchardgrass: Use 2 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. Apply when grasses are actively growing and when most plants have developed to the boot-to-early seedhead stage. For partial control in pasture or hay crop renovation, use 1 to 1.5 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing weeds when most are 4 to 12 inches high. Wait a minimum of 7 days after treatment before tillage.

Orchardgrass (sods going to no-till corn): Use 1 to 1.5 quarts of Quali-Pro Glyphosate T&O per acre plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Orchardgrass should be at least 12 inches high for spring treatment, and 6 inches high for fall treatment. Wait a minimum of 3 days before planting. For best results, a sequential application of atrazine is necessary. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Texas Blueweed: Apply 3 to 4 quarts of Quali-Pro Glyphosate T&O per acre east of the Mississippi River, and 4 to 5 quarts per acre west of the Mississippi River. Treat when weeds are actively growing and at full bloom or beyond. If weeds are under drought stress, do not treat—good soil moisture is needed for active growth. (If new leaves are developing, this indicates active growth.) Apply in late summer or fall for optimum results. Fall applications must be made before a killing frost. Wait a minimum of 7 days after treatment before tillage.

Brackenfern: Using a broadcast spray, apply 3 to 4 quarts of Quali-Pro Glyphosate T&O per acre. With hand-held equipment, use a 1- to 1.5-percent solution. Treatments should be applied to fully expanded fronds that are a minimum of 18 inches in height.

Bursage, woollyleaf: For control, apply 2 quarts of Quali-Pro Glyphosate T&O plus 1 pint of Banvel per acre. For partial control, apply 1 quart of Quali-Pro Glyphosate T&O plus 1 pint of Banvel per acre. Add 0.5- to 1-percent nonionic surfactant by total spray volume and apply in 3 to 20 gallons of water per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Canarygrass, reed; Timothy; Wheatgrass, western: Apply 2 to 3 quarts of Quali-Pro Glyphosate T&O per acre to actively growing plants when most have reached the boot-to-head stage of development. Wait a minimum of 7 days after treatment before tillage.

Cogongrass: Use 3 to 5 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant applied in 10 to 40 gallons of water per acre. Apply in late summer or fall when weeds are a minimum of 18 inches high and actively growing. Wait a minimum of 7 days after application before mowing or tillage. Because of uneven stages of growth and the dense nature of Cogongrass, good spray coverage is often difficult. Therefore, repeat treatments may be needed to maintain adequate control.

Dandelion / Dock, curly: Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Wait 7 or more days after application before tillage.

Another option for control: Apply 16 fluid ounces of Quali-Pro Glyphosate T&O plus 0.5 pound of 2,4-D active ingredient plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10

gallons of water per acre. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Dogbane, hemp: Apply 4 quarts of Quali-Pro Glyphosate T&O per acre when most weeds are actively growing and have reached the late bud to flower stage of development. Before treating after a crop harvest or mowing, allow weeds to regrow to a mature stage of development. Wait a minimum of 7 days after treatment before tillage.

For suppression, using ground applications only, apply 16 fluid ounces of Quali-Pro Glyphosate T&O plus 0.5 pounds of 2,4-D active ingredient plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply after maximum emergence of dogbane has occurred. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Fescue, tall: Apply 3 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.

For applications in fall only: Use 1 quart of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply in fall when weeds are actively growing and have reached 6 to 12 inches of new growth. Wait a minimum of 7 days after treatment before tillage. To improve long-term control, and to control seedlings germinating after fall treatments (or the spring that follows), apply a sequential application of 1 pint per acre of Quali-Pro Glyphosate T&O plus nonionic surfactant.

Guineagrass: Apply 3 quarts of Quali-Pro Glyphosate T&O per acre, or when using hand-held equipment, use a 1-percent solution. Apply when guineagrass is actively growing and when it has reached at least the 7-leaf stage of development. If using hand-held equipment, ensure thorough coverage. Wait a minimum of 7 days after treatment before tillage.

Johnsongrass; Ryegrass, perennial: Use 1 to 3 quarts of Quali-Pro Glyphosate T&O per acre. Use 1 to 2 quarts of Quali-Pro Glyphosate T&O per acre in annual cropping systems. Use 1 quart of this product plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Or use 2 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. In noncrop areas, or in areas where annual tillage is not performed (no-till), use 2 to 3 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. Optimum results are obtained when applied to actively growing plants when most have reached the boot-to-head stage of development, or in fall before frost. Wait a minimum of 7 days after treatment before tillage. If using the 1-quart-per-acre rate, do not tank mix with residual herbicides.

For johnsongrass burndown, use 1 pint per acre plus 0.5- to 1-percent nonionic surfactant in 3 to 10 gallons of water per acre before the plants grow to a height of 12 inches. Wait a minimum of 3 days after application before tillage.

For spot treatment of johnsongrass (partial control or suppression): When Johnsongrass has reached a height of 12 to 18 inches, apply a 1-percent solution of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume. Complete and uniform coverage is essential.

Kikuyugrass: Use 2 to 3 quarts of Quali-Pro Glyphosate T&O per acre. Spray when the majority of kikuyugrass is actively growing and is at a minimum of 8 inches high (at the 3- or 4-leaf stage of development). Wait a minimum of 3 days after treatment before tillage.

Knapweed; Horseradish: Use 4 quarts of Quali-Pro Glyphosate T&O per acre when weeds are actively growing and when most have reached the late bud to flower stage of development. If treating after crop harvest or after mowing, allow weeds to regrow to a mature stage of development, then apply. Apply treatment in late summer or fall for optimum results. Wait a minimum of 7 days after treatment before tillage.

Lantana: Apply Quali-Pro Glyphosate T&O as a 1- to 1.25-percent solution using hand-held equipment only. Treat actively growing lantana when it is at or beyond the bloom stage of development. The higher application rate of Quali-Pro Glyphosate T&O should be applied for weeds that have reached the woody stage of development. Wait a minimum of 7 days after treatment before tillage.

Milkweed, common: Use 3 quarts of Quali-Pro Glyphosate T&O per acre when milkweed is actively growing and when most of the weeds have reached the late bud to flower stage of development. If applying after small grain harvest or after mowing, allow milkweed to regrow to a mature stage prior to applying. Wait a minimum of 7 days after treatment before tillage.

Muhly, wirestem: Use 1 to 2 quarts of Quali-Pro Glyphosate T&O per acre. Apply 1 quart of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of Quali-Pro Glyphosate T&O per acre in 10 to 40 gallons of water per acre, or when applied in pasture, sod, or noncrop areas. Spray when weed is actively growing and at least 8 inches high. Do not till between harvest and fall applications, or in the fall or spring prior to spring treatments. Wait a minimum of 3 days after treatment before tillage. Quali-Pro Glyphosate T&O will not provide residual control of wirestem muhly from seeds which germinate after this product is applied. When using the 1-quart-per-acre rate, do not tank mix with residual herbicides.

Nightshade, silverleaf: For control, use 2 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Do not apply until at least 60 percent of the plants have berries. If applying in fall, do so before a killing frost. Wait a minimum of 7 days after treatment before tillage. If the weeds are under drought stress, do not treat—good soil moisture is needed for active growth.

Nutsedge—purple, yellow: Using a broadcast spray, apply 3 quarts of Quali-Pro Glyphosate T&O per acre. If using hand-held equipment, use a 1- to 2-percent solution to control existing

nutsedge plants and immature nutlets that are attached to treated plants. Apply when weeds are in flower or when new nutlets are seen at rhizome tips. Nutlets that have not yet germinated will not be controlled and may germinate after treatment. For long-term control of ungerminated tubers, repeat treatments will be necessary.

For control, apply sequential applications of 1 to 2 quarts of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Applications should be made when a majority of the weeds are in the 3- to 5-leaf stage of development (when under 6 inches tall). Repeat this application as necessary when newly emerging weeds reach the 3- to 5-leaf stage of growth. For long-term control, subsequent treatments are required.

For suppression or partial control of existing weeds, use 1 pint to 2 quarts of Quali-Pro Glyphosate T&O per acre plus 0.5- to 1-percent nonionic surfactant in 3 to 40 gallons of water per acre. Apply when plants are 6 inches tall or less and have 3 to 5 leaves. Additional treatments will be necessary to control subsequent emerging weeds or regrowth of existing weeds. Wait a minimum of 7 days after treatment before mowing or tillage.

Pampasgrass / Ice plant: Using hand-held equipment, apply Quali-Pro Glyphosate T&O as a 1.5- to 2-percent solution to weeds that are actively growing. Apply when pampasgrass is at or beyond the boot stage of development. Thorough coverage is essential.

Phragmites: For partial control in Florida and those counties of states that border the Gulf of Mexico, use 5 quarts per acre using broadcast spray equipment, or use a 2-percent solution applied by hand-held equipment. For partial control in other areas of the U.S., use 3 quarts per acre using broadcast spray equipment or a 1-percent solution applied by hand-held equipment. For optimum results, treat in late summer or fall, or when weeds are actively growing and in the full bloom stage of development. If treated before or after this stage, reduced control may result. The dense nature of the weeds and the uneven stages of growth may prevent good spray coverage; therefore, repeat treatments may be needed to maintain control. Visual control symptoms are slow to develop.

Quackgrass-In Annual Cropping Systems, or in Pastures and Sodds Followed by Deep Tillage: Use 1 to 2 quarts of Quali-Pro Glyphosate T&O per acre. If using the 1-quart rate: Apply 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre; do not tank mix with residual herbicides. If using the 2-quart rate: Apply treatment in 10 to 40 gallons of water per acre. To apply, wait until quackgrass is actively growing and is from 6 to 8 inches high. Do not till between harvest and fall applications, or in fall or spring prior to spring glyphosate treatments. Wait a minimum of 3 days after treatment before tillage. For best results in pastures or sods, a moldboard plow should be used.

Quackgrass-Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application: Use 2 to 3 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. Quackgrass should be higher than 8 inches tall and actively growing when treated. Do not till between harvest and fall applica-

tions, in fall or spring prior to spring glyphosate treatments. Wait a minimum of 3 days after treatment before tillage.

Redvine: For suppression, use 24 fluid ounces of Quali-Pro Glyphosate T&O per acre at each of two applications 7 to 14 days apart, or a single application of 2 quarts of Quali-Pro Glyphosate T&O per acre. Recommended rates should be applied in 5 to 10 gallons of water per acre, plus 0.5- to 1-percent nonionic surfactant by total volume. Apply to actively growing plants in September or early October, when plants are a minimum of 18 inches tall and have been growing 45 to 60 days since the last tillage. Apply treatments a minimum of 1 week before a killing frost.

Reed, giant: For control, use a 2-percent solution of Quali-Pro Glyphosate T&O when plants are actively growing. For optimum results, apply in late summer to fall.

Smartweed, swamp: Use 3 to 5 quarts of Quali-Pro Glyphosate T&O per acre when plants are actively growing, and when most have reached the early bud stage of development. Wait a minimum of 7 days after treatment before tillage.

Another option for control: Use 16 fluid ounces of Quali-Pro Glyphosate T&O with 0.5 pound of 2,4-D active ingredient, plus 0.5- to 1-percent nonionic surfactant by total volume in 3 to 10 gallons of water per acre in late summer or fall. Treatments should be applied when weeds are actively growing and when the majority of plants have reached the early bud stage of development. Wait a minimum of 7 days after treatment before tillage. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Spurge, leafy: For suppression, use 16 fluid ounces of Quali-Pro Glyphosate T&O with 0.5 pound of 2,4-D active ingredient, plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in late summer or fall. Apply when plants are actively growing. If weeds have been mowed prior to treatment, apply when the majority of the weeds are 12 inches in height. Wait a minimum of 7 days after treatment before tillage. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Starthistle, yellow: For best results, apply during periods of active growth, including the rosette, bolting, and early flower stages of development. When using "spray-to-wet" applications, apply Quali-Pro Glyphosate T&O as a 2-percent solution. When using broadcast applications, use 2 quarts of Quali-Pro Glyphosate T&O per acre in 10 to 40 gallons of water per acre.

Sweet Potato, wild; Thistle, artichoke: Using hand-held equipment, apply Quali-Pro Glyphosate T&O as a 2-percent solution when weeds are actively growing or are beyond the bloom stage of growth. Additional applications may be required. The weeds should be allowed to reach the recommended stage of development prior to re-treatment. Wait a minimum of 7 days after treatment before tillage.

Thistle, Canada: Use 2 to 3 quarts of Quali-Pro Glyphosate T&O per acre applied to actively growing thistles when the majority are at or beyond the bud stage of development. In the late summer or fall after harvest, tillage, or mowing, allow a minimum of 4 weeks for initiation of active growth and rosette development before applying this product. If treating in fall, Quali-Pro Glyphosate T&O must be applied before a killing frost. Wait a minimum of 3 days after treatment before tillage.

For suppression of Canada thistle, use 1 quart per acre of Quali-Pro Glyphosate T&O, or 1 pint of Quali-Pro Glyphosate T&O plus 0.5 pound a.i. 2,4-D per acre, plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Torpedograss: For partial control, use 4 to 5 quarts of Quali-Pro Glyphosate T&O per acre. For applications to be effective, torpedograss must be actively growing, when the majority of weeds are at or beyond the seedhead stage of development. To maintain control, repeat applications are necessary. When applying in fall, treatments must be made before frost. Wait a minimum of 7 days after treatment before tillage.

Trumpet creeper: For control, use 2 quarts of Quali-Pro Glyphosate T&O per acre in 5 to 10 gallons of water per acre. Apply treatments in late September or October to actively growing plants that are a minimum of 18 inches in height and have been growing 45 to 60 days since the last tillage. Applications should be made a minimum of 1 week before a killing frost.

For other perennials listed on this label: Use 3 to 5 quarts of Quali-Pro Glyphosate T&O per acre when perennial weeds are actively growing and most have reached early head or early bud stage of development. Wait a minimum of 7 days after treatment before tillage.

WOODY BRUSH AND TREES

When applying Quali-Pro Glyphosate T&O as directed under the conditions described herein, this herbicide CONTROLS or PARTIALLY CONTROLS the woody brush, trees, and plants listed below.

For application instructions and a list of woody brush and trees controlled for Industrial, Turf, Ornamental uses, see the section titled **NONCROP USES**.

Alder
Alnus spp.

Ash*
Fraxinus spp.

Aspen, quaking
Populus tremuloides

Bearmat (Bearclover)
Chamaebatia foliolosa

Beech
Fagus grandifolia

Birch
Betula spp.

Blackberry
Rubus spp.

Blackgum
Nyssa spp.

Bracken
Peridium spp.

Broom:
French
Cytisus monspessulanus

Scotch
Cytisus scoparius

Buckwheat, California*
Eriogonum fasciculatum

Cascara*
Rhamnus purshiana

Catsclaw*
Acacia greggii

Ceanothus*
Ceanothus spp.

Chamise
Adenostoma fasciculatum

Cherry:
Bitter
Prunus emarginata

Black
Prunus serotina

Pin
Prunus pensylvanica

Coyote brush
Baccharis consanguinea

Creeper, Virginia*
Parthenocissus quinquefolia

Dewberry
Rubus trivialis

Dogwood*
Cornus spp.

Elderberry
Sambucus spp.

Elm*
Ulmus spp.

Eucalyptus
Eucalyptus spp.

Gorse
Ulex europaeus

Hasardia*
Haplopappus squamosus

Hawthorn
Crataegus spp.

Hazel
Corylus spp.

Hickory*
Carya spp.

Holly, Florida / Brazilian Peppertree*
Schinus terebinthifolius

Honeysuckle
Lonicera spp.

Hornbeam, American*
Carpinus caroliniana

Kudzu
Pueraria lobata

Locust, black*
Robinia pseudoacacia

Madrone
Arbutus menziesii

Manzanita
Arctostaphylos spp.

Maple:
Red**
Acer rubrum

Sugar
Acer saccharum

Vine*
Acer circinatum

Monkey Flower*
Mimulus guttatus

Oak:
Black*
Quercus velutina

Northern Pin
Quercus palustris

Post
Quercus stellata

Red
Quercus rubra

Southern Red
Quercus falcata

White*
Quercus alba

Persimmon*
Diospyros spp.

Pine
Pinus spp.

Poison Ivy
Rhus radicans

Poison Oak
Rhus toxicodendron

Poplar, yellow*
Liriodendron tulipifera

Raspberry
Rubus spp.

Redbud, eastern
Cercis canadensis

Rose, multiflora
Rosa multiflora

Russian-olive
Elaeagnus angustifolia

Sage; black, white
Salvia spp.

Sagebrush, California
Artemisia californica

Salmonberry
Rubus spectabilis

Salt cedar
Tamarix spp.

Sassafras
Sassafras albidum

Sourwood
Oxydendrum arboreum

Sumac:
Poison*
Rhus vernix

Smooth*
Rhus glabra

Winged*
Rhus copallina

Sweetgum
Liquidambar styraciflua

Swordfern*
Polystichum munitum

Tallowtree, Chinese
Sapium sebiferum

Tan Oak
Lithocarpus densiflorus

Thimbleberry
Rubus parviflorus

Tobacco, tree*
Nicotiana glauca

Trumpet creeper
Campsis radicans

Waxmyrtle, southern*
Myrica cerifera

Willow
Salix spp.

NOTE: Do not apply to brush that has been mowed or tilled or to trees that have been cut until regrowth has reached the recommended stages of development.

Quali-Pro Glyphosate T&O should be applied when plants are actively growing and after full leaf expansion, unless otherwise directed. The higher rate of application should be used for larger plants and/or dense growth areas. On vines, use the higher rate for plants that have reached the woody stage of development. For optimum results, apply Quali-Pro Glyphosate T&O in late summer or fall after fruit has formed.

Best results are obtained in arid areas when Quali-Pro Glyphosate T&O is applied in spring to early summer, when brush species have a high moisture content and are flowering.

Thorough coverage is essential when using hand-held equipment. Control symptoms may not appear before frost or senescence with fall applications. Wait a minimum of 7 days after treatment before tillage, mowing or removal. Additional treatments may be needed to control plants that regenerate from underground parts or seed. When applying on undesirable deciduous species, some autumn colors are acceptable, as long as no major leaf drop has occurred. Reduced effectiveness may result if fall treatments are made after a frost.

Review the sections of this label titled **DIRECTIONS FOR USE** and **MIXING, ADDITIVES**, and **APPLICATION INSTRUCTIONS** for detailed application information.

To control or partially control the woody brush and trees listed below, apply Quali-Pro Glyphosate T&O as directed.

Alder; Dewberry; Honeysuckle; Post Oak; Raspberry: For control, use 3 to 4 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or as a 1- to 1.5-percent solution using hand-held equipment.

Aspen, quaking; Cherry; Bitter, Black, Pin; Hawthorn; Oak, Southern Red; Sweetgum; Trumpet creeper: For control, use 2 to 3 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or as a 1- to 1.5-percent solution using hand held equipment.

Birch; Elderberry; Hazel; Salmonberry; Thimbleberry: For control, use 2 quarts of Quali-Pro Glyphosate T&O per acre as a broadcast spray, or as a 1-percent solution using hand-held equipment.

Blackberry: For control, use 3 to 4 quarts of Quali-Pro Glyphosate T&O per acre as a broadcast spray, or 1- to 1.5-percent solution with hand-held equipment. Apply after plants have reached full leaf maturity. For best results, apply in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4-percent solution of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

* Partial control

** See below for information on control or partial control

Broom: French, Scotch: For control, use hand-held equipment to apply a 1.5 to 2 percent solution of Quali-Pro Glyphosate T&O.

Buckwheat, California; Hasardia; Monkey Flower; Tobacco, tree: For partial control, use hand-held equipment to apply a foliar spray of a 1- to 2-percent solution of Quali-Pro Glyphosate T&O. Foliage must be thoroughly covered to obtain best results.

Catsclaw: For partial control, use a 1- to 1.5-percent solution of Quali-Pro Glyphosate T&O applied by hand-held equipment.

Coyote Brush: For control, use hand-held equipment and spray a 1.5- to 2-percent solution of Quali-Pro Glyphosate T&O when at least 1/2 of the new leaves are at the full growth stage.

Eucalyptus: For controlling eucalyptus resprouts, use hand-held equipment and spray a 2-percent solution of Quali-Pro Glyphosate T&O when resprouts are 6 to 12 feet in height and plants are actively growing. Complete coverage is required. Avoid use on plants that are drought-stressed.

Kudzu: For control, use 4 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or use hand-held equipment to apply a 2-percent solution. To maintain adequate control, additional applications will be necessary.

Madrone resprouts: For suppression or partial control, use a 2-percent solution of Quali-Pro Glyphosate T&O applied to resprouts that are less than 3 to 6 feet tall. For optimum results, apply in spring or early summer.

Maple, Red: For control, use hand-held equipment and apply a 1- to 1.5-percent solution of Quali-Pro Glyphosate T&O when a minimum of 1/2 of the new leaves of the maple are fully developed. For partial control, use a broadcast spray of 2 to 4 quarts of Quali-Pro Glyphosate T&O per acre.

Maple, Sugar; Oak, Northern Pin; Oak, Red: For control, use hand-held equipment and apply a 1- to 1.5-percent solution of Quali-Pro Glyphosate T&O when a minimum of 1/2 of the new leaves are at the full development stage.

Poison Ivy; Poison Oak: For control, use 4 to 5 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray or, when using hand-held equipment, use a 2-percent solution. To maintain adequate control, additional applications may be necessary. If applying in fall, treatments must be applied before leaves lose their green color.

Rose, multiflora: For control, use 2 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or if using hand-held equipment, apply a 1-percent solution. Treat multiflora rose before leaves begin to deteriorate from leaf-feeding insects.

Sage, black; Sagebrush, California; Chamise; Tallowtree, Chinese: For control, apply a 1-percent solution of Quali-Pro Glyphosate T&O as a foliar spray with hand-held equipment. Thorough coverage is essential.

Tan Oak resprouts: For suppression or partial control, use a 2-percent solution of Quali-Pro Glyphosate T&O applied to resprouts that are less than 3 to 6 feet tall. For optimum results, apply in fall.

Willow: For control, use 3 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or if using hand-held equipment, use a 1-percent solution.

Other Trees and Woody Brush listed on this label: For partial control, use 2 to 5 quarts of Quali-Pro Glyphosate T&O per acre applied as a broadcast spray, or if using hand-held equipment, use a 1- to 2-percent solution.

NONCROP USES

Review the sections of this label titled **GENERAL INFORMATION** and **MIXING, ADDITIVES, and APPLICATION INSTRUCTIONS** for important information about Quali-Pro Glyphosate T&O. Review the following **NONCROP** sections for specific recommended use information.

EXERCISE EXTREME CAUTION TO AVOID SPRAY CONTACT WITH THE FOLLOWING: FOLIAGE; EXPOSED NONWOODY ROOTS OR GREEN STEMS; CROP FRUITS; DESIRABLE TURF-GRASSES; SHRUBS; TREES; OR OTHER DESIRABLE VEGETATION. DESTRUCTION OR SEVERE DAMAGE MAY RESULT.

Additional treatments may be needed to adequately control weeds that regenerate from seeds or underground parts.

If multiple treatments are needed, use a maximum of 10.6 quarts of Quali-Pro Glyphosate T&O per acre per year. **NOTE:** THE MAXIMUM USE RATES DESCRIBED IN THIS LABEL APPLY TO QUALI-PRO GLYPHOSATE T&O WHEN COMBINED WITH ALL OTHER HERBICIDES CONTAINING THE ACTIVE INGREDIENTS GLYPHOSATE OR SULFOSATE—WHETHER APPLIED AS A MIXTURE OR APPLIED SEPARATELY. USERS MUST CAREFULLY CALCULATE ALL APPLICATION RATES AND MAKE SURE THAT TOTAL USAGE OF QUALI-PRO GLYPHOSATE T&O AND OTHER PRODUCTS CONTAINING GLYPHOSATE OR SULFOSATE DOES NOT EXCEED THE RECOMMENDED MAXIMUM USAGE RATES.

Quali-Pro Glyphosate T&O does not provide residual control of weeds. Users should follow a label-approved herbicide program for subsequent weed control.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

INDUSTRIAL, RECREATIONAL, AND PUBLIC AREAS

When applied as directed under the **NONCROP USES** section of this label, and under the corresponding conditions described, Quali-Pro Glyphosate T&O provides control of those perennial and annual weeds described on this label, in areas such as the following:

Airports
Ditch banks
Dry ditches

Parks
Petroleum tank farms and pumping installations
Pipelines

Dry canals	Power and telephone rights-of-way
Fencerows	Railroads
Golf courses	Roadsides
Highways	Schools
Industrial plant sites	Storage areas
Lumber yards	Utility substations
Parking areas	Other public areas and similar industrial or noncrop areas

Review the **WEEDS CONTROLLED** section of this label for application rates and other instructions for control of perennial weeds, annual weeds, trees, and woody brush.

Recirculating sprayers, wiper applicators, or shielded applicators may be used to apply Quali-Pro Glyphosate T&O on any noncrop area described herein. Review the **SELECTIVE EQUIPMENT** section of this label's **APPLICATION EQUIPMENT AND TECHNIQUES** section for equipment calibration and proper use information.

TANK MIXTURES FOR INDUSTRIAL SITES AND FORESTRY SITE PREPARATIONS

Quali-Pro Glyphosate T&O Plus OUST® (or a registered generic alternative)

Use on industrial sites such as the following: Airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, railroads, roadsides, storage areas, and other similar sites where bare ground is desired.

The Quali-Pro Glyphosate T&O plus Oust mixture may also be used as a site preparation treatment for sites to be planted with the following: Jack pine, loblolly pine, red pine, slash pine, and Virginia pine. When applied as directed herein for **NONCROP USES** under the conditions described, Quali-Pro Glyphosate T&O plus Oust controls annual weeds listed in the **WEEDS CONTROLLED** sections of the label for this product and Oust, and controls or partially controls the perennial weeds listed below.

Make applications of 1 to 2 quarts of Quali-Pro Glyphosate T&O with 2 to 4 ounces of Oust in 10 to 40 gallons of spray solution per acre as a broadcast spray to weeds that are actively growing.

A Quali-Pro Glyphosate T&O plus Oust mixture may be applied by aerial equipment in site prep operations. When applying aerially, use the recommended rates in 5 to 15 gallons of spray solution per acre.

Do not apply this product plus Oust tank mixtures by air in California.

For controlling annual weeds, use the lower rates of Quali-Pro Glyphosate T&O and Oust. For control of the perennial weeds listed herein, use the higher rates of both products. For partial control, use the lower listed rates.

Bahiagrass
Paspalum notatum

Bermudagrass*
Cynodon dactylon

Broomsedge
Andropogon virginicus

Dock, curly
Rumex crispus

Dogfennel
Eupatorium capilliflorum

Fescue, tall
Festuca arundinacea

Johnsongrass**
Sorghum halepense

Poorjoe**
Diodia teres

Quackgrass
Agropyron repens

Trumpetcreeper*
Campsis radicans

Vaseygrass
Paspalum urvillei

Vervain, blue
Verbena hastata

* Suppression at higher rates only.

** Control at lower rates.

Review and carefully observe all cautionary statements and other information appearing on the labels of all of the herbicides used.

TANK MIXTURES FOR NONCROP SITES

When applied as tank mixtures, Quali-Pro Glyphosate T&O controls emerged annual weeds and partially controls emerged perennial weeds listed in this label. When applying as a tank mixture, the residual herbicides listed below (or their registered generic equivalents) will provide preemergence control of those weeds listed in the individual herbicide product labels.

- Quali-Pro Glyphosate T&O with DIURON
- Quali-Pro Glyphosate T&O with KROVAR™ I
- Quali-Pro Glyphosate T&O with KROVAR II
- Quali-Pro Glyphosate T&O with RONSTAR™ 50WP
- Quali-Pro Glyphosate T&O with SIMAZINE, PRINCEP™ CALIBER™ 90
- Quali-Pro Glyphosate T&O with SIMAZINE 4L
- Quali-Pro Glyphosate T&O with SIMAZINE 80W
- Quali-Pro Glyphosate T&O with SURFLAN® 75W
- Quali-Pro Glyphosate T&O with ORYZALIN 4 A.S.

When using residual herbicides in tank mixtures, add an agriculturally approved nonionic surfactant at the rate of 0.5 to 1 percent by volume of spray solution. Review the **MIXING, ADDITIVES** and **APPLICATION INSTRUCTIONS** portion of this label prior to preparing these tank mixtures.

Review and carefully observe all label claims, cautionary statements, recommended usage rates, and all other information on the labels of all of the products used in these mixtures. Use the most restrictive label directions for each product in the mixture.

CONTROL OF EMERGED WEEDS

NOTE: Review this label's **HAND-HELD AND HIGH-VOLUME EQUIPMENT** section for recommended rates when using handgun or back sprayer equipment.

Annual Weeds: Use 1 quart of Quali-Pro Glyphosate T&O per acre in these tank mixtures when weeds are smaller than 6 inches in height, and 1.5 quarts of Quali-Pro Glyphosate T&O per acre when weeds are higher than 6 inches.

Perennial Weeds: For partial control using these tank mixtures, use 2 to 5 quarts of Quali-Pro Glyphosate T&O per acre. Review and follow the recommendations listed in this label's **WEEDS CONTROLLED** section for rate of application and stage of growth information specific to perennial weeds.

PREEMERGENCE WEED CONTROL

Review individual product labels for specific rates, carrier volumes, specific noncrop sites, and precautionary statement information applicable to preemergence weed control.

Mix only the quantity of spray solution that is to be used that same day. These tank mixtures should not be permitted to stand overnight—reduced weed control may result.

INDUSTRIAL, TURF, AND ORNAMENTAL USE RECOMMENDATIONS

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, AND EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE INJURY OR DESTRUCTION IS LIKELY.

Detailed instructions are provided below by site.

Applications may be made to control any weeds listed in the annual, perennial, and woody brush tables, unless otherwise stated. In addition, refer to the **SELECTIVE EQUIPMENT** section of this label.

APPLICATION RATES

APPLICATION	AMOUNT OF QUALI-PRO GLYPHOSATE T&O	SPRAY VOLUME GALLONS/ACRE
<u>Spray-To-Wet</u> Handgun or Backpack	1-4 Percent by Volume***	Spray-to-Wet*
<u>Low-Volume Directed Spray</u> Backpack	9 – 18 Percent by Volume ***	15 to 25 **
Modified High Volume	35 to 7 Percent by Volume ***	40 to 60 **

* For applications made on a spray-to-wet basis, spray coverage should be complete and uniform. Do not spray to the point of runoff.

** For low-volume directed spray applications, coverage should be uniform with a minimum of 50 to 75 percent of the foliage contacted. Coverage of the top one-half of the plant is critical for best results. Low-volume directed applications with backpacks work best when treating weeds and brush that do not exceed 10 feet tall. For taller brush and weeds, high-volume handguns may be modified by reducing nozzle size and spray pressure to produce a low-volume directed spray. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is dense and thick, or where there are multiple sprouts. To achieve best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop.

*** For example, to make a 1-percent solution, use one gallon of Quali-Pro Glyphosate T&O plus 99 gallons of water. Use four gallons plus 96 gallons for a 4-percent solution, etc.

CUT STUMPS

Use Quali-Pro Glyphosate T&O to control woody vegetation by applying to freshly cut stumps or resprouts of undesirable trees. Treatment of cut stumps may be made on any site listed on this label. Use suitable application equipment to ensure coverage of the entire cambium. Trees or resprouts must be cut close to the soil surface. Apply a 50- to 100-percent solution of this product to the freshly-cut surface **immediately after** cutting, as delays in application may result in reduced performance. Applications should be made during periods of active growth and full leaf expansion to ensure optimum results.

DO NOT TREAT CUT STUMPS IF THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO OR SHARE THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees of similar age, height, and spacing may have shared roots. Whether grafted or shared, injury is likely to occur to nontreated stems/trees when one or more trees sharing common roots are treated.

Quali-Pro Glyphosate T&O, when used according to instructions for cut stump applications, will CONTROL, PARTIALLY CONTROL, or SUPPRESS many types of tree species and woody brush. Some of these species are as follows:

Alder
Alnus spp.

Eucalyptus
Eucalyptus spp.

Madrone
Arbutus menziesii

Oak
Quercus spp.

Reed, giant
Arundo donax

Saltcedar
Tamarisk spp.

Sweetgum
Liquidambar styraciflua

Tan Oak
Lithocarpus densiflorus

Willow
Salix spp.

FORESTRY SITE PREPARATION

This product is recommended for the control or partial control of woody brush, trees, and herbaceous weeds in forests, for maintaining logging roads, and for use in preparing or establishing wildlife openings within forest sites.

This product is recommended for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars, and silvicultural nursery sites.

See **APPLICATION RATES** table in **HAND-HELD EQUIPMENT** section of this label for applications using different types of equipment.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When using tank mixes, read and carefully observe the label claims, cautionary statements, and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: When using tank mixes for forestry site preparation, make sure the tank mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Use any recommended rate of this product in a tank mix with the following products (or their registered generic equivalents) for forestry site preparation.

PRODUCT	BROADCAST RATES
Arsenal Applicators Concentrate	2 – 16 fluid oz. / acre
Escort	0.5 – 3.5 oz. / acre
Chopper™	4 – 32 fluid oz. / acre
Garlon 4	1 – 4 qts. / acre
Oust®	1 – 4 oz. / acre
SPRAY-TO-WET RATES	
Arsenal Applicators Concentrate	0.03 – 0.5 percent by volume
LOW-VOLUME DIRECTED SPRAY RATES	
Arsenal Applicators Concentrate	0.1 – 0.5 percent by volume

For controlling dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release.

GENERAL NONCROP AREAS AND INDUSTRIAL SITES

This product may be used in areas such as airports; apartment complexes; Christmas tree farms; ditch banks; dry ditches; dry canals; fencerows; golf courses; industrial sites; lumber yards; manufacturing sites; office complexes; ornamental nurseries; parks; parking areas; petroleum tank farms and pumping installations; railroads; recreational areas; residential areas; roadsides; sod or turf seed farms; schools; storage areas; substations; warehouse areas; other public areas; and similar industrial and noncrop sites.

Grazing Restrictions: Quali-Pro Glyphosate T&O may be used to treat undesirable vegetation in rights-of-way that pass through pastures, rangeland, and forestry sites that are being grazed. For use in tank mix applications, comply with all restrictions appearing on the tank mix product labels.

There are no grazing restrictions for the following labeled applications of Quali-Pro Glyphosate T&O:

- Where the spray can be directed onto undesirable woody trees and brush, such as in handgun spray-to-wet or low-volume directed spray treatments.
- For tree injection or frill applications and for treatments on cut stumps.

For broadcast applications, observe the following restrictions:

- For application rates of greater than 6 but less than 10 quarts per acre, no more than 15 percent of the available grazing area may be treated.
- For application rates of no more than 6 quarts per acre, no more

- than 25 percent of the available grazing area may be treated.
- All restrictions outlined above apply to lactating dairy animals. No other restrictions apply to lactating dairy animals.

These recommendations do not apply to rangeland outside of rights-of-way.

General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general noncrop areas using any application equipment described in this label. Use to trim-and-edge around objects in noncrop sites, for spot treatment of unwanted vegetation, and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. Use prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used to maintain bare ground as weeds emerge.

TANK MIXTURES: This product may be used in tank mixes with the following products (or their registered generic equivalents) if the tank mix product is registered for use on such noncrop sites. Refer to the tank mix product's labels for approved noncrop sites and application rates.

Arsenal™	Pendulum™ 3.3 EC
Barricade™ 65WG	Pendulum WDG
Clarity	Plateau™
Diuron	Princep™ DF
Endurance™	Princep™ Liquid
Escort®	Ronstar™ 50WP
Garlon™ 3A	Sahara™
Garlon 4	Simazine
Hyvar X	Spike 80DF
Karmex™ DF	Surflan®
Krovar™ I DF	Telar™
Manage®	Vanquish™
Oust®	2,4-D

Do not apply this product with dicamba tank mixtures by air in California.

Brush Control Tank Mixtures

Tank mixtures of Quali-Pro Glyphosate T&O may be used to increase the spectrum of control for herbaceous weeds, trees, and woody brush. When tank mixing, read and carefully observe all label claims, cautionary statements and information on the labels of all products used in the mixture. Use according to the most restrictive precautionary statements for each product. Any recommended rate of this product may be used in a tank mix as stated in this label.

For controlling herbaceous weeds, the lower recommended tank mixture rates should be used. For controlling dense stands or tough-to-control woody brush and trees, the higher recommended rates should be used.

NOTE: For side-trimming treatments, this product should be used

alone or in tank mixture with Garlon 4 (or a registered generic equivalent). For all products listed in the chart below, registered generic equivalents may also be used.

PRODUCT	BROADCAST RATES
Arsenal 2WSL	6 – 32 fluid oz. / acre
Escort	1 – 2 oz. / acre
Garlon 3A*, Garlon 4	1 – 4 qts. / acre
SPRAY-TO-WET RATES	
Arsenal 2WSL	0.06 – 0.12 percent by volume
Escort	1 – 2 oz. / acre
LOW-VOLUME DIRECTED SPRAY RATES	
Arsenal 2WSL	0.1 – 0.5 percent by volume
Escort	1 – 2 oz. / acre

* Ensure that Garlon 3A (or a registered generic equivalent) is mixed thoroughly with water in accordance with label directions before adding Quali-Pro Glyphosate T&O. Spray mixture must be agitating when this product is added to avoid spray compatibility problems.

Chemical Mowing - Perennials

Quali-Pro Glyphosate T&O will suppress perennial grasses listed in this section to serve as a substitute for mowing. When treating tall fescue, fine fescue, orchardgrass, quackgrass, or reed canarygrass covers, use 8 fluid ounces of this product per acre. When treating Kentucky bluegrass, use 6 fluid ounces of this product per acre. Apply these treatments in 10 to 20 gallons of spray solution per acre.

Use only where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing - Annuals

For growth suppression of annual grasses such as annual ryegrass, wild barley, and wild oats growing in coarse turf in industrial areas or on roadsides, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Make applications when annual grasses are actively growing and before seedheads are in the boot stage of development. Note that treatments may injure the desired grasses.

Bromus Species and Medusahead in Industrial Sites, Pastures and Rangelands

Bromus species: Quali-Pro Glyphosate T&O may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial sites, rangeland and pastures. Apply 8 to 16 fluid ounces of Quali-Pro Glyphosate T&O per acre on a broadcast basis.

Treatment should coincide with early seedhead emergence of the most mature plants for best results. Delaying application until this growth stage will maximize the emergence of other weedy grass flushes. Apply Quali-Pro Glyphosate T&O to the same site each

year until seed banks are depleted and the desirable perennial grasses become reestablished on the site.

Medusahead: For treating medusahead, apply 16 fluid ounces of Quali-Pro Glyphosate T&O per acre as soon as plants are actively growing, and prior to the 4-leaf stage. Apply in the fall or spring.

Use ground or aerial equipment when applying to brome and medusahead. Use fixed-wing or helicopter equipment when applying by air, and apply in 2 to 10 gallons of water per acre. When using ground equipment, apply in 10 to 20 gallons of water per acre. When applied in accordance with this label, there are no grazing restrictions.

Dormant Turfgrass

Quali-Pro Glyphosate T&O may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Apply only when turf is dormant and before spring greenup.

Apply 8 to 64 fluid ounces of Quali-Pro Glyphosate T&O per acre. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where some temporary injury or discoloration can be tolerated and where bermudagrass or bahiagrass are desirable ground covers.

Applying treatments in excess of 16 fluid ounces per acre may cause injury or delayed greenup in highly maintained areas, such as golf courses and lawns. Do not apply tank mixtures of Quali-Pro Glyphosate T&O plus Oust or Outrider in highly maintained turfgrass areas. For further uses, refer to the **ROADSIDES** section below, which provides rates for treating dormant bermudagrass and bahiagrass.

Actively Growing Bermudagrass

Quali-Pro Glyphosate T&O may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. In highly maintained turfgrass areas, do not apply more than 16 fluid ounces of this product per acre, and do not apply tank mixtures of this product plus Oust. For additional uses, refer to the **ROADSIDES** section below, which gives rates for actively growing bermudagrass treatments. Use only in areas where temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

Use Quali-Pro Glyphosate T&O to control most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, determine if any regrowth has occurred from escaped underground plant parts before planting or sodding. Where repeat treatments are necessary, ensure that sufficient regrowth has occurred prior to application. Summer or fall applications provide the best control for warm-season grasses such as bermudagrass. If existing vegetation is growing under mowed turfgrass management, apply Quali-Pro Glyphosate T&O after omitting a minimum of one regular mowing to allow sufficient growth for good interception of the spray.

Before treatment, do not disturb soil or underground plant parts. Delay tillage or renovation techniques such as vertical mowing, coring, or slicing for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted once the above procedures have been followed.

For instances of unwanted vegetation growing in existing turfgrass, hand-held equipment may be used for spot treatment. To control sod remnants or other unwanted vegetation after sod is harvested, broadcast or hand-held equipment may be used.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks after applying this product.

DORMANT RANGELAND

Quali-Pro Glyphosate T&O controls or suppresses many weeds in dormant rangeland, including the following: Downy brome, cheat grass, cereal rye, medusahead rye, and jointed goatgrass.

In early spring when weeds have greened up but desirable grasses (such as crested and tall wheatgrass) are still truly dormant, apply Quali-Pro Glyphosate T&O at the rate of 8 to 16 ounces per acre.

Some slight discoloration of the desirable grasses may result, but they will regreen and regrow under moist soil conditions, as the effects of Quali-Pro Glyphosate T&O dissipate.

For dormant rangeland applications, do not use additional surfactant or ammonium sulfate.

HABITAT MANAGEMENT

Habitat Restoration and Maintenance

Use Quali-Pro Glyphosate T&O to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Treatments can be made to allow recovery of native plant species, before planting desirable native species, and for similar broad-spectrum vegetation control requirements. Use spot treatments to selectively remove unwanted plants for habitat management and enhancement. However, extreme care should be taken to ensure Quali-Pro Glyphosate T&O does not contact desirable plants.

Wildlife Food Plots

Quali-Pro Glyphosate T&O may be used as a site preparation treatment prior to planting wildlife food plots. Users may plant any wildlife food species after applying this product, or native species may be allowed to repopulate the area. If tillage is necessary to prepare a seedbed, wait 7 days after application before tillage to facilitate translocation into underground plant parts.

INJECTION AND FRILL (WOODY BRUSH AND TREES)

Use Quali-Pro Glyphosate T&O to control woody brush and trees by injection or frill applications. When applying this product, use suitable equipment that penetrates into the living tissue. Use the equiv-

alent of 0.04 fluid ounce (1 ml) of Quali-Pro Glyphosate T&O per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50- to 100-percent concentration of Quali-Pro Glyphosate T&O either to a continuous frill around the tree or as cuts that are evenly spaced around the tree and below all branches. As a tree's diameter increases in size, apply diluted material to a continuous frill or more closely spaced cuttings for better results. Do not use application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, use a 100-percent concentration of this product and make the frill or cuts at an oblique angle to produce a cupping effect. For best results, make applications during periods of active growth and after full leaf expansion.

ORNAMENTALS, PLANT NURSERIES, AND CHRISTMAS TREES

UNLESS OTHERWISE SPECIFIED, THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES.

Exercise care to avoid contact of spray, drift, or mist with foliage or green bark of established ornamental species.

Post-Directed, Trim-and-Edge

When applied as instructed herein, Quali-Pro Glyphosate T&O controls undesirable vegetation listed on this label prior to planting, within and around greenhouses and shadehouses, and as a post-directed spray around Christmas trees.

Quali-Pro Glyphosate T&O can be used as a post-directed spray around established woody ornamental species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, hollies, jojoba, lilac, magnolia, maple, oak, pine, poplar, privet, spruce, and yew. Quali-Pro Glyphosate T&O is also effective when used to trim and edge around trees, buildings, roads and sidewalks, potted plants, and other objects in nursery settings.

Review the **WEEDS CONTROLLED** sections of this label for specific instructions and rates of application for controlling various perennial and annual weeds. For Christmas trees, if making multiple applications, Do not use more than 10.6 quarts of Quali-Pro Glyphosate T&O per acre per year.

NOTE: Protect desirable plants from the spray solution by using shields or coverings made of impermeable material, such as cardboard.

Site Preparation

Use Quali-Pro Glyphosate T&O prior to planting any ornamental, nursery, or Christmas tree species. Exercise extreme caution to keep spray off nontarget plants during site preparation applications.

Wiper Applications

Quali-Pro Glyphosate T&O may be applied through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established eucalyptus or poplar trees. Review the **SELECTIVE EQUIPMENT** section of this label for additional information about the correct use of wiper applicators.

Greenhouse/Shadehouse

Control weeds growing in and around greenhouses and shadehouses by using Quali-Pro Glyphosate T&O. When doing so, turn off air circulation fans, and remove desirable vegetation prior to application.

Christmas Tree Plantations in Oregon and Washington ONLY

NOTE: If this product is improperly applied, it has the potential to cause severe injury to Christmas trees. Follow all labeled directions.

Quali-Pro Glyphosate T&O may be applied as a broadcast spray over established Christmas trees. To prevent drift onto nearby desirable crops or vegetation, ensure that adequate buffers are maintained. For additional application precautions, read the entire section titled **APPLICATION EQUIPMENT AND TECHNIQUES** on the Quali-Pro Glyphosate T&O main label.

The following Christmas tree species are approved for this application:

- Douglas Fir (*Pseudotsuga menziesii*)
- Fir species (*Abies* spp.)
- Spruce species (*Picea* spp.)

Do not apply this product until trees have completed at least a full growing season since planting or transplanting. Do not apply within 1 full year prior to tree harvest.

In the fall, applications may only be made after the formation of final conifer resting buds. Final resting buds must be in the dormant stage and fully hardened. If applications are made at any other time, unacceptable Christmas tree injury may occur.

Avoid spray pattern overlap, as injury may result.

Apply 1 quart of Quali-Pro Glyphosate T&O per acre in 5 to 30 gallons of water per acre.

NOTE: Adding surfactants, additives containing surfactants, or any other additives to this product may result in severe Christmas tree injury.

In some areas, this product may be used at rates from 1 to 2 quarts per acre. Contact your local Makhteshim Agan of North America, Inc. representative for specific recommendations if you require rates that exceed 1 quart per acre.

Drift control additives are not recommended, as they may increase Christmas tree injury. Using other herbicides tank mixed with Quali-Pro Glyphosate T&O is not recommended, as Christmas trees could be severely injured.

SILVICULTURAL SITES AND RIGHTS-OF-WAY

DO NOT USE QUALI-PRO GLYPHOSATE T&O AS AN OVER-THE-TOP BROADCAST SPRAY IN SILVICULTURAL NURSERIES.

When applied as directed for noncrop uses under the conditions

described herein, Quali-Pro Glyphosate T&O will control the undesirable vegetation listed on this label. Quali-Pro Glyphosate T&O will also suppress or control the undesirable vegetation listed on this label when it is applied at the recommended rates for release of established coniferous species listed on this label.

See the **WEEDS CONTROLLED** sections of this label for specific rates of application and instructions for controlling various brush, annual and perennial weeds. See the **Conifer Release** part of this section of the label for specific rates of application for release of listed coniferous species.

Do not apply more than 10.6 quarts of Quali-Pro Glyphosate T&O per acre per year where repeat applications are necessary.

Aerial Application

For silvicultural site preparation, conifer release, and rights-of-way treatments, this product may be applied using aerial spray equipment. Review the **APPLICATION EQUIPMENT AND TECHNIQUES** part of the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for detailed information on how to apply this product aerially.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHT-OF-WAY SITES IN CALIFORNIA.

Site Preparation

After preplant applications of Quali-Pro Glyphosate T&O, any silvicultural species may be planted.

Postdirected Spray

Use as a spray on the foliage of undesirable vegetation in established silvicultural sites. Exercise care to avoid contact of spray, mist, or drift with foliage or green bark of desirable species.

Conifer Release

For conifer release, apply only where conifers have been established for a period of one year or more. Do not disturb vegetation prior to treatment or until visual symptoms appear after application. Symptoms of treatment are slow to appear, especially in woody species that have been treated in late fall. Injury may occur to conifers that have been treated for release, especially where spray patterns overlap, where the higher rates are applied, or when applications are made during periods of active conifer growth. **For conifer release applications, do not use additional surfactant.**

Make applications after formation of final conifer resting buds in the fall, or prior to initial bud swelling in the spring. Some autumn colors on undesirable deciduous species are acceptable so long as no major leaf drop has occurred. Use the rates listed below for conifer release to control or partially control those weeds described in the **WEEDS CONTROLLED** sections of this label.

For release of the conifer species listed below:

Douglas Fir
Pseudotsuga menziesii

Fir
Abies spp.

Hemlock
Tsuga spp.

Pines*
Pinus spp.

Spruce
Picea spp.

*Includes all species except eastern white pine, slash pine, or loblolly pine.

Apply 1.5 to 2 quarts of Quali-Pro Glyphosate T&O per acre except for west of the crest of the Cascade Mountains in Washington and Oregon. When applying in spring west of the crest of the Cascade Mountains, apply 1 quart of Quali-Pro Glyphosate T&O per acre prior to conifer bud swell for controlling annual weeds. When applying in fall west of the crest of the Cascade Mountains in Washington and Oregon, apply 1 to 1.5 quarts of this product per acre prior to any major leaf drop of deciduous species.

Apply 1 quart of this product per acre for release of western hemlock.

For release of these conifer species:

Loblolly pine
Pinus taeda

Eastern white pine
Pinus strobes

Slash pine
Pinus elliotii

Late-Season Application

During early autumn, apply 1.5 to 2 quarts of Quali-Pro Glyphosate T&O in at least 5 gallons of spray solution per acre. Applying before September 1 or when conditions are conducive to rapid growth of conifers will create the possibility of increased injury in the form of needle and/or tip burn. Injury may decrease with later applications. Some autumn colors are acceptable at the time of treatment. Treat before frost or leaf drop of undesirable plants. Applications made in accordance with label directions will release loblolly pine, slash pine and eastern white pine by reducing competition from the following species:

Ash
Fraxinus spp.

Cherry:
Black
Prunus serotina

Pin
Prunus pensylvanica

Elm
Ulmus spp.

Hawthorn
Crataegus spp.

Locust, black
Robinia pseudoacacia

Maple, red
Acer rubra

Oak:
Black
Quercus velutina

Post
Quercus stellata

Southern Red
Quercus falcata

White
Quercus alba

Persimmon
Diospyros spp.

Poplar, yellow
Liriodendron tulipifera

Sassafras
Sassafras albidum

Sourwood
Oxydendrum arboreum

Sumac:
Poison
Rhus vernix

Smooth
Rhus glabra

Winged
Rhus copallina

Sweetgum
Liquidambar styraciflua

Treat only the sites where woody brush and trees listed in this label make up the majority of the undesirable species.

Conifer Release From Herbaceous Weeds: Quali-Pro Glyphosate T&O Plus Oust Tank Mixtures

To release **loblolly pines** from herbaceous weeds, tank mixtures of Quali-Pro Glyphosate T&O plus Oust (or a registered generic alternative) will control annual weeds listed in the **WEEDS CONTROLLED**

sections of this and the Oust label, and will partially control those perennial weeds that are listed below.

Mix and apply 16 to 24 fluid ounces of Quali-Pro Glyphosate T&O with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Apply to weeds that are actively growing as a broadcast spray over the top of the young loblolly pines.

If desired, this tank mixture may be applied using aerial equipment. When using aerial equipment, use the recommended rate in 5 to 15 gallons of spray solution per acre.

Do not apply Quali-Pro Glyphosate T&O plus Oust tank mixtures by air in California.

For controlling the annual weeds listed below that are 12 inches in height (or runner length on annual vines), use the lower rates of both Quali-Pro Glyphosate T&O and Oust. When annual weeds are in more advanced stages of growth and are approaching flower or seed formation, use the higher rates of both products.

Use the higher rates of both products for partial control of the perennial weeds listed below. Use the lower rates to suppress growth.

Bahiagrass
Paspalum notatum

Broomsedge
Andropogon virginicus

Dock, curly
Rumex crispus

Dogfennel
Eupatorium capilliflorum

Fescue, tall
Festuca arundinacea

Johnsongrass*
Sorghum halepense

Poorjoe*
Diodia teres

Trumpet creeper**
Campsis radicans

Vaseygrass
Paspalum urvillei

Vervain, blue
Verbena hastata

* Provides control at higher rates.

** Provides suppression only at higher rates.

If treatment takes place when young trees are under stress from drought, flood water, insects or disease, pine damage may occur or can be accentuated.

Review and carefully observe the cautionary statements and all other information appearing on the labels of all of the herbicides used.

NOTE TO USER: Do not use Quali-Pro Glyphosate T&O in areas where adverse impact on federally designated endangered or threatened plants or aquatic species is likely.

Before treatment, users must determine that no such species are located in or immediately adjacent to the area of application.

PARKS, RECREATIONAL, AND RESIDENTIAL AREAS

Quali-Pro Glyphosate T&O may be applied in parks, recreational, and residential areas using any application equipment described in this label. It may be used to trim-and-edge around trees, paths, fences, and around buildings, sidewalks, and other objects. This product may be used for spot treatment of unwanted vegetation, and may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to laying asphalt or beginning construction projects, or prior to planting an area to ornamentals, flowers, or turfgrass (sod or seed).

All of the instructions in the section titled **GENERAL NONCROP AREAS AND INDUSTRIAL SITES** apply to park and recreation areas.

RAILROADS

All of the instructions in the section titled **GENERAL NONCROP AREAS AND INDUSTRIAL SITES** apply to railroads.

Bare Ground, Ballast and Shoulders, Crossings, Spot Treatments

Use Quali-Pro Glyphosate T&O to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be needed, as weeds emerge, to maintain bare ground. Quali-Pro Glyphosate T&O may also be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. Use up to 80 gallons of spray solution per acre for crossing applications

TANK MIXTURES: Tank mix Quali-Pro Glyphosate T&O with the following products (or their registered generic equivalents) only if the specific product is registered for ballast, shoulder, spot, bare ground, and crossing treatments:

Arsenal	Krovax I
DFClarity	Oust
Diuron	Sahara
Escort	Spike™
Garlon 3A	Telar
Garlon 4	Vanquish
Hvyar™ X	2,4-D

Brush Control

Control woody brush and trees on railroad rights-of-way by applying Quali-Pro Glyphosate T&O as follows: Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boom-less nozzles. A maximum of 80 gallons of spray solution per acre

may be used. Apply a 0.75- to 2-percent solution of Quali-Pro Glyphosate T&O when using high-volume spray-to-wet applications. When using low-volume directed sprays for spot treatment, apply a 5- to 10-percent solution of this product

TANK MIXTURES: For enhanced control of woody brush and trees, mix Quali-Pro Glyphosate T&O with the following products (or their registered generic equivalents):

Arsenal	Garlon 4
Escort	Tordon™ K
Garlon 3A	

Bermudagrass Release

Quali-Pro Glyphosate T&O may be used for control or partial control of many annual and perennial weeds for effective release of actively growing bermudagrass. Mix 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length); use the higher rate as weeds grow or as they approach flower or seedhead formation. Applying Quali-Pro Glyphosate T&O at these rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpetcreeper
Fescue, tall	Vaseygrass

TANK MIXTURES: Quali-Pro Glyphosate T&O may be tank mixed with Oust (or a registered generic equivalent) by mixing no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust per acre. To control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label, use the lower rates of each product. The higher rates should be used as annual weeds increase in size and as they approach the flower or seedhead stages. These rates will also provide partial control of these perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestem, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpetcreeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use this product only on well-established bermudagrass. Although bermudagrass injury may result from the treatment, regrowth will occur under moist conditions. Do not repeat applications in the same season as severe injury may occur.

ROADSIDES

All of the instructions provided in the **GENERAL NONCROP AREAS AND INDUSTRIAL SITES** section apply to roadsides.

TANK MIXTURES: Quali-Pro Glyphosate T&O may be tank mixed with the following products (or their registered generic equivalents) if the specific product is registered for shoulder, guardrail, spot and

bare ground treatments:

Clarity	Princep Liquid
Diuron	Ronstar 50WP
Endurance	Sahara
Escort	Simazine
Krovat I DF	Surflan
Oust	Telar
Pendulum 3.3 EC	Vanquish
Pendulum WDG	2,4-D
Princep DF	

Refer to the **GENERAL NONCROP AREAS AND INDUSTRIAL SITES** section of this label for general tank mixing instructions.

Shoulder Treatments

Quali-Pro Glyphosate T&O may be used on road shoulders using boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails and Other Obstacles to Mowing

Use Quali-Pro Glyphosate T&O to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

Use Quali-Pro Glyphosate T&O as a spot treatment to control unwanted vegetation growing along roadsides.

Release of Bermudagrass or Bahiagrass

Dormant Applications

Quali-Pro Glyphosate T&O may be used to control or partially control tall fescue and many winter annual weeds for effective release of dormant bermudagrass or bahiagrass. Apply only when turf is dormant and before spring greenup. This product may also be tank mixed with Oust or Outrider for residual control. **NOTE:** Tank mixtures of this product with Oust may delay greenup.

To obtain best results on winter annuals, apply when plants are in an early growth stage (below 6 inches in height) after most have germinated. To obtain best results on tall fescue, apply when fescue is at or beyond the 4- to 6-leaf stage.

Apply 8 to 64 fluid ounces of Quali-Pro Glyphosate T&O in a tank mixture with 0.75 to 1.3 ounces of Outrider herbicide per acre. Read and observe all label directions for Outrider herbicide.

Apply 8 to 64 fluid ounces of Quali-Pro Glyphosate T&O per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust. Apply recommended rates in 10 to 40 gallons of water per acre. This product should only be used in areas where bermudagrass or bahiagrass are desirable ground covers, and where some temporary injury or discoloration can be tolerated. To minimize injury and to avoid delays in greenup, add no more than 1 ounce of Oust per acre on bermudagrass and no more than 0.5 ounce of Oust per acre on bahiagrass. Avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

Use Quali-Pro Glyphosate T&O to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 1 to 3 pints of Quali-Pro Glyphosate T&O in 10 to 40 gallons of spray solution per acre. The lower rate should be used when treating annual weeds below 6 inches in height (or runner length). The higher rate should be used as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the perennial species listed below:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpet creeper
Fescue, tall	Vaseygrass

TANK MIXTURES: Quali-Pro Glyphosate T&O may be tank mixed with Outrider to control or partially control Johnsongrass and other weeds listed on the Outrider label. Mix and apply 8 to 32 fluid ounces of this product with 0.75 to 1.3 ounces of Outrider. The higher rates of both products should be used to control perennial weeds or annual weeds greater than 6 inches in height.

Quali-Pro Glyphosate T&O may be tank mixed with Oust. If tank mixed, do not exceed 1 to 2 pints of this product with 1 to 2 ounces of Oust per acre. The lower rates of each product should be used to control those annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. The higher rates should be used as annual weeds increase in size and as they approach the flower or seedhead stages. These rates will also provide partial control of the perennial weeds listed below:

Bahiagrass	Fescue, tall
Bluestem, silver	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Trumpet creeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

Use only on bermudagrass that is well established. However, bermudagrass injury may result from treatment, but regrowth will occur under moist conditions. Do not repeat applications of the tank mix in the same season, as severe injury may occur.

Actively Growing Bahiagrass

To suppress vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of approximately 3 to 4 inches. This application must be made before seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, then apply 2 to 4 fluid ounces per acre approximately 45 days later. Do not exceed 2 applications per year.

TANK MIXTURES: Quali-Pro Glyphosate T&O may be used to control or partially control johnsongrass and other weeds listed on the Outrider label in actively growing bahiagrass. Mix and apply 1.5 to 5 fluid ounces of Quali-Pro Glyphosate T&O with 0.75 to 1.3

ounces of Outrider per acre. The higher rates should be used for control of perennial weeds or annual weeds that exceed 6 inches in height. Use only on bahiagrass that is well established.

A tank mixture of Quali-Pro Glyphosate T&O plus Oust may be used. Mix and apply 6 fluid ounces of this product plus 0.25 ounce of Oust per acre 1 to 2 weeks after an initial spring mowing. Do not exceed one application per year.

UTILITY SITES

Quali-Pro Glyphosate T&O is recommended for use along electrical power, pipeline and telephone rights-of-way, and also in other sites associated with these rights-of-way, including substations, roadsides, railroads, or similar rights-of-way that run in conjunction with utilities.

Quali-Pro Glyphosate T&O may be used in preparing or establishing wildlife openings within utility sites, maintaining access roads, and for side trimming along utility rights-of-way.

TANK MIXTURES: Quali-Pro Glyphosate T&O used in tank mixtures increases the spectrum of control for herbaceous weeds, woody brush, and trees.

IMPORTANT: When tank mixing, read and carefully observe all label claims, cautionary statements, and other information on the labels of all products used. Products should be used according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of Quali-Pro Glyphosate T&O may be used in a tank mix.

Use the lower recommended tank mixture rates for control of herbaceous weeds. Use the higher recommended rates for control of dense stands or tough-to-control woody brush and trees. For all products listed in the chart below, registered generic equivalents may also be used.

NOTE: For side-trimming treatments, this product should be used alone or in tank mixture with Garlon 4.

PRODUCT	*BROADCAST RATES	USE SITES
Arsenal 2WSL	6 – 32 fluid oz. / acre	Utility Sites
Escort	1 – 2 oz. / acre	Utility Sites
Garlon 3A**, Garlon 4	1 – 4 quarts / acre	Utility Sites / Side Trimming
Oust	1 – 4 oz. / acre	Utility Sites
	SPRAY-TO-WET RATES	USE SITES
Arsenal 2WSL	0.06 – 0.1 percent by volume	Utility Sites
Escort	1 – 2 oz. / acre	Utility Sites
	LOW-VOLUME DIRECTED SPRAY RATE	USE SITES
Arsenal 2WSL	0.1 – 0.5 percent by vol- ume	Utility Sites
Escort	1 – 2 oz. / acre	Utility Sites

*Or a registered generic equivalent.

** Garlon 3A (or the registered generic equivalent) must be thoroughly mixed with water according to label directions before adding

Quali-Pro Glyphosate T&O. To avoid spray compatibility problems, have spray mixture agitating at the time this product is added.

Bare Ground, Trim-and-Edge

Use Quali-Pro Glyphosate T&O in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation, and to eliminate unwanted weeds that are growing in established ornamental plantings or shrub beds. This product may be used before planting a utility site with ornamentals, flowers, or turf-grass (sod or seed), or prior to beginning construction projects.

To maintain bare ground, repeat applications of this product may be used as weeds emerge.

TANK MIXTURES: Quali-Pro Glyphosate T&O may be tank mixed with the products (or their registered generic equivalents) listed below. Refer to and observe all instructions in these products' labels for approved noncrop sites and application rates.

Arsenal	Plateau™
Banvel	Princep™ DF
Barricade™ 65WG	Princep™ Liquid
Diuron	Ronstar™ 50WP
Endurance™	Sahara™
Escort	Simazine
Garlon 3A	Surflan®

WEEDS CONTROLLED—INDUSTRIAL, TURF, ORNAMENTAL USES

If weed growth is heavy or dense, or if weeds are growing in an undisturbed (non-cultivated) area, always use the higher rate of this product per acre within the recommended range.

Reduced performance may result when treating weeds that are heavily covered with dust. Allow regrowth to occur prior to treatment of weeds that have been mowed, grazed, or cut.

Review the following label sections for recommended rates for the control of perennial and annual weeds, woody brush, and trees. For difficult-to-control perennial weeds, woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, obtain improved results by using Quali-Pro Glyphosate T&O at the rate of 5 to 10 quarts per acre.

ANNUAL WEEDS

Use 1 quart of Quali-Pro Glyphosate T&O per acre if weeds are less than 6 inches in height or runner length. Use 1.5 quarts to 4 quarts per acre if weeds exceed 6 inches in height or runner length, or when weeds are growing under stressed conditions.

When using spray-to-wet applications, apply a 0.5-percent solution of Quali-Pro Glyphosate T&O to weeds less than 6 inches in height or runner length. Apply before seedhead formation in grass, or prior to bud formation in broadleaf weeds. For annual weeds that exceed 6 inches in height, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. The higher rate should be

used for tough-to-control species or for weeds over 24 inches tall.

WEED SPECIES

Annoda, spurred
Barley*
Barnyardgrass*
Bittercress*
Black nightshade*
Bluegrass, annual*
Bluegrass, bulbous*
Bassia, fivehook
Brome, downy*
Brome, Japanese*
Browntop panicum*
Buttercup*
Carolina foxtail*
Carolina geranium
Castor bean
Cheatgrass*
Cheeseweed (*Malva parviflora*)
Chervil*
Chickweed*
Cocklebur*
Copperleaf, hophornbeam
Corn*
Corn speedwell*
Crabgrass*
Dwarf dandelion*
Eastern manna grass*
Eclipta*
Fall panicum*
Falsedandelion*
Falseflax, smallseed*
Fiddleneck
Field pennycress*
Filaree
Fleabane, annual*
Fleabane, hairy (*Conyza bonariensis*)*
Fleabane, rough*
Florida pusley
Foxtail*
Goatgrass, jointed*
Goosegrass
Grain sorghum (milo)*
Groundsel, common*
Hemp sesbania
Henbit
Horseweed/Marestail (*Conyza canadensis*)**
Itchgrass*
Johnsongrass, seedling
Junglerice
Knotweed
Kochia
Lambs quarters*
Little barley*
London rocket*

WEED SPECIES (Cont.)

Mayweed
Medusahead*
Morningglory (*Ipomoea* spp.)
Mustard, blue*
Mustard, tansy*
Mustard, tumble*
Mustard, wild*
Oats
Pigweed*
Plains/Tickseed coreopsis*
Prickly lettuce*
Puncturevine
Purslane, common
Ragweed, common*
Ragweed, giant
Red rice
Russian thistle
Rye*
Ryegrass*
Sandbur, field*
Shattercane*
Shepherd's purse*
Sicklepod
Signalgrass, broadleaf*
Smartweed, ladythumb*
Smartweed, Pennsylvania*
Sowthistle, annual
Spanishneedles
Speedwell, purslane*
Sprangletop*
Spurge, annual
Spurge, prostrate*
Spurge, spotted*
Spurry, umbrella*
Starthistle, yellow
Stinkgrass*
Sunflower*
Tea weed/ Prickly sida
Texas panicum*
Velvetleaf
Virginia copperleaf
Virginia pepperweed*
Wheat*
Wild oats*
Witchgrass*
Woolly cupgrass*
Yellow rocket

* When applying with field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles), these species will be controlled or partially controlled using 1 pint of Quali-Pro Glyphosate T&O per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Nozzles that ensure thorough coverage of foliage should be used, and treatment should be made when weeds are in an early growth stage.

**Resistance to some biotypes has been observed. If users

encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

PERENNIAL WEEDS

For best results, apply treatment after perennial weeds reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For use on nonflowering plants, best results are obtained when plants reach a mature stage of growth. In many situations, treatments must be made prior to these growth stages. Under these conditions, use the higher application rate within the recommended range.

When using spray-to-wet treatments by hand-held equipment, thorough coverage is essential. When using hand-held equipment for low-volume directed spot treatments, apply a 5- to 10-percent solution of Quali-Pro Glyphosate T&O.

Allow a minimum of 7 days after application before tillage, unless otherwise specified.

WEED SPECIES	APPLICATION RATE (Quarts/Acre)	HAND-HELD % SOLUTION
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2 - 4	1 - 2
Bahiagrass	3 - 5	2
Beachgrass, European (Ammophila Arenaria)	—	5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4 - 5	2
Bluegrass, Kentucky	2	2
Blueweed, Texas	4 - 5	2
Brackenfern	3 - 4	1 - 1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf	—	2
Canarygrass, reed	2 - 3	2
Cattail	3 - 5	2
Clover; red, white	3 - 5	2
Cogongrass	3 - 5	2
Dallisgrass	3 - 5	2
Dandelion	3 - 5	2
Dock, curly	3 - 5	2
Dogbane, hemp	4	2
Fescue (except tall)	3 - 5	2
Fescue, tall	1 - 3	2
German ivy	2 - 4	1 - 2
Guineagrass	3	1
Horsenettle	3 - 5	2
Horseradish	4	2
Iceplant	2	1.5 - 2
Jerusalem artichoke	3 - 5	2
Johnsongrass	2 - 3	1
Kikuyugrass	2 - 3	2
Knapweed	4	2
Lantana	—	1 - 1.25

WEED SPECIES

	APPLICATION RATE (Quarts/Acre)	HAND-HELD % SOLUTION
Lespedeza	3 - 5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3 - 5	2
Napiergrass	3 - 5	2
Nightshade, silverleaf	2	2
Nutsedge; purple, yellow	3	1 - 2
Orchardgrass	2	2
Pampasgrass	3 - 5	1.5 - 2
Paragrass	3 - 5	2
Pepperweed, perennial	4	2
Phragmites*	3 - 5	1 - 2
Poison hemlock	2 - 4	1 - 2
Quackgrass	2 - 3	2
Redvine*	2	2
Reed, giant	4 - 5	2
Ryegrass, perennial	2 - 3	1
Smartweed, swamp	3 - 5	2
Spurge, leafy*	—	2
Sweet potato, wild*	—	2
Thistle, artichoke	2 - 3	1 - 2
Thistle, Canada	2 - 3	2
Timothy	2 - 3	2
Torpedograss*	4 - 5	2
Trumpet creeper*	2 - 3	2
Vaseygrass	3 - 5	2
Velvetgrass	3 - 5	2
Wheatgrass, western	2 - 3	2

* Partial Control

WOODY BRUSH AND TREES

Quali-Pro Glyphosate T&O should be applied after full leaf expansion, unless otherwise directed. For larger plants and/or dense areas of growth, use the higher rate. On vines, the higher rate should be used on plants that have reached the woody stage of growth. For best results, apply in late summer or fall after fruit formation.

For best results in arid areas, apply in the spring to early summer when brush species are at high moisture content and are flowering.

When applying with hand-held equipment, ensure thorough coverage for spray-to-wet treatments. When using hand-held equipment for low-volume directed-spray spot treatments, use a 5- to 10-percent solution of this product.

For fall treatments, symptoms may not appear prior to frost or senescence.

Allow a minimum of 7 days after application before tillage, mowing or removal, unless otherwise specified. Repeat treatments may be needed to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable if no major leaf drop has occurred. Reduced performance may result from fall treatments made following a frost.

WEED SPECIES	APPLICATION RATE (Quarts/Acre)	HAND-HELD % SOLUTION
Alder	3-4	1-1.5
Ash*	2-5	1-2
Aspen, quaking	2-3	1-1.5
Bearclover (Bearmat)*	2-5	1-2
Beech*	2-5	1-2
Birch	2	1
Blackberry	3-4	1-1.5
Blackgum	2-5	1-2
Bracken	2-5	1-2
Broom; French, Scotch	2-5	1.5-2
Buckwheat, California*	2-4	1-2
Cascara*	2-5	1-2
Catsclaw*	—	1-1.5
Ceanothus*	2-5	1-2
Chamise*	2-5	1
Cherry; bitter, black, pin	2-3	1-1.5
Coyote brush	3-4	1.5-2
Dewberry	2-5	1
Dogwood*	2-5	1-2
Elderberry	2	1
Elm*	2-5	1-2
Eucalyptus	—	2
Gorse*	2-5	1-2
Hasardia*	2-4	1-2
Hawthorn	2-3	1-1.5
Hazel	2	1
Hickory*	2-5	1-2
Honeysuckle	3-4	1-1.5
Hornbeam, American*	2-5	1-2
Kudzu	4	2
Locust, black*	2-4	1-2
Madrone resprouts*	—	2
Manzanita*	2-5	1-2
Maple, red	2-4	1-1.5
Maple, sugar	—	1-1.5
Monkey flower*	2-4	1-2
Oak; black, white*	2-4	1-2
Oak, post	3-4	1-1.5
Oak; northern, pin	2-4	1-1.5
Oak, Scrub*	2-4	1-1.5
Oak; southern red	2-3	1-1.5
Peppertree, Brazilian (Florida Holly)*	2-5	1-2
Persimmon*	2-5	1-2
Pine	2-5	1-2
Poison ivy	4-5	2
Poison oak	4-5	2
Poplar, yellow*	2-5	1-2
Redbud, eastern	2-5	1-2
Rose, multiflora	2	1
Russian olive*	2-5	1-2
Sage, black	2-4	1
Sage, white*	2-4	1-2
Sage brush, California	2-4	1
Salmonberry	2	1

WEED SPECIES	APPLICATION RATE (Quarts/Acre)	HAND-HELD % SOLUTION
Saltcedar*	2-5	1-2
Sassafras*	2-5	1-2
Sourwood*	2-5	1-2
Sumac; laurel, poison, smooth, sugarbush, winged*	2-4	1-2
Sweetgum	2-3	1-1.5
Swordfern*	2-5	1-2
Tallowtree, Chinese	—	1
Tan oak resprouts*	—	2
Thimbleberry	2	1
Tobacco, tree*	2-4	1-2
Trumpet creeper	2-3	1-1.5
Vine maple*	2-5	1-2
Virginia creeper	2-5	1-2
Waxmyrtle, southern*	2-5	1-2
Willow	3	1
Yerbasanta*	—	2

*Partial Control

FARMSTEAD WEED CONTROL

When applied as directed for **NONCROP USES** under the conditions described, Quali-Pro Glyphosate T&O controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts, and for general nonselective farmstead weed control.

Review the **WEEDS CONTROLLED** sections of this label for instructions and specific rates of application to control various perennial and annual weeds.

FARM DITCHES

When used in accordance with instructions, Quali-Pro Glyphosate T&O will suppress perennial grasses growing along farm ditches. A rate of 6 to 8 fluid ounces per acre should be used. For treating fine fescue, tall (coarse) fescue, orchardgrass, or quackgrass covers, 8 fluid ounces per acre should be used.

For best results, ammonium sulfate may be added at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces of Quali-Pro Glyphosate T&O per acre without ammonium sulfate when treating Kentucky bluegrass.

Use 10 to 20 gallons of spray solution per acre for actively growing perennial grass covers. Flat-fan nozzles should be used for optimum spray coverage and distribution.

Add a nonionic surfactant at a rate of 0.5 percent of the spray solution.

For control or suppression of broadleaf weeds, Quali-Pro Glyphosate T&O should be tank mixed with a labeled broadleaf weed herbicide. When using tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

CONSERVATION RESERVE PROGRAM (CRP ACRES)

Use Quali-Pro Glyphosate T&O to control undesirable vegetation when rotating out of CRP acres, or for suppressing competitive growth and production of seeds of undesirable vegetation in CRP acreage.

Review the **WEEDS CONTROLLED** section of this label for application rates for various perennial and annual weeds.

Use conventional spray equipment or wiper applicators for CRP uses.

When using broadcast spray equipment for selective applications, use 12 to 16 ounces of Quali-Pro Glyphosate T&O per acre in early spring before desirable CRP grasses (such as crested and tall wheatgrass) break dormancy and initiate green growth. Wait until desirable perennial grasses reach dormancy before applying in late fall.

NOTE: Some stunting of CRP perennial grasses will occur if treatments are made when plants have not reached dormancy.

TURFGRASSES AND GRASSES FOR SEED PRODUCTION

Preplant and Renovation

When applied as directed for **NONCROP USES** under conditions described, Quali-Pro Glyphosate T&O controls most existing vegetation prior to the planting or renovation of grass seed production areas.

Review the **WEEDS CONTROLLED** sections of this label for specific instructions and rates of application for the control of various trees, woody brush, and perennial and annual weeds.

For best results in controlling existing vegetation, users should delay planting to see if any regrowth occurs from escaped underground plant parts. Where additional treatments are needed, there must be sufficient regrowth prior to application. Summer or fall applications provide best control of warm-season grasses, such as bermudagrass.

DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE APPLYING TREATMENT. Unless otherwise specified, tillage, vertical mowing, slicing, coring, or other renovation techniques must be delayed for at least 7 days after application to facilitate proper translocation into underground plant parts.

Turfgrasses

Where existing vegetation is growing in a field or other unmowed area, apply Quali-Pro Glyphosate T&O to actively growing weeds at the stages of growth that are listed in this label's **WEEDS CONTROLLED** sections.

Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting a minimum of one regular mowing. This will allow sufficient growth for good interception of the spray.

Desirable turfgrasses may be planted after following the procedures described above.

Grasses for Seed Production

Apply Quali-Pro Glyphosate T&O to actively growing weeds at the stages of growth recommended in the **WEEDS CONTROLLED** section of this label before planting or renovating turf or forage grass areas that are grown for seed production.

Do not permit grazing or feeding of treated areas for a minimum of 8 weeks after treatment.

ANNUAL WEED CONTROL IN DORMANT BAHIAGRASS AND BERMUDAGRASS

When applied as directed for **NONCROP USES** under the conditions described herein, Quali-Pro Glyphosate T&O controls or suppresses many winter annual weeds and tall fescue for effective release of dormant bahiagrass and bermudagrass turf. For recommended rates and volumes on the species to be suppressed or controlled, review the rate table for Quali-Pro Glyphosate T&O Alone under the **RELEASE OF BERMUDAGRASS AND BAHIAGRASS** section of this label. Apply only when turf is dormant and before spring greenup. Broadcast applications or spot treatments of this product of more than 16 fluid ounces per acre may result in delayed greenup or injury in highly maintained turfgrass areas such as lawns, golf courses, etc. **DO NOT APPLY TANK MIXTURES** of Quali-Pro Glyphosate T&O plus Oust in turfgrass areas that are highly maintained.

RELEASE OF BAHIAGRASS OR BERMUDAGRASS

NOTE: Apply only in areas where bahiagrass or bermudagrass are desirable ground covers and where temporary discoloration or injury can be tolerated. Use tank mixtures of Quali-Pro Glyphosate T&O plus Oust (or a registered generic alternative) only on highways, railroads, utility plant sites, or other right-of-way areas.

When this product is applied as directed for **NONCROP USES** under the conditions described, it provides control or suppression of tall fescue and many winter annual weeds for effective release of dormant bahiagrass or bermudagrass. Quali-Pro Glyphosate T&O may be tank mixed with Oust as recommended to provide residual control. Apply to dormant bahiagrass or bermudagrass. Tank mixtures of Quali-Pro Glyphosate T&O plus Oust could delay greenup. To minimize injury and to avoid delays in greenup, do not exceed 1 ounce of Oust per acre on bermudagrass, or do not exceed 0.5 ounce per acre on bahiagrass, or apply when these grasses are in a semi-dormant condition.

To obtain best results on winter annuals, apply when plants are in an early growth stage (less than 6 inches tall) after most have germinated. To obtain best results on tall fescue, apply when fescue is either in or beyond the 4- to 6-leaf stage.

Weeds Controlled

Listed below are the rate recommendations for control or suppression of winter annuals and tall fescue.

Users may apply the recommended rates of Quali-Pro Glyphosate T&O alone or as a tank mixture in 10 to 25 gallons of water, plus 0.5- to 1-percent nonionic surfactant by total spray volume per acre.

Contact your Makhteshim Agan of North America, Inc. sales representative for the best recommendation regarding the mixture of weeds within your geographic area.

Weeds Controlled or Suppressed with Quali-Pro Glyphosate T&O Alone*

NOTE: C = Control S = Suppression

WEED SPECIES	GLYPHOSATE T&O FLUID OUNCES/ACRE					
	8	12	16	24	32	64
Barley, little (<i>Hordeum Pusillum</i>)	S	C	C	C	C	C
Bedstraw, catchweed (<i>Galium aparine</i>)	S	C	C	C	C	C
Bluegrass, annual (<i>Poa annua</i>)	S	C	C	C	C	C
Chervil (<i>Chaerophyllum tainturieri</i>)	S	C	C	C	C	C
Chickweed, common (<i>Stellaria media</i>)	S	C	C	C	C	C
Clover, crimson (<i>Trifolium incarnatum</i>)	–	S	S	C	C	C
Clover, large hop (<i>Trifolium incarnatum</i>)	–	S	S	C	C	C
Fescue, tall (<i>Festuca arundinaceae</i>)	–	–	–	–	S	S
Geranium, Carolina (<i>Geranium carolinianum</i>)	–	–	S	S	C	C
Henbit (<i>Lamium amplexicaule</i>)	–	S	C	C	C	C
Ryegrass, Italian (<i>Lolium multiflorum</i>)	–	–	S	C	C	C
Speedwell, corn (<i>Veronica arvensis</i>)	S	C	C	C	C	C
Vetch, common (<i>Vicia sativa</i>)	–	–	S	C	C	C

* These rates only apply to those sites where an established competitive turf is present.

Weeds Controlled or Suppressed with Quali-Pro Glyphosate T&O Plus OUST*

NOTE: C = Control S = Suppression

Quali-Pro Glyphosate T&O Fluid Ounces/Acre	Quali-Pro Glyphosate T&O PLUS OUST						
	8	12	12	16	16	12	16
	+	+	+	+	+	+	+
OUST Ounces/Acre	1/4	1/4	1/2	1/4	1/2	1	1
Barley, little (<i>Hordeum Pusillum</i>)	C	C	C	C	C	C	C
Bedstraw, catchweed (<i>Galium aparine</i>)	C	C	C	C	C	C	C
Bluegrass, annual (<i>Poa annua</i>)	S	C	C	C	C	C	C
Chervil (<i>Chaerophyllum tainturieri</i>)	C	C	C	C	C	C	C
Chickweed, common (<i>Stellaria media</i>)	S	C	C	C	C	C	C
Clover, crimson (<i>Trifolium incarnatum</i>)	S	S	S	S	C	C	C
Clover, large hop (<i>Trifolium incarnatum</i>)	–	–	S	S	S	C	C
Fescue, tall (<i>Festuca arundinaceae</i>)	–	–	–	–	–	S	S
Geranium, Carolina (<i>Geranium carolinianum</i>)	–	S	S	C	C	C	C
Henbit (<i>Lamium amplexicaule</i>)	–	S	C	C	C	C	C
Ryegrass, Italian (<i>Lolium multiflorum</i>)	–	S	S	C	C	C	C
Speedwell, corn (<i>Veronica arvensis</i>)	S	C	C	C	C	C	C
Vetch, common (<i>Vicia sativa</i>)	C	C	C	C	C	C	C

* These rates (or mixtures of rates) only apply to those sites where an established competitive turf is present.

Release of Actively Growing Bermudagrass

When applied as directed, Quali-Pro Glyphosate T&O will aid in the release of bermudagrass by providing control of annual species listed in this and the Oust labels' **WEEDS CONTROLLED** sections, and partial control or suppression of certain perennial weeds.

Use 1 to 3 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre for control or suppression of those annual species listed on this label. When treating annual weeds below 6 inches in height (or length of runner in annual vines), use the lower rate. As weeds increase in size or as they approach flower or seedhead formation, use the higher rate.

For partial control of the perennial species listed below, use the higher rate of this product. For suppression of growth, use the lower rates. To obtain best results, review this label's **WEEDS CONTROLLED** section for proper stage of growth.

Bahiagrass

Paspalum notatum

Bluestem, silver

Andropogon saccharoides

Fescue, tall

Festuca arundinacea

Johnsongrass*

Sorghum halepense

Trumpet creeper**

Campsis radicans

Vaseygrass

Paspalum urvillei

* Provides control at the higher rates.

** Provides suppression only at higher rates.

Quali-Pro Glyphosate T&O may be tank mixed with Oust. If tank mixed, do not exceed 1 to 2 pints per acre of Quali-Pro Glyphosate T&O with 1 to 2 ounces of Oust per acre.

Apply the lower rates of both mixtures to control annual weeds less than 6 inches in height (or runner length in annual vines) that are listed in this and the Oust labels' **WEEDS CONTROLLED** sections. Apply the higher rates as annual weeds increase in size and as they approach the seedhead or flower stages.

To provide partial control of the perennial weeds listed below, use the higher rates of Quali-Pro Glyphosate T&O. For suppression of growth, use the lower rates.

Bahiagrass

Paspalum notatum

Bluestem, silver

Andropogon saccharoides

Broomsedge

Andropogon virginicus

Dock, curly

Rumex crispus

Dogfennel

Eupatorium capilliflorum

Fescue, tall

Festuca arundinacea

Johnsongrass*

Sorghum halepense

Poorjoe*

Diodia teres

Trumpet creeper**

Campsis radicans

Vaseygrass

Paspalum urvillei

Vervain, blue

Verbena hastata

* Provides suppression only at higher rates.

** Provides control at the higher rates.

Quali-Pro Glyphosate T&O should be applied only on well-established bermudagrass. Injury to bermudagrass may result from treatment, but regrowth will occur under moist conditions. Severe injury may result from repeat applications in the same season and are therefore not recommended.

Review and carefully observe all cautionary statements and other information that appears on the labels of all herbicides used.

COOL-SEASON TURF GROWTH REGULATION

When applied as directed herein, Quali-Pro Glyphosate T&O suppresses growth and seedhead development of the listed turf species in industrial sites.

Quali-Pro Glyphosate T&O may be used for management of coarse turf on roadside rights-of-way or other industrial areas. Do not apply on high-quality turf or other areas where some turf color changes are not acceptable. Slight turf discoloration may result, but turf will regreen and regrow under moist conditions as the effects of Quali-Pro Glyphosate T&O dissipate.

Mix and apply 4 to 6 fluid ounces of Quali-Pro Glyphosate T&O per acre alone or in a recommended tank mixture. Use spray volumes of 10 to 40 gallons per acre.

When using Quali-Pro Glyphosate T&O, mix and apply 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

Quali-Pro Glyphosate T&O can be used for growth and seedhead suppression as stated below.

Tall Fescue and Smooth Brome

To obtain best results, apply Quali-Pro Glyphosate T&O in a recommended tank mixture to actively growing turfgrasses after greenup in spring. For suppression of seedheads, treatments must be applied before the boot-to-seedhead stage of development. Applications made from the time of seedhead emergence until maturity may result in turf injury or discoloration.

After mowing or the removal of seedheads, Quali-Pro Glyphosate T&O – in a recommended tank mixture – may also be used to suppress the growth of certain turfgrasses. Turf must be permitted to recover from stress caused by drought, heat, or mowing before making applications. If applications are made to turf under stress, turf discoloration or injury may occur.

Annual Grasses

To suppress growth of some annual grasses such as wild barley, annual ryegrass, and wild oats, apply 3 to 4 fluid ounces of Quali-Pro Glyphosate T&O in 10 to 40 gallons of spray solution per acre. Apply when annual grasses are actively growing and before the seedheads are in the boot stage of development. If treatments are made after seedhead emergence, injury to the desired grasses may occur.

Tank Mixtures: For the tank mixtures mentioned below, review each product label for weeds controlled and their correct stage of application. Do not treat turf that is under stress.

Tank mixtures plus 2,4-D Amine: To obtain additional weed control benefits, up to 1 pound active ingredient per acre of 2,4-D amine may be added to the tank mixtures listed below. Consult the label for 2,4-D amine for weeds controlled.

Tall Fescue

Quali-Pro Glyphosate T&O plus Telar™: For the suppression of tall fescue growth and seedheads, and to control or partially control some annual weeds, apply this tank mixture after greenup and before the boot-to-seedhead stage of development. Use no more than 0.5 ounces of Telar per acre.

For turf-growth suppression, this tank mixture can also be applied after mowing or removal of tall fescue seedheads. Make only one of the above-mentioned applications per growing season.

Quali-Pro Glyphosate T&O plus Oust: For the suppression of tall fescue growth and seedheads, and to control or partially control some annual weeds, apply this tank mixture after greenup and before the boot-to-seedhead stage of development. Use no more than 0.25 ounces of Oust per acre.

Quali-Pro Glyphosate T&O plus Escort™: For turf-growth suppression and control or partial control of some annual weeds, this tank mixture can be applied after mowing or after the removal of tall fescue seedheads. Use no more than 1/3 ounces of Escort per acre.

NOTE: Quali-Pro Glyphosate T&O is not registered for use with Escort in the state of California. Registered generic alternatives may be substituted for the brand name products listed herein.

Smooth Brome

Quali-Pro Glyphosate T&O plus Oust: For control or partial control of some annual weeds, and for suppression of smooth brome growth and seedheads, apply this tank mixture after greenup and before the boot-to-seedhead stage of development. Use no more than 0.25 ounce of Oust per acre.

BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When Quali-Pro Glyphosate T&O is applied as directed in the specified noncrop areas (roadsides, airports, golf course roughs, and plant sites), it will significantly inhibit seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications, and for approximately 120 days with sequential applications.

Apply Quali-Pro Glyphosate T&O from 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of approximately 3 to 4 inches. Applications must be made before seedheads emerge. Apply 6 fluid ounces of Quali-Pro Glyphosate T&O per acre plus 0.5- to 1-percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

To extend the period of seedhead and vegetative growth suppression, sequential applications of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume may be made at approximately 45-day intervals. For continued seedhead suppression, sequential applications must be made before seedheads emerge. Do not exceed 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of Quali-Pro Glyphosate T&O per acre plus nonionic surfactant. Approximately 45 days after the last application, a second sequential application of 2 to 4 fluid ounces per acre plus nonionic surfactant may be made.

For seedhead inhibition and vegetative suppression, a tank mixture of this product plus Oust **may be applied only on roadsides**. Apply 6 fluid ounces of Quali-Pro Glyphosate T&O per acre plus 0.25 ounces per acre of Oust, plus 0.5- to 1-percent nonionic surfactant by total spray volume 1 to 2 weeks after an initial mowing in spring. When using Quali-Pro Glyphosate T&O plus Oust for suppression of bahiagrass, apply only once per year.

CROPPING SYSTEMS

When used in accordance with the instructions and under the conditions described in this label's **CROPPING SYSTEMS** section, Quali-Pro Glyphosate T&O controls those perennial and annual weeds listed on this label. Use prior to the emergence of direct-seeded crops or prior to transplanting the crops listed on this label.

For product performance information, please review the sections of this label titled **GENERAL INFORMATION** and **MIXING, ADDITIVES** and **APPLICATION INSTRUCTIONS**.

EXERCISE EXTREME CAUTION TO AVOID SPRAY CONTACT WITH FOLIAGE, GREEN STEMS, CROP FRUITS, OR OTHER DESIRABLE PLANTS. DESTRUCTION OR SEVERE DAMAGE MAY RESULT.

To control weeds regenerating from seed or underground parts, multiple treatments may be needed. Unless otherwise stated on this label, all treatments must be made before the crop emerges in accordance with this label's instructions.

Unless otherwise stated in a specific crop section of this label, the

maximum, combined total of all treatments cannot exceed 8 quarts per acre of Quali-Pro Glyphosate T&O per year.

NOTE: THE MAXIMUM USE RATES DESCRIBED IN THIS LABEL APPLY TO QUALI-PRO GLYPHOSATE T&O WHEN COMBINED WITH ALL OTHER HERBICIDES CONTAINING THE ACTIVE INGREDIENTS GLYPHOSATE OR SULFOSATE—WHETHER APPLIED AS A MIXTURE OR APPLIED SEPARATELY. USERS MUST CAREFULLY CALCULATE ALL APPLICATION RATES AND MAKE SURE THAT TOTAL USAGE OF QUALI-PRO GLYPHOSATE T&O AND OTHER PRODUCTS CONTAINING GLYPHOSATE OR SULFOSATE DOES NOT EXCEED THE RECOMMENDED MAXIMUM USAGE RATES.

For any crop not listed below, applications must be made at least 30 days prior to planting.

Do not harvest or feed treated vegetation for a minimum of 8 weeks after application. If applying spot treatments or using selective equipment, wait a minimum of 14 days before grazing domestic livestock or harvesting legumes or grasses.

See the following **CROPPING SYSTEMS** sections for specific recommended uses.

ROW CROPS

CORN (ALL)*
COTTON*
PEANUTS
SORGHUM (MILO)*
SOYBEANS*
SUGARCANE*

CEREAL GRAINS

BARLEY*
BUCKWHEAT*
MILLET (PEARL, PROSO)*
OATS*
RICE**
RYE*
TRITICALE*
WHEAT (ALL)*
WILD RICE*

CITRUS

CALAMONDIN
CHIRONJA
CITRON
GRAPEFRUIT
KUMQUAT
LEMON
LIME
MANDARIN ORANGE
ORANGE (ALL)
PUMMELO
TANGELO
TANGERINE
TANGORS

TREE NUTS

ALMOND
BEECHNUT
BRAZIL NUT
BUTTERNUT
CASHEW
CHESTNUT
CHINQUAPIN
FILBERT (HAZELNUT)
HICKORY NUT
MACADAMIA
PECAN
PISTACHIO
WALNUT (BLACK, ENGLISH)

VINE CROPS

GRAPES
KIWI FRUIT

TREE FRUITS

APPLE
APRICOTS
CHERRY (SWEET, SOUR)
LOQUAT
MAYHAW
NECTARINE
OLIVE
PEACH
PEAR
PLUM/PRUNE (ALL)
QUINCE

VEGETABLES

ARTICHOKE,
JERUSALEM
ASPARAGUS*
BEANS (ALL)
BEET GREENS
BEETS (RED,SUGAR)
BROCCOLI (ALL)
BRUSSELS SPROUTS
CABBAGE (ALL)
CABBAGE, CHINESE
CANTALOUPE***
CARROT
CAULIFLOWER
CASABA MELON***
CELERIAC
CELERY
CHARD, SWISS
CHICORY
COLLARDS
CRENSHAW MELON***
CUCUMBER***
EGGPLANT***
ENDIVE
GARLIC***

GOURDS***
 GROUND CHERRY***
 HONEYDEW MELON***
 HONEY BALL MELON***
 HORSERADISH
 KALE
 KOHLRABI
 LEEK
 LENTILS
 LETTUCE
 MANGO MELON***
 MELONS (ALL)***
 MUSKMELON***
 MUSTARD GREENS
 OKRA
 ONION
 PARSLEY
 PARSNIPS
 PEAS (ALL)
 PEPPER (ALL)***
 PERSIAN MELON***
 POTATO (IRISH,SWEET)
 PUMPKIN***
 RADISH
 RAPE GREENS
 RHUBARB
 RUTABAGA
 SHALLOT
 SPINACH (ALL)
 SQUASH (SUMMER,WINTER)***
 TOMATILLO***
 TOMATO***†
 TURNIP
 WATERCRESS***
 WATERMELON***
 YAMS

SMALL FRUITS AND BERRIES

BLACKBERRY
 BLUEBERRY
 BOYSENBERRY
 CRANBERRY
 CURRANT
 DEWBERRY
 ELDERBERRY
 GOOSEBERRY
 HUCKLEBERRY
 LOGANBERRY
 OLALLIEBERRY
 RASPBERRY (BLACK,RED)

FORAGE CROPS AND LEGUMES

ALFALFA*
 FORAGE GRASSES*
 FORAGE LEGUMES*

TROPICAL CROPS

ACEROLA
 ATEMOYA
 AVOCADO
 BANANA
 BREADFRUIT
 CANISTEL
 CARAMBOLA
 CHERIMOYA
 COCOA BEANS
 COFFEE
 DATES
 FIGS
 GENIP
 GUAVA
 JABOTICABA
 JACKFRUIT
 LONGAN
 LYCHEE
 MANGO
 PAPAYA
 PASSION FRUIT
 PERSIMMONS
 PLANTAINS
 PINEAPPLE****
 POMEGRANATE
 SAPODILLA
 SAPOTE (BLACK, MAMEY, WHITE)
 SOURSOP
 SUGAR APPLE
 TAMARIND
 TEA

* Spot treatments allowed.

** Do not treat rice fields or levees when the fields contain flood water.

*** Apply before planting only. Wait a minimum of 3 days between application and planting.

**** Do not feed or graze treated pineapple forage following application.

† Use on direct-seeded crops only.

NOTE: When applying Quali-Pro Glyphosate T&O before transplanting crops into plastic mulch, take care to remove residues of this product—which can cause crop injury—from the plastic before transplanting. A single 1/2 inch application of water, by either natural rainfall or a sprinkler irrigation system, will remove residues. If treatments are applied at emergence, injury or death to the emerged seedlings will result.

Spot Treatment (only for those crops marked with an *): Treatments made to growing crops must be made before the following: Heading of small grains and milo; silking of corn; initial pod set in soybeans; or boll opening on cotton.

For information regarding forage grasses and forage legumes, please review **SPOT TREATMENT** in the **PASTURES** section of

CROPPING SYSTEMS of this label.

For information on dilution and application rates using boom or hand-held equipment, please review the **MIXING, ADDITIVES and APPLICATION INSTRUCTIONS** along with the **WEEDS CONTROLLED** sections of this label.

NOTE: FOR TREATING FORAGE GRASSES AND FORAGE LEGUMES, DO NOT APPLY TREATMENT TO MORE THAN ONE-TENTH OF ANY ACRE AT ONE TIME. FOR ALL OTHER CROPS, NO MORE THAN 10 PERCENT OF THE TOTAL FIELD AREA TO BE HARVESTED SHOULD BE TREATED.

CROPS THAT ARE SPRAYED IN TREATED AREAS WILL BE KILLED. EXERCISE CAUTION TO AVOID DRIFT OR SPRAY OUTSIDE THE TARGET AREA—PLANT INJURY OR DESTRUCTION WILL RESULT.

Selective Equipment: For cotton or soybeans, Quali-Pro Glyphosate T&O may be applied through shielded applicators, wiper applicators, or recirculating sprayers. Shielded and wiper applicators may also be used in treating grapes and tree crops. Wiper applicators may be used in rutabagas, forage grasses and forage legumes, wheat, pasture sites, and grain sorghum (milo).

For specific information on proper use and calibration of equipment, please review the **SELECTIVE EQUIPMENT** section of the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label.

For the following crops, allow—at a minimum—the noted time interval between application and harvest:

CROPS	MINIMUM INTERVAL BETWEEN TREATMENT AND HARVEST
Apples, Citrus, Pear	1 day
Nut Crops	3 days
Cotton, Soybeans	7 days
Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherry, Grapes, Dates, Jaboticaba, Jackfruit, Longan, Lychee, Passion Fruit, Persimmons, Rutabagas, Sapodilla, Sapote, Soursop, Sugar Apple, Tamarind	14 days
Stone Fruit	17 days
Wheat ¹	35 days
Sorghum (milo) ^{1, 2}	40 days

¹ Do not use roller applications.

² Do not graze or feed treated milo fodder. Do not ensile treated vegetation.

ASPARAGUS

When used in asparagus in accordance with the instructions and under the conditions described under **CROPPING SYSTEMS**, Quali-Pro Glyphosate T&O controls those weeds listed on this label.

Review the **WEEDS CONTROLLED** section of this label for specific

instructions and application rates for controlling various perennial and annual weeds.

Prior to Crop Emergence: For controlling emerged annual and perennial weeds mentioned on this label, apply Quali-Pro Glyphosate T&O before crop emergence. DO NOT APPLY THIS PRODUCT WITHIN ONE WEEK BEFORE THE FIRST ASPARAGUS SPEARS EMERGE.

Spot Treatment: Quali-Pro Glyphosate T&O should be applied immediately after cutting, but before the emergence of new asparagus spears. No more than 10 percent of the total field area to be harvested should be treated. Wait at least 5 days after treatment before harvesting.

Postharvest: Quali-Pro Glyphosate T&O should be applied after the final harvest and all asparagus spears have been removed. If spears are permitted to regrow, wait until ferns have developed before applying Quali-Pro Glyphosate T&O. If treatments are delayed as such, they should be applied as directed sprays or as shielded sprays to avoid herbicide contact with ferns, stems, or spears. IF QUALI-PRO GLYPHOSATE T&O COMES INTO DIRECT CONTACT WITH ASPARAGUS, SERIOUS CROP INJURY OR DESTRUCTION MAY RESULT.

NOTE: Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

BERRIES AND SMALL FRUITS

For cranberries, wiper applicators may be used in accordance with the instructions in this section.

For berries other than cranberries, use Quali-Pro Glyphosate T&O as a preplant broadcast application, or as a directed spray or wiper application post-planting.

For important product performance information, please review the **GENERAL INFORMATION and MIXING, ADDITIVES and APPLICATION INSTRUCTIONS** sections of this label.

For specific information on recommended use and calibration of equipment, please review the **SELECTIVE EQUIPMENT** section of the label section titled **APPLICATION EQUIPMENT AND TECHNIQUES**.

Do not harvest cranberries less than 30 days after the last application of Quali-Pro Glyphosate T&O. For other small fruits, allow at least 14 days after the last application of Quali-Pro Glyphosate T&O before harvest.

For Wick or Other Wiper Applicators: Prepare a 20-percent solution by mixing 1 gallon of Quali-Pro Glyphosate T&O in 4 gallons of water. For severe infestations, reduce ground speed of the application equipment so that adequate amounts of Quali-Pro Glyphosate

T&O are wiped on weeds. A second application going in the opposite direction may be beneficial.

DO NOT ALLOW QUALI-PRO GLYPHOSATE T&O SOLUTIONS TO COME INTO CONTACT WITH DESIRABLE VEGETATION, INCLUDING CANES, FOLIAGE, OR GREEN SHOOTS.

CORN

Hooded Sprayers: For controlling weeds between rows of corn, Quali-Pro Glyphosate T&O may be used through hooded sprayers – using only those hooded sprayers that completely enclose the spray pattern.

A hooded sprayer is one type of shielded applicator. When a hooded sprayer is used, the crop is shielded from the spray solution because the spray pattern is completely enclosed by a hood on the top and on all 4 sides. It is extremely important that hooded sprayers are set up and operated in a way that avoids bouncing or raising the hoods off the ground in any way. If hoods are raised, spray may escape and contact the crop, causing crop damage or destruction. Spray hoods must be operated on the ground or while skimming across the ground, and users must adjust tractor speed to avoid spray hood bouncing. Do not operate hooded sprayers on rough or sloping ground where the spray hoods might be raised off the ground.

If applying Quali-Pro Glyphosate T&O to corn that is growing on raised beds, the hood must be designed to completely enclose the spray solution. If necessary, the front and rear flaps of the hoods should be extended so that they reach the ground in deep furrows.

To minimize damage to desirable crops and to obtain best results, observe the following:

1. Spray hoods must be operated on the ground or skimming across the ground.
2. Apply no more than 1 quart of Quali-Pro Glyphosate T&O per acre per application.
3. Corn must be a minimum of 12 inches tall (measured without extending leaves).
4. Leave at least an 8-inch untreated strip over the drill row. (Example: if the crop row is 38 inches wide, the maximum width of the spray hood should be 30 inches.)
5. Do not exceed a tractor speed of 5 mph.
6. Do not apply if wind speed exceeds 10 mph.
7. Use low-drift nozzles.

Crop injury or destruction may result if the foliage of treated weeds comes into contact with the leaves of the desirable crop. Therefore, Quali-Pro Glyphosate T&O should not be applied if the leaves of the crop are growing in direct contact with the weeds to be killed. Droplets, foam, splatter, or mist from the Quali-Pro Glyphosate T&O solution could contact the crop and cause stunting, discoloration, or crop destruction.

SEVERE DAMAGE OR DESTRUCTION MAY RESULT IF THIS PRODUCT COMES INTO CONTACT WITH ANY VEGETATION ON WHICH APPLICATION WAS NOT INTENDED. SUCH DAMAGE SHALL BE THE SOLE RESPONSIBILITY OF THE APPLICATOR.

Please review the **WEEDS CONTROLLED** section of this label for specific instructions and application rates for controlling various perennial and annual weeds.

If Quali-Pro Glyphosate T&O treatments are applied to corn using hooded sprayers, do not graze or feed corn forage or fodder to livestock.

Do not apply more than 3 quarts of Quali-Pro Glyphosate T&O per acre per year for hooded sprayer applications.

FALLOW AND REDUCED-TILLAGE SYSTEMS

FOR AERIAL APPLICATION IN CALIFORNIA AND ARKANSAS, REFER TO THE SECTION TITLED SUPPLEMENTAL USES AT THE END OF THIS LABEL.

For controlling annual weeds prior to emergence of labeled crops, use Quali-Pro Glyphosate T&O in fallow and reduced tillage systems. For specific application rates and instructions, please review the **WEEDS CONTROLLED** sections of this label. Treatments of Quali-Pro Glyphosate T&O may be applied using aerial or ground spray equipment. Review the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for further information.

●	TANK MIXTURES	●
●	QUALI-PRO GLYPHOSATE T&O plus BANVEL plus NONIONIC SURFACTANT	●
●	QUALI-PRO GLYPHOSATE T&O plus 2,4-D plus NONIONIC SURFACTANT	●
●	QUALI-PRO GLYPHOSATE T&O plus GOAL™ plus NONIONIC SURFACTANT	●

IN CALIFORNIA, DO NOT APPLY 2,4-D OR BANVEL TANK MIXTURES BY AIR.

For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Treatments of Banvel or 2,4-D must be applied a minimum of 7 days before planting corn.

A mixture of Banvel with Quali-Pro Glyphosate T&O may provide short-term residual control of some weed species. However, some crop injury may result if Banvel is applied within 45 days of planting. Please review the 2,4-D and Banvel labels for use instructions and cropping restrictions.

Quali-Pro Glyphosate T&O Herbicide plus Goal (or Generic Equivalent) Tank Mixtures

For controlling the weeds listed below, use Quali-Pro Glyphosate T&O alone or in a tank mixture with Goal plus 0.5- to 1-percent nonionic surfactant by total spray volume.

Treatments should be applied when weeds are actively growing and at the height/recommended stage of growth specified below. Avoid application when weeds are under moisture stress, when dust is on foliage, or when straw canopy is covering the weeds, as unsatisfactory control may result.

QUALI-PRO GLYPHOSATE T&O @ 12 FL. OZ. PER ACRE

Species	Maximum length / height in inches
Wheat	18
Barley	12
Bluegrass, annual	6
Barnyardgrass	6
Rye	6

QUALI-PRO GLYPHOSATE T&O @ 16 FL. OZ. PER ACRE

Species Annual Weeds Above, Plus:	Maximum length / height in inches
Ryegrass, annual	6
Chickweed	6
Groundsel	6
Marestail*	6
Rocket, London	6
Shepherd's purse	6
Crabgrass	12
Johnsongrass, seedling	12
Lambsquarters	12
Oats, wild	12
Pigweed, redroot	12
Mustards	12

NOTE: 32 fluid ounces of Quali-Pro Glyphosate T&O per acre should be used where heavy weed densities exist.

* Resistance to some biotypes has been observed. If users encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

QUALI-PRO GLYPHOSATE T&O @ 12 FL. OZ. / Acre

plus
GOAL** @ 2 – 4 FL. OZ. / Acre

Species Annual Weeds Above, Plus:	Maximum length / height in inches
Cheeseweed, common	3
Chickweed	3
Groundsel	3
Rocket, London	6
Shepherd's purse	6

QUALI-PRO GLYPHOSATE T&O @ 16 FL. OZ. / Acre

plus
GOAL** @ 2 – 4 FL. OZ. / Acre

Species
Annual Weeds Above, Plus: Maximum length / height in inches

Cheeseweed, common	6
Chickweed	12
Groundsel	6
Rocket, London	12
Shepherd's purse	12

** The higher rate of Goal should be used when weeds approach maximum height or stands are dense.

NOTE: Mix 32 fluid ounces of Quali-Pro Glyphosate T&O per acre with 2-4 ounces of Goal per acre on heavy weed densities.

Ground or aerial spray equipment may be used to apply recommended tank mixtures. Please review this label's **WEEDS CONTROLLED** sections for instructions and specific application rates.

ECOFARMING SYSTEMS

The uses listed in this section are not registered for use in California.

The "Ecofarming System" consists of a rotation of winter wheat, corn/sorghum, and ecofallow.

The following tank mixtures may be used for controlling emerged annual weeds prior to planting corn or sorghum in the Ecofarming System. Review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Quali-Pro Glyphosate T&O @ 16 to 20 FL. OZ. per acre

plus
2,4-D at 0.375 to 0.5 pound a.i. per acre
plus

Atrazine at 0.75 to 1 pound a.i. per acre
plus

Lasso® herbicide at 2.5 to 3 quarts per acre

The tank mixture listed above should be applied in a liquid fertilizer carrier of 28-0-0 or 32-0-0 analysis at 20 to 30 gallons per acre. Water may be used to dilute the liquid fertilizer to attain the desired carrier volume.

WEEDS CONTROLLED: This tank mixture will control the weeds listed below up to a maximum of 4 inches in height.

Brome, downy
Bromus tectorum

Cheat
Bromus secalinus

Foxtail, green
Setaria viridis

Foxtail, yellow
Setaria lutescens

Kochia*
Kochia scoparia

Lettuce, prickly
Lactuca serriola

Pigweed, redroot
Amaranthus retroflexus

Thistle, Russian
Salsola kali

Wheat, volunteer
Triticum aestivum

* For improved results when applying to kochia, add Banvel at a rate of 4 fluid ounces per acre (0.125 pound of active ingredient per acre) to the above tank mixture.

Users may reduce the risk of crop injury from 2,4-D or Banvel by applying this mixture 7 to 14 days prior to planting.

Refer to the label booklet for Lasso herbicide for preemergence weed control with this tank mixture.

Prior to mixing/applying this tank mix, review the specific product labels of all tank mix products for cautionary statements and crop rotation restrictions.

AID TO TILLAGE

Quali-Pro Glyphosate T&O will control downy brome, volunteer wheat, cheat, foxtail, and tansy mustard when used in conjunction with preplant tillage practices. Apply 8 fluid ounces of Quali-Pro Glyphosate T&O plus 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and before they reach 6 inches in height. Treatments must be followed by conventional tillage practices within 15 days after treatment and before regrowth occurs. Allow a minimum of 1 day after treatment before tillage. Reduced effectiveness may result if tank mixtures with residual herbicides are used.

POSTHARVEST GRAIN SORGHUM, SORGHUM REGROWTH CONTROL

Quali-Pro Glyphosate T&O may be applied to grain sorghum (milo) after harvest for suppression or control of regrowth. For control, apply 1 quart of Quali-Pro Glyphosate T&O per acre. For suppression, use 1.5 pints of this product per acre. In both cases, use a 0.5-percent nonionic surfactant in 3 to 10 gallons of spray solution per acre.

PASTURES

Apply prior to planting forage grasses and legumes.

Pasture or Hay Crop Renovation: Apply Quali-Pro Glyphosate T&O as a broadcast spray prior to planting forage grasses or legumes to control the perennial and annual weeds listed in this label. Domestic livestock should be removed prior to application. Wait 8 weeks after treatment before grazing or harvesting.

Spot Treatment: Quali-Pro Glyphosate T&O can be used as a spot treatment when applied as recommended herein. Spot treatments may be applied to control perennial and annual weeds listed in this label when growing in pastures, forage grasses, and forage legumes composed of bermudagrass, bahiagrass, brome, bluegrass, orchardgrass, fescue, ryegrass, wheatgrass, timothy, clover, or alfalfa.

Wiper Application: Quali-Pro Glyphosate T&O, when applied according to instructions, controls or suppresses the weeds listed under **WIPER APPLICATORS AND SPONGE BARS** in the **SELECTIVE EQUIPMENT** section of this label.

For both spot treatment and wiper application, apply only in areas where the movement of domestic livestock can be controlled. Treat no more than one-tenth of any acre at one time. Additional treatments may be made in the same area at intervals of 30 days. Domestic livestock should be removed prior to application. Wait 2 weeks after treatment before grazing or harvesting.

SUGARCANE

When applied as directed for **CROPPING SYSTEMS**, under the conditions described, Quali-Pro Glyphosate T&O controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields prior to the emergence of plant cane. This product will also control undesirable sugarcane.

NOTE: Where multiple treatments are needed, do not apply more than 10.6 quarts of Quali-Pro Glyphosate T&O per acre per year. Do not apply this product to plants in or around canals, ditches, or ponds containing water that will be used for irrigation.

Broadcast Treatment: Apply Quali-Pro Glyphosate T&O to emerged weeds in 10 to 40 gallons of water per acre prior to the emergence of plant cane.

For control of various perennial and annual weeds, please see this label's **WEEDS CONTROLLED** sections for specific instructions and rates of application.

For removing last stubble or ratoon cane, apply 4 to 5 quarts of Quali-Pro Glyphosate T&O in 10 to 40 gallons of water per acre. Apply to new growth having a minimum of 7 or more new leaves. Wait a minimum of 7 days after treatment before tillage.

Spot Treatment in or Around Sugarcane Fields: For use with hand-held equipment, review the dilution and application rate information contained in the **MIXING, ADDITIVES and APPLICATION INSTRUCTIONS** and **WEEDS CONTROLLED** sections of this label.

To control diseased or volunteer sugarcane, apply a 1-percent solution of Quali-Pro Glyphosate T&O in water and spray to wet foliage of plants to be controlled.

NOTE: Diseased or volunteer sugarcane should have at least 7 new leaves prior to application.

Avoid spray contact with healthy cane plants since severe damage or destruction may result.

Do not feed or graze treated sugarcane forage following application.

CONSERVATION TILLAGE, MINIMUM TILLAGE, AND NO-TILL SYSTEMS

CORN AND SOYBEAN Tank Mixtures

The uses listed in this section are not registered for use in California.

When applied according to the instructions and under the conditions described herein, tank mixtures listed in this section will control many emerged weeds and provide preemergence control of many annual weeds where soybeans or corn will be planted directly into a cover crop, established sod, or in previous crop residues.

Review crop rotation and cautionary statement information listed on the labels of all products used in these tank mixtures. Review the **MIXING, ADDITIVES**, and **APPLICATION INSTRUCTIONS** section of this label for mixing instructions.

Apply these tank mixtures in 10 to 20 gallons of water, or in 10 to 60 gallons of nitrogen solution per acre before, during, or after planting. Do not apply these mixtures after crop emergence.

When tank mixing Quali-Pro Glyphosate T&O with residual herbicides, add an agriculturally approved nonionic surfactant at the rate of 0.5 to 1 percent by volume of spray solution. Adding 1 to 2 percent dry ammonium sulfate by weight may improve the effectiveness of this product.

NOTE: Do not exceed 4 quarts of Quali-Pro Glyphosate T&O per acre when using these tank mixtures.

CORN

For residual control, this product may be tank mixed with the following herbicides or combination of herbicides:

Atrazine	Lasso/Alachlor
Bicep Magnum™	Micro-Tech®
Bullet®	Partner®
Cyanazine	Prowl™
Dual Magnum™	Simazine
Lariat®	

For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

To improve burndown, tank mix Quali-Pro Glyphosate T&O with dicamba or 2,4-D. Treatments of 2,4-D or dicamba must be made a minimum of 7 days before planting corn. Please review the **WEEDS CONTROLLED** sections of this label for specific rate information.

SOYBEANS

For residual control, apply Quali-Pro Glyphosate T&O with the following herbicides or combination of herbicides:

Canopy™	Partner
Command™	Preview™
Dual Magnum	Prowl
Gemini™	Pursuit™
Lasso/Alachlor	Pursuit Plus™
Lexone™	Scepter™
Linuron	Sencor™
Lorox™ Plus	Squadron™
Micro-Tech	Turbo™

For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

To improve burndown, tank mix Quali-Pro Glyphosate T&O with 2,4-D or 2,4-DB. (Review the 2,4-D label for intervals between application and planting.)

CORN AND SOYBEANS

Annual Weeds: For difficult-to-control weeds such as barnyardgrass, crabgrass, fall panicum, broadleaf signalgrass, and shattercane that are up to 2 inches tall, and for Pennsylvania smartweed that is up to 6 inches tall, apply Quali-Pro Glyphosate T&O at the rate of 2 pints per acre in these tank mixtures. For other annual weeds on this label, apply 1 to 1.5 pints of Quali-Pro Glyphosate T&O per acre when weeds are smaller than 6 inches tall, and 2 to 3 pints per acre when weeds exceed 6 inches tall. Review this label's **WEEDS CONTROLLED** sections for a complete list of annual weeds controlled.

Perennial Weeds: When using minimum tillage systems at normal application times, perennial weeds may not be at the proper stage of development for control. To determine the proper stage of devel-

opment for perennial weeds, review the **WEEDS CONTROLLED** sections of this label.

Under these conditions, use 2 to 4 quarts of Quali-Pro Glyphosate T&O per acre in the tank mixtures mentioned above to obtain top kill and reduce competition from many emerged perennial broadleaf and grass weeds. See the **WEEDS CONTROLLED** sections of this label for information on emerged perennial weeds controlled.

To obtain the desired stage of growth, it may be necessary to apply Quali-Pro Glyphosate T&O alone in the late summer or fall and then follow with a label-approved, seedling weed-control program at planting.

NOTE: USE OF THESE TANK MIXTURES TO CONTROL JOHNSONGRASS OR BERMUDAGRASS IN MINIMUM TILLAGE SYSTEMS IS NOT RECOMMENDED. To control bermudagrass, review the **CONTROL OF PERENNIAL WEEDS** section of this label and follow its instructions, then use a label-approved, seedling weed-control program in a minimum-tillage or conventional-tillage system. To control johnsongrass, review the **CONTROL OF PERENNIAL WEEDS** section and follow its instructions, then use a label-approved, seedling weed-control program with conventional tillage.

PREHARVEST APPLICATIONS

Quali-Pro Glyphosate T&O will control those perennial and annual weeds listed on this label when applied prior to the harvest of soybeans, cotton, grain sorghum (milo), and wheat. Apply as directed and under the conditions described.

Please review the **WEEDS CONTROLLED** sections of this label for specific rates and application instructions to control various perennial and annual weeds.

Use ground or aerial equipment to apply this product. However, DO NOT EXCEED 1 QUART OF QUALI-PRO GLYPHOSATE T&O PER ACRE WHEN APPLYING BY AIR. For specific instructions on ground and aerial applications, please review the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label.

NOTE: Quali-Pro Glyphosate T&O may not be applied to crops grown for seed—a reduction of vigor or of germination may result.

Quali-Pro Glyphosate T&O is not registered in California on preharvest grain sorghum (milo).

SOYBEANS

Apply Quali-Pro Glyphosate T&O after pods have set and have lost all green color. Allow at least 7 days between the application of Quali-Pro Glyphosate T&O and harvesting. Exercise caution to avoid excessive seed shatter loss from ground application equipment.

Do not permit livestock to graze, and do not harvest treated crop for feed within 25 days after the last preharvest application.

FOR PREHARVEST APPLICATIONS, DO NOT APPLY MORE THAN 6 QUARTS OF QUALI-PRO GLYPHOSATE T&O PER ACRE.

COTTON

Apply Quali-Pro Glyphosate T&O after sufficient bolls have developed. If applications are made prior to this time, maximum yield potential could be affected.

Broadcast Applications: Use either aerial or ground spray equipment to apply Quali-Pro Glyphosate T&O. When using broadcast equipment by ground application, apply Quali-Pro Glyphosate T&O in 10 to 20 gallons of water per acre. When applying by air, apply treatments in 3 to 10 gallons of water per acre.

When treatments are applied prior to cotton harvest, Quali-Pro Glyphosate T&O provides weed control and cotton regrowth inhibition. For cotton regrowth inhibition, apply 1 to 2 quarts of Quali-Pro Glyphosate T&O in 3 to 10 gallons of water per acre. For preharvest applications, do not exceed 2 quarts of this product per acre. THE USE OF ADDITIVES FOR PREHARVEST TREATMENTS TO COTTON IS PROHIBITED.

Users may tank mix Quali-Pro Glyphosate T&O with Folex™, DEF™ 6, or Prep™ to improve cotton leaf drop. For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Allow at least 7 days between the last treatment and cotton harvest.

Do not feed treated cotton forage to livestock or permit livestock to graze treated cotton forage or hay after preharvest applications.

GRAIN SORGHUM (MILO)

Apply at a minimum of 7 days prior to harvest and at 30 percent or less grain moisture.

Apply up to 2 quarts of Quali-Pro Glyphosate T&O per acre.

WHEAT

Treatments should be applied at a minimum of 7 days prior to harvest, and after the hard-dough stage of grain (a maximum of 30 percent grain moisture).

FOR PREHARVEST APPLICATIONS TO WHEAT, DO NOT EXCEED 1 QUART QUALI-PRO GLYPHOSATE T&O PER ACRE.

TREE AND VINE CROPS

Use Quali-Pro Glyphosate T&O to control weeds in established vineyards, orchards, and groves, or for site preparation prior to transplanting those crops listed in this section. Unless directed otherwise in this section, treatments can be applied with Controlled Droplet Applicator (CDA), shielded sprayers, boom equipment, hand-held and high-volume wands, lances, orchard guns, or with wiper equipment. For specific information regarding the use of application equipment, please review the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label.

Refer to the **WEEDS CONTROLLED** sections of this label and to specific recommendations in this section for rates to be used.

STRIPS

FOR PERENNIAL AND ANNUAL WEEDS IN STRIPS OF TREE AND VINE CROPS.

Tank Mixtures with Residual Herbicides

The following mixtures control emerged annual weeds and provide control or suppression of emerged perennial weeds listed on this label. The residual herbicides listed below will provide preemergence control of weeds listed on the individual herbicide product labels.

- QUALI-PRO GLYPHOSATE T&O Herbicide plus GOAL 2XL
- QUALI-PRO GLYPHOSATE T&O Herbicide plus KARMEX™ DF
- QUALI-PRO GLYPHOSATE T&O Herbicide plus KROVAR I
- QUALI-PRO GLYPHOSATE T&O Herbicide plus KROVAR II
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SIMAZINE, PRINCEP CALIBER 90
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SIMAZINE 4L
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SIMAZINE 80W
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SOLICAM™ 80DF
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SURFLAN AS or Oryzalin 4 A.S.
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SURFLAN 75W
- QUALI-PRO GLYPHOSATE T&O Herbicide plus SIMAZINE (80W, or 4L, or PRINCEP CALIBER 90) plus Oryzalin 4 A.S. or SURFLAN (AS or 75W)
- QUALI-PRO GLYPHOSATE T&O Herbicide plus GOAL 2XL plus Oryzalin 4 A.S. or SURFLAN (AS or 75W)
- QUALI-PRO GLYPHOSATE T&O Herbicide plus GOAL 2XL plus SIMAZINE (80W, or 4L, or PRINCEP CALIBER 90)
- QUALI-PRO GLYPHOSATE T&O Herbicide plus GOAL 2XL plus Oryzalin 4 A.S. or SURFLAN (AS or 75W) plus SIMAZINE (80W, 4L, or PRINCEP CALIBER 90)

For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements. Read and carefully observe the cautionary statements, claims, rates, and all other information on product labels of all herbicides used.

These tank mixtures cannot be applied in Puerto Rico.

When tank mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution.

Recommended Rates

Annual Weeds: Use 1 to 5 quarts of Quali-Pro Glyphosate T&O per acre in these tank mixtures. Rates at the high end of the recommend range should be used if weeds are stressed, are higher than 12 inches tall, or are growing in dense populations.

Perennial Weeds: For control or suppression of perennial weeds,

apply 1 pint to 5 quarts of Quali-Pro Glyphosate T&O per acre. Review and follow the instructions in the **WEEDS CONTROLLED** sections of this label for application rates and stage of growth information for specific perennial weeds.

QUALI-PRO GLYPHOSATE T&O Herbicide plus GOAL plus SIMAZINE/Oryzalin 4 A.S. (or SURFLAN)

For postemergence control of the weeds listed below, mix Quali-Pro Glyphosate T&O with low rates of Goal in 3-way or 4-way mixtures with simazine and/or Oryzalin 4 A.S. (or Surflan). For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Review the labels for simazine and Oryzalin 4 A.S. (or Surflan) for important information, including preemergence rates and weeds controlled.

Use 3 to 40 gallons of water when applying these tank mixtures. Add a nonionic surfactant to the spray solution at the rate of 0.5 to 1 percent by total spray volume.

For controlling the following weeds, apply 1 to 5 quarts of Quali-Pro Glyphosate T&O per acre, plus 4 to 48 fluid ounces of Goal per acre plus the applicable rates of simazine and/or Surflan (or Oryzalin 4 A.S.) listed on their respective labels. For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Barley, wild
Hordeum leporinum
Bluegrass, annual
Poa annua
Cheeseweed, common
Malva spp.
Chickweed, common
Stellaria media
Filaree*
Erodium spp.
Fleabane, hairy
Conyza bonariensis
Groundsel, common
Senecio vulgaris
Horseweed/Marestail**
Conyza canadensis
Nettle, stinging
Urtica dioica
Pineappleweed
Matricaria matricariodes
Rocket, London
Sisymbrium irio
Shepherd's purse
Capsella bursa-pastoris
Sowthistle, annual
Sonchus oleraceus

* Use at least 1.5 quarts of Quali-Pro Glyphosate T&O in these mixtures.

** Resistance to some biotypes has been observed. If users encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

NOTE: For preemergence weed control, these recommendations do not preclude the use of higher, labeled rates of Goal in these tank mixtures.

PERENNIAL GRASS SUPPRESSION ORCHARD FLOORS

When applied in accordance with instructions, Quali-Pro Glyphosate T&O will suppress vegetative growth as described below.

Bahiagrass

Quali-Pro Glyphosate T&O will significantly inhibit seedhead emergence and will suppress vegetative growth for approximately 45 days with a single application, and with sequential applications, approximately 120 days. Apply Quali-Pro Glyphosate T&O approximately 1-2 weeks after full greenup, or after mowing to a uniform height of 3 to 4 inches. Treatments must be made before seedhead emergence. Apply 6 fluid ounces of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

To extend the period of seedhead and vegetative growth suppression, sequential applications of Quali-Pro Glyphosate T&O plus nonionic surfactant may be made at intervals of approximately 45 days. To continue suppression of seedheads, sequential applications must be made before seedheads emerge. Apply a maximum of 2 sequential applications annually. For the first sequential application, apply 4 fluid ounces of Quali-Pro Glyphosate T&O with nonionic surfactant. For the second sequential application, 2 to 4 ounces may be used at approximately 45 days after the first application.

Bermudagrass

For burndown, apply 1 to 2 quarts of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. East of the Rocky Mountains: Use 1 quart of Quali-Pro Glyphosate T&O in 3 to 20 gallons of water per acre. West of the Rocky Mountains: Use 1 to 2 quarts of Quali-Pro Glyphosate T&O in 3 to 10 gallons of water per acre. Apply only if reduction of the bermudagrass stand can be tolerated. If burndown is necessary before harvest, allow a minimum of 21 days to ensure enough time for burndown to take place.

For suppression only (east of the Rocky Mountains): No earlier than 1 to 2 weeks after full greenup, apply 6 to 16 fluid ounces of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. If mowing before applying treatment, ensure a height of at least 3 inches is maintained. For shaded conditions or where a lesser degree of suppression is needed, apply at a rate of 6 to 10 ounces of this product plus nonionic surfactant. In areas where bermudagrass injury and stand reduction can be tolerated, sequential applications may be made when regrowth occurs.

For suppression only (west of the Rocky Mountains): No earlier

than 1 to 2 weeks after full greenup, apply 16 fluid ounces of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches tall. If mowing before applying treatment, ensure a height of at least 3 inches is maintained. In areas where bermudagrass injury and stand reduction can be tolerated, sequential applications may be made when regrowth occurs.

Cool-Season Grass Covers

For suppressing quackgrass, orchardgrass, fine fescue, and tall fescue, apply 8 fluid ounces of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Improved results may be obtained by adding ammonium sulfate to the spray solution at the rate of 2 percent by weight or 17 pounds per 100 gallons of spray solution.

For suppressing Kentucky bluegrass covers, apply 6 fluid ounces of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant. Do not add ammonium sulfate to the mix.

For optimum results, mow cool-season grass covers in spring to level their height, then apply the recommended treatment rate of Quali-Pro Glyphosate T&O at 3 to 4 days after mowing. Treatment should be avoided for cool-season grass covers under poor growing conditions, such as disease, insect damage, or drought stress (drip irrigation).

LOW-VOLUME APPLICATION (FLORIDA AND TEXAS)

For burndown or control of listed weeds, apply the recommended rates of Quali-Pro Glyphosate T&O with 0.5- to 1-percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. In areas where weed foliage is dense, use 10 to 30 gallons of water per acre.

Annual Weeds

Goatweed: Apply 2 to 3 quarts of Quali-Pro Glyphosate T&O per acre with 17 pounds of ammonium sulfate per 100 gallons of water plus 0.5- to 1-percent nonionic surfactant by total spray volume. Apply in 20 to 30 gallons of water per acre when plants are actively growing. When plants are 8 inches tall, use 2 quarts per acre. When plants are taller than 8 inches, use 3 quarts per acre. Adding Krovar II or Karmex when goatweed is taller than 8 inches may improve control. For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements.

Perennial Weeds

Apply when weeds are actively growing and are at the growth stages described in the **PERENNIAL WEEDS CONTROLLED** section of this label. After mowing, allow weeds to regrow to the recommended stage of development before treating.

WEED SPECIES	QUALI-PRO GLYPHOSATE T&O RATE PER ACRE				
	1 Qt.	2 Qts.	3 Qts.	5 Qts.	
Bermudagrass	B	—	PC	C	
Guineagrass: Texas and Florida Ridge	B	C	C	C	
Guineagrass: Florida Flatwoods	—	B	C	C	
Paragrass	B	C	C	C	
Torpedograss	S	—	PC	C	

S = Suppression PC = Partial Control B = Burndown C = Control

TREE CROPS

Citrus Fruits²: Calamondin, chironja, citron, grapefruit, kumquat, lemon, lime, mandarin orange, orange, pummelo, tangelo, tangerine, tangors.

Nuts²: Almond, beechnut, Brazil nut, butternut, cashew, chestnuts, chinquapin, filbert, hazel nut, hickory nut, macadamia, pecan, pistachio, walnut.

Pome Fruit²: Apple, loquat, mayhaw, pear, quince.

Stone Fruit²: Apricots, cherries, nectarines, olives, peaches, plums/prunes.

For use on cherries, any application equipment described in this section can be used in all states.

For use on citron and olives, only apply as a directed spray.

Any application equipment listed in this section may be used in apricots, nectarines, peaches, and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington, except for peaches grown in the states specified in the following paragraph. In all other states, use wiper equipment only.

For **PEACHES** grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee, apply only with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.**

Tropical Fruit: Acerola¹, atemoya¹, avocado¹, banana², breadfruit¹, canistel¹, carambola¹, cherimoya¹, cocoa beans¹, coffee², dates¹, figs¹, genip¹, guava¹, jaboticaba¹, jackfruit¹, longan¹, lychee¹, mango¹, mayhaw¹, papaya², passion fruit¹, persimmons¹, plantains², pomegranate¹, sapodilla¹, sapote¹, soursoy¹, sugar apple¹, tamarind¹, tea¹. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

NOTES:

¹ Wait at least 14 days after last application to harvest.

² Wait at least 3 days after last application to harvest.

³ Wait at least 17 days after last application to harvest.

⁴ Wait at least 28 days after last application to harvest.

⁵ Wait at least 1 day after last application to harvest.

VINE CROPS

Grapes: Any variety of table, wine, or raisin grape may be treated with any equipment listed in this section.

Kiwi Fruit

Applications should not be made when green shoots, canes, or foliage are in the spray zone.

Allow a minimum of 14 days between last application and harvest.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

SUPPLEMENTAL USES

The additional label information that follows applies to the specific states, applications, uses, crops, and conditions provided in each. Information within these supplemental instructions is in addition to the instructions, uses, and precautions mentioned above. **In the event of a conflict between information in the original label above and the information in the supplemental instructions below, follow the information specified in the supplemental instructions.**

Supplemental uses include the following:

1. POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE ROUNDUP READY® GENE
2. IN-CROP APPLICATIONS TO COTTON WITH THE ROUNDUP READY® GENE
3. POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE
4. CANOLA WITH THE ROUNDUP READY® GENE
5. AERIAL APPLICATIONS IN FRESNO COUNTY, CALIFORNIA ONLY (From February 15 through March 31 only)
6. AERIAL APPLICATION IN CALIFORNIA ONLY
7. AERIAL APPLICATIONS IN ARKANSAS ONLY

SUPPLEMENTAL USE # 1

POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE ROUNDUP READY® GENE

NOTE: See **GENERAL INFORMATION** and **MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS** sections of the label booklet for essential product performance information.

GENERAL INFORMATION

FOR POST-EMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

Applying this product to soybean varieties which are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene, since severe injury or destruction will result.

The Roundup Ready designation indicates that the soybean contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready soybeans may be obtained from your seed supplier.

APPLICATION INSTRUCTIONS

This product may be applied postemergence to Roundup Ready soybeans from the cracking stage throughout flowering. Allow a minimum of 14 days between final application and harvest or feeding of soybean grain, forage or hay.

MAXIMUM ALLOWABLE YEARLY RATES

Cropping Season: Do not use more than 8 quarts (256 fluid ounces) Quali-Pro Glyphosate T&O per acre.

Preplant, Preemergence: Do not use more than 5 quarts (160 fluid ounces) Quali-Pro Glyphosate T&O per acre prior to crop emergence.

In-crop: Do not use more than a total of 3 quarts (96 fluid ounces) Quali-Pro Glyphosate T&O per acre in single or multiple in-crop applications from cracking throughout the flowering stage.

Preharvest: Do not use more than 1 quart (32 fluid ounces) Quali-Pro Glyphosate T&O per acre applied after loss of green color in soybean pods until 14 days before harvest.

When applied as directed, Quali-Pro Glyphosate T&O will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications.

Precautions/Restrictions

The combined total application from crop emergence through harvest must not exceed 3 quarts (96 fluid ounces) per acre. Do not use more than 2 quarts (64 fluid ounces) Quali-Pro Glyphosate T&O per acre for any single in-crop application. Do not use a combined total of more than 2 quarts (64 fluid ounces) Quali-Pro Glyphosate T&O per acre during flowering. Allow a minimum of 14 days between final application and harvest or feeding of soybean grain, forage or hay.

There are no rotational crop restrictions following applications of this product.

For ground applications: Use the recommended rates of Quali-Pro Glyphosate T&O in 5 to 20 gallons of spray solution per acre as

a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles which provide a flat fan pattern. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of Quali-Pro Glyphosate T&O in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre unless otherwise directed. **DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. MAINTAIN APPROPRIATE BUFFER ZONES TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION.**

ANNUAL WEED RATE TABLES

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the **WEEDS CONTROLLED** sections of this label booklet for rate recommendations for specific annual weeds.

Makhteshim Agan of North America, Inc. will not warrant crop safety or weed control when Roundup Ready soybeans are treated with herbicides not specified in these supplemental instructions. Because of the potential for: 1) crop injury, 2) poor weed control from antagonism, and/or 3) rotational crop restrictions, herbicides not specified in these supplemental instructions should not be used, whether applied preemergence or applied postemergence as a tank mixture with Quali-Pro Glyphosate T&O herbicide.

Quali-Pro Glyphosate T&O may be used up to 64 fluid ounces per acre in any single application for control of annual weeds, where heavy weed densities exist.

NOTE: The following recommendations are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of 16 to 64 fluid ounces per acre of Quali-Pro Glyphosate T&O can be used to control existing weeds prior to crop emergence.

MIDWEST / MID-ATLANTIC RECOMMENDATIONS

Narrow-row or drilled soybeans: A single in-crop application of Quali-Pro Glyphosate T&O will provide effective control of labeled weeds. For best results, an initial application of 32 fluid ounces per acre on 4- to 8-inch weeds is recommended. Weeds will generally be 4 to 8 inches tall 3 to 5 weeks after planting. If the initial application is delayed and weeds are 8 to 18 inches tall, use 48 fluid ounces Quali-Pro Glyphosate T&O per acre for best results.

Under adverse growing conditions such as drought, hail, wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 24 to 32 fluid ounces per acre may be necessary to control late flushes of weeds.

Wide-row soybeans: An in-crop application of Quali-Pro Glyphosate T&O will provide effective control of the initial stand of labeled weeds.

For best results, an initial application of 32 fluid ounces per acre, on 4- to 8-inch weeds is recommended. Weeds will generally be 4 to 8 inches tall 3 to 5 weeks after planting. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

Initial and Sequential Applications (If Needed)

Weed Height (inches)	Rate (fl. oz. Quali-Pro Glyphosate T&O per acre)
1 to 3	24
4 to 8	32
8 to 18	48

Giant ragweed: Apply 32 fluid ounces Quali-Pro Glyphosate T&O per acre when weeds are 8 to 12 inches tall to avoid the need for sequential application.

Black nightshade, Pennsylvania smartweed, velvetleaf, and waterhemp: Apply 32 fluid ounces Quali-Pro Glyphosate T&O per acre to weeds 3 to 6 inches tall and 48 fluid ounces per acre when weeds are up to 12 inches tall. For morningglory species, apply 32 fluid ounces per acre when weeds are up to 4 inches tall, and 48 fluid ounces per acre when weeds are up to 6 inches tall.

Some weeds (such as black nightshade, woolly cupgrass, shatter-cane, wild proso millet, burcumber, and giant ragweed) with multiple germination times may require a sequential application of Quali-Pro Glyphosate T&O. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 24 fluid ounces of Quali-Pro Glyphosate T&O per acre for sequential applications.

SOUTHEAST RECOMMENDATIONS

Narrow-row, drilled, or wide-row soybeans: An in-crop application of Quali-Pro Glyphosate T&O will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 32 fluid ounces per acre, on 3- to 6-inch weeds is recommended. Weeds will generally be 3 to 6 inches tall 2 to 3 weeks after planting.

Initial Treatment

Weed Height (inches)	Rate (fl. oz. Quali-Pro Glyphosate T&O per acre)
3 to 6	32
6 to 12	48

Under adverse growing conditions such as drought, hail, wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of Quali-Pro Glyphosate T&O at 16 to 32 fluid ounces per acre may be necessary to control late flushes of weeds.

Sequential Application (If Needed)

Weed Height (inches)	Rate (fl. oz. Quali-Pro Glyphosate T&O per acre)
2 to 3	16
3 to 6	24
6 to 12	32

Florida pusley, hemp sesbania, and spurred anoda: Apply 32 fluid ounces per acre to weeds 2 to 4 inches for the initial application. Apply 32 fluid ounces per acre when these weeds are 3 to 6 inches tall if a sequential application is necessary.

Morningglory, black nightshade, groundcherry, and Pennsylvania smartweed: Apply 24 fluid ounces per acre on 1- to 3-inch weeds, 32 fluid ounces per acre on 3- to 6-inch weeds, or 48 fluid ounces per acre on 6- to 12-inch weeds for the initial application.

Some weeds (such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod) with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

DELTA/MID-SOUTH RECOMMENDATIONS

Narrow-row, drilled, or wide-row soybeans: An in-crop application of this product will provide effective control of the initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. For best results, an initial application of 32 fluid ounces per acre, on 2- to 4-inch weeds is recommended. Weeds will generally be 2 to 4 inches tall 2 to 3 weeks after planting.

Initial Treatment

Weed Height (inches)	Rate (fl. oz. Quali-Pro Glyphosate T&O per acre)
2 to 4	32
5 to 12	48

Sequential Application (If Needed)

Weed Height (inches)	Rate (fl. oz. Quali-Pro Glyphosate T&O per acre)
2 to 3	16
3 to 6	24
6 to 12	32

Hemp sesbania and spurred anoda: Apply a sequential treatment of 32 fluid ounces per acre on 3- to 6-inch weeds if necessary.

Some weeds (such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod) with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applica-

tions. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications.

PERENNIAL WEEDS RATE RECOMMENDATIONS

A 32 to 64 fluid ounces per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestail* (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redivine, trumpet-creeper, swamp smartweed, and wirestem muhly.

* Resistance to some biotypes has been observed. If users encounter resistant biotypes, they should contact their local extension service for tank mix recommendations.

For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with Quali-Pro Glyphosate T&O herbicide. For additional information on perennial weeds, see the **WEEDS CONTROLLED** sections of this label. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

NOTE: Nonionic surfactants which are labeled for use with post-emergence herbicides may be used. When using additional surfactant, use 0.5-percent surfactant concentration (2 quarts per 100 gallons of spray solution) for surfactants which contain at least 70 percent active ingredient or a 1-percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient.

The addition of certain surfactants to this product may result in some crop response including leaf necrosis, leaf chlorosis, or leaf speckling due to the surfactant added to the spray mixture. Read and carefully observe cautionary statements and other information appearing on the surfactant label.

SUPPLEMENTAL USE # 2

IN-CROP APPLICATIONS TO COTTON WITH THE ROUNDUP READY® GENE

NOTE: See **GENERAL INFORMATION** and **MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS** sections of the label booklet for essential product performance information. The use of additional surfactant in the spray solution may result in crop injury and reduced yield and is not recommended for over-the-top applications of this product to Roundup Ready cotton.

GENERAL INFORMATION

ATTENTION: FOR USE ONLY OVER-THE-TOP OF OR DIRECT-ED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY GENE. **NOTE:**

SEVERE INJURY OR DEATH WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN CROPS WITH THE ROUNDUP READY GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

ROUNDUP READY COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

APPLICATION INSTRUCTIONS

This product will control many troublesome weeds with over-the-top, post-directed, hooded sprayer, or preharvest applications in Roundup Ready cotton.

Maximum Allowable Yearly Rates

1. Combined total per year for all applications	8 quarts/acre
2. Preplant, Preemergence applications	5 quarts/acre
3. Total in-crop applications from cracking to lay-by	4 quarts/acre
4. Maximum preharvest application rate	2 quarts/acre

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications, apply this product in 3 to 15 gallons of water per acre.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR, UNLESS OTHERWISE DIRECTED.

FOR AERIAL APPLICATION IN CALIFORNIA OR ARKANSAS, REFER TO THE SUPPLEMENTAL USE INSTRUCTIONS AT THE END OF THIS LABEL FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS, AND REQUIREMENTS. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE. Do not apply during low-level inversion conditions, when winds are gusty, or under any other conditions which favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

There are no rotational crop restrictions following applications of this product.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready cotton. Follow the cleaning proce-

dures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations, and care should be taken to thoroughly clean all equipment prior to use.

In addition to uses listed for cotton in the **CROPPING SYSTEMS** section of this label, the following applications can be made:

Over-the-top applications: This product may be applied by aerial or ground application equipment postemergence to Roundup Ready cotton from the ground cracking stage until the four-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the four-leaf (node) stage of development may result in boll loss, delayed maturity, and/or yield loss. Any single over-the-top broadcast application should not exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four-leaf (node) stage of development. Sequential over-the-top applications of this product must be at least 10 days apart, and cotton must have at least two nodes of incremental growth between applications. **The use of additional surfactant in the spray solution may result in crop injury and reduced yield and is not recommended for over-the-top applications of this product to Roundup Ready cotton.**

NOTE: Always plant into a weed-free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16 to 48 fluid ounces per acre of this product.

Post-directed or hooded applications: This product may be applied using precision post-directed or hooded sprayers to Roundup Ready cotton through lay-by. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches). Any single post-directed application should not exceed 1 quart Quali-Pro Glyphosate T&O per acre. No more than two applications should be made from the fifth leaf through lay-by. Sequential in-crop applications of this product must be at least 10 days apart, and cotton must have at least two nodes of incremental growth between applications.

ATTENTION: USE OF QUALI-PRO GLYPHOSATE T&O IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY COTTON; HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES, AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY, AND/OR YIELD LOSS.

Salvage Treatment: This treatment may be used after the four-leaf stage of development and should only be used where weeds threat-

en to cause the loss of the crop. One quart per acre may be applied either as an over-the-top applications or as a post-directed treatments sprayed higher on the cotton plants and over the weeds.

NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY, AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to **WEEDS CONTROLLED** sections of this label. Quali-Pro Glyphosate T&O applied at 1 quart per acre will burn down or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common bermudagrass, silverleaf nightshade, trumpet creeper, and redivine. Fall preharvest applications may be required for control of these perennial weeds.

Tank mixtures with other herbicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

Preharvest applications: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20-percent boll crack. For application rates, please see the **WEEDS CONTROLLED** sections of this label booklet. This product may be applied using either aerial or ground spray equipment. Aerial and ground applications may be made up to a maximum of 2 quarts per acre. Allow a minimum of 7 days between final application and harvest of cotton or feeding of cotton forage or hay. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF QUALI-PRO GLYPHOSATE T&O TO ROUNDUP READY COTTON IS PROHIBITED. NOTE:** Quali-Pro Glyphosate T&O will not enhance the performance of harvest aids when applied to Roundup Ready cotton. **DO NOT APPLY QUALI-PRO GLYPHOSATE T&O PREHARVEST TO CROPS GROWN FOR SEED.**

SUPPLEMENTAL USE # 3

FOR POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES OTHER THAN CORN WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See **GENERAL INFORMATION** and **MIXING** sections of this label booklet for essential product performance information.

GENERAL INFORMATION

USE THIS PRODUCT ONLY ON CORN HYBRIDS DESIGNATED AS CONTAINING THE ROUNDUP READY GENE.

Applying this product to corn hybrids which are not designated as Roundup Ready will result in severe crop injury and yield loss.

The Roundup Ready designation indicates that the corn contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready corn may be obtained from your seed supplier.

APPLICATION INSTRUCTIONS

This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of this product must not exceed 1 quart per acre. Sequential in-crop applications of this product from emergence through the V8 stage or 30 inches must not exceed 2 quarts per acre per growing season.

MAXIMUM YEARLY RATES ALLOWED

Preplant: Do not apply more than 5 quarts per acre prior to crop emergence.

In-crop: Do not use more than a combined total of 2 quarts Quali-Pro Glyphosate T&O per acre in multiple in-crop applications from emergence through the V8 stage or 30 inches.

Preharvest: Do not apply more than 1 quart Quali-Pro Glyphosate T&O per acre after maximum kernal fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest.

Cropping Season: Do not exceed 8 quarts Quali-Pro Glyphosate T&O per acre as a combined total per year for all applications.

When applied as directed, this product controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Applications should be made to actively growing weeds before they reach the maximum size listed in the **WEEDS CONTROLLED** sections of this label booklet. Refer to the **MIXING** section of the label booklet for proper use instructions.

The addition of 1- to 2-percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet, Micro-Tech, or Partner herbicides. For all tank mixes, review the tank mix herbicides' individual product labels for specific crops, application rates, geographical restrictions, and precautionary statements. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including fertilizers and micro-nutrients, are not recommended with this product since this may result in increased potential for crop injury.

Allow a minimum of 50 days between application of this product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 10 days between in-crop applications of this product. There are no rotational crop restrictions following applications of this product.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See **WEEDS CONTROLLED** sections on this label. AVOID DRIFT—DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

WEED-CONTROL RECOMMENDATIONS

Apply 24 to 32 fluid ounces of Quali-Pro Glyphosate T&O per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the **WEEDS CONTROLLED** sections of this label booklet for rate recommendations for specific annual weeds. Quali-Pro Glyphosate T&O applied at up to 1 quart per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpet creeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the **PERENNIAL WEED** section of the **WEEDS CONTROLLED** sections of this label booklet.

Preemergence Followed by Postemergence Weed-Control Program

This product may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. A single in-crop application of this product at the recommended rate will provide control of emerged weeds listed on the label. This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first.

Postemergence Only Weed-Control Program

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first.

This product may be applied in tank mixture with a labeled rate of Harness™, Harness Xtra, Harness Xtra 5.6L, Micro-Tech, Bullet, Partner, Permit™, or atrazine herbicides. Refer to the specific product label and observe all precautionary statements and limitations on the label for all products used in tank mixtures, including specific crops, application timing restrictions, application rate, geographical restrictions, soil restrictions, minimum re-cropping interval, and rotational guidelines—the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

Tank Mix Partner	Maximum Height Of Corn For Application
Harness	11 inches
Harness Xtra	
Harness Xtra 5.6L	
Bullet*	5 inches
Micro-Tech*	
Partner*	
Permit	24 inches
Atrazine	12 inches

*Bullet, Micro-Tech and Partner are not registered for use as a postemergence application in Texas.

Bullet, Harness, Micro-Tech and Partner are registered trademarks of Monsanto Company. Permit is a trademark of, and used under license from, Nissan Chemical Industries, Ltd.

SUPPLEMENTAL USE # 4

CANOLA WITH THE ROUNDUP READY® GENE

See **GENERAL INFORMATION** and **MIXING** sections of this label booklet for essential product performance information.

GENERAL INFORMATION

USE ONLY ON CANOLA WHICH CONTAINS THE ROUNDUP READY GENE. DO NOT USE THIS PRODUCT ON CANOLA WITH THE ROUNDUP READY GENE PLANTED IN THE FOLLOWING

STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA, AND WEST VIRGINIA.

Applying this product to canola which is not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene since severe injury or destruction will result.

The Roundup Ready designation indicates the canola contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready canola may be obtained from your seed supplier.

USE RECOMMENDATIONS

This product will control many troublesome emerged weeds when applied preplant, preemergent, and/or with over-the-top applications in Roundup Ready canola. Allow a minimum of 60 days between last application and canola harvest.

Maximum Allowable Combined Yearly Rates

Preplant and preemergence applications	2 quarts/acre
Total in-crop application from emergence to 6 leaf	1 quart/acre

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

For aerial applications apply this product in 3 to 15 gallons of water per acre.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE. DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine-spray

droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

There are no rotational crop restrictions following applications of this product.

Sprayer Preparation

It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready canola. Follow the cleaning procedures specified on the label of the product(s) previously used. Canola can be very sensitive to many herbicides at extremely low concentrations, and care should be taken to thoroughly clean all equipment prior to use.

Preplant or Preemergent Applications

This product may be applied by aerial or ground application equipment prior to planting or emergence of canola. The maximum combined application rate from all preplant and preemergent applications should not exceed 2 quarts per acre per season.

NOTE: In no-till and stale seedbed systems, always use a burndown treatment to control existing weeds before canola emerges. Apply a preplant burndown treatment of 16 to 32 fluid ounces per acre of this product.

Over-the-top Applications

This product may be applied by aerial or ground application equipment postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Single Application: Apply 16 to 24 ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications which may result in temporary yellowing, delayed flowering, and or growth reduction. Similar injury may result when applications of more than 16 ounces per acre are applied after the 4-leaf stage.

Sequential Applications: Apply 16 ounces per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications are recommended for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass.

This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 32 ounces per acre.

WEED-CONTROL RECOMMENDATIONS

For specific rates of application and instructions for control of various annual and perennial weeds, refer to the **WEEDS CONTROLLED** sections of this label booklet.

Tank mixtures with other herbicides, insecticides, or fungicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

SUPPLEMENTAL USE # 5

AERIAL APPLICATIONS IN FRESNO COUNTY, CALIFORNIA ONLY

(From February 15 through March 31 only)

NOTE: For aerial application outside these dates, refer to the **AERIAL APPLICATION IN CALIFORNIA ONLY** statewide supplemental label below.

See the **GENERAL INFORMATION** and **MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS** sections of this label booklet for essential product information.

See the **CROPPING SYSTEMS** section of this label booklet for specific recommendations on the use of this product.

APPLICABLE AREA

This supplement applies only to the area contained inside the following boundaries within Fresno County, California.

North:	Fresno County line
South:	Fresno County line
East:	State Highway 99
West:	Fresno County line

GENERAL INFORMATION

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of Quali-Pro Glyphosate T&O. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Recommendations

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of Quali-Pro Glyphosate T&O is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at Night: Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

SUPPLEMENTAL USE # 6

AERIAL APPLICATION IN CALIFORNIA ONLY

See **GENERAL INFORMATION** and **MIXING** sections of this label booklet for essential product information.

See the **CROPS** section of this label booklet for specific recommendations on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL "ROUNDUP READY" CROPS), PLANTS, TREES, OR OTHER DESIRABLE VEGETATION, SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Aerial applications of this product are allowed in the following situations:

1. In fallow and reduced-tillage systems prior to the emergence or transplanting of labeled crops.
2. In alfalfa and pasture renovation applications.
3. Application to brush and chaparral. Refer to the current supplemental label for directions for this use.
4. Preharvest in alfalfa, corn, cotton, wheat, and Roundup Ready corn. Observe this label's specific preharvest application instructions for each individual crop.

Do not plant subsequent crops other than those listed in this label booklet for 30 days following application.

When applied as recommended under the conditions described, this

product controls annual and perennial weeds listed in this label booklet.

DO NOT EXCEED A MAXIMUM RATE OF 2 QUARTS PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED-TILLAGE SYSTEMS, ALFALFA AND PASTURE RENOVATION AND BRUSH AND CHAPARRAL APPLICATIONS.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN ALFALFA, CORN, COTTON, WHEAT, AND ROUNDUP READY CORN PRIOR TO HARVEST.

AERIAL EQUIPMENT

Use the recommended rates of this product in 3 to 15 gallons of water per acre. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

AVOID DRIFT—DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine-spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift-control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure Uniform Application: To avoid streaking, uneven, or overlapped application, use appropriate marking devices. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

SUPPLEMENTAL USE # 7

AERIAL APPLICATIONS IN ARKANSAS ONLY

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS, AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the **GENERAL INFORMATION** and **MIXING** sections of this label booklet for essential product performance information.

USE DIRECTIONS

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE-SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment setup to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan or rotor. In many cases, reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the airstream and never discharge downwards more than 45 degrees on fixed-wing aircraft or forward of the prevailing air flow on rotary-winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when winds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine-spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Keep container closed to prevent spills and contamination. Store product in original container only.

PESTICIDE DISPOSAL: Wastes that result from using Quali-Pro Glyphosate T&O that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal, or in accordance with all applicable Federal, state, or local procedures. Empty containers retain vapor and product residues. Follow all labeled safeguards until container is cleaned, reconditioned, or destroyed.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable Container: Refillable container. Refill this container with glyphogan only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Disposal of this container must be in compliance with state and local regulations.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product.

Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro is a registered trademark of Makhteshim Agan of North America, Inc.

Bullet, Lasso, Micro-Tech, Partner, and Lariat are registered trademarks of Monsanto Company.

Oust, Krovar, Escort, Telar, Canopy, Gemini, Lexone, Lorox Plus, Preview, and Karmex are trademarks of E.I. duPont de Nemours and Company.

Dual MAGNUM, Bicep MAGNUM, Princep, and Caliber are trademarks of Novartis Corporation.

Goal is a registered trademark of Dow Agrosciences.

Surflan is a registered trademark of UPL.

Banvel and Solicam are trademarks of BASF, Inc.

Ronstar, Folex, and Prep are trademarks of Rhone-Poulenc, Inc.

Sencor and Turbo are trademarks of Bayer AG.

Prowl, Pursuit, Pursuit Plus, Scepter, and Squadron are trademarks of American Cyanamid Company.

Command is a trademark of FMC Corporation.

DEF is a trademark of Mobay Chemical Company.

Microfoil is a trademark of Union Carbide Agricultural Products.

Thru-Valve is a trademark of Waldrum Specialties, Inc.

QUALI-PRO[®]

Glyphosate T&O

Herbicide

ACTIVE INGREDIENT:	% BY WT.
*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt	41.0%
OTHER INGREDIENTS:	59.0%
TOTAL	100.0%

*This product contains 480 grams per liter or four pounds per U.S. gallon of glyphosate—the active ingredient—in the form of its isopropylamine salt. This equals 356 grams per liter, or three pounds per U.S. gallon of the acid, glyphosate.

EPA Reg. No. 66222-208

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Immediately hold eyelids open and flush with plenty of water for at least 15 minutes.• Remove contact lenses, if present, after first 5 minutes then continue rinsing eyes.• Get medical attention.
IF INHALED:	<ul style="list-style-type: none">• Remove individual to fresh air.• If not breathing, give artificial respiration, preferably mouth-to-mouth.• Get medical attention.
IF SWALLOWED:	<ul style="list-style-type: none">• This product will cause gastrointestinal tract irritation.• Immediately dilute by swallowing water or milk.• Get medical attention.• NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.
HAVE THE PRODUCT CONTAINER OR LABEL WITH YOU WHEN CALLING A POISON CONTROL CENTER OR DOCTOR, OR GOING FOR TREATMENT.	
For medical emergencies involving this product, call Prosar at 1-877-250-9291.	

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

EPA 120605/Notif 012910/Trans 010410/Rev C

EPA Est. No. 37429-GA-001^{BT}; 37429-GA-002^{BO}; 5905-AR-001^{HA}
Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

Net Contents: 2.5 Gallons