

# Specimen Label

## AquaPier<sup>\*</sup> Aquatic Herbicide



AquaPier may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. AquaPier may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water. AquaPier may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites.

### ACTIVE INGREDIENT

Isopropylamine salt of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl-5-oxo-1*H*-imidazol-2-yl)-3-pyridinecarboxylic acid]<sup>†</sup> . . . . . 27.8%

OTHER INGREDIENTS . . . . . 72.2%

TOTAL . . . . . 100.0%

<sup>†</sup>Equivalent to 22.6% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl-5-oxo-1*H*-imidazol-2-yl)-3 pyridinecarboxylic acid or 2 pounds acid equivalent per gallon.

## Keep Out of Reach of Children CAUTION/PRECAUCIÓN

**PRECAUCIÓN AL USUARIO:** Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**CAUTION!** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing and wash before reuse.

First Aid	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor for further treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• DO NOT induce vomiting unless told to by a poison control center or doctor.</li><li>• DO NOT give anything to an unconscious person.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>EMERGENCY NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

Refer to inside of label booklet for additional **Precautionary Information** and **Directions for Use** including **Storage and Disposal**.

**Notice:** Read the entire label before using. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer", "Terms of Sale" and "Limitation of Liability" inside label booklet.

If you wish to obtain additional product information, please visit our web site at [www.sepro.com](http://www.sepro.com).

EPA Reg. No. 74477-6-67690  
FPL011507

<sup>\*</sup>Trademark SePRO Corporation, Carmel, IN 46032 U.S.A.

## Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves made of any waterproof material; and
- Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are given for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENVIRONMENTAL HAZARDS

DO NOT apply to water except as specified in this label. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss may cause the suffocation of some aquatic organisms. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift precautions on the label.

## PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of AquaPier should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

**DO NOT** mix, store or apply AquaPier or spray solutions of AquaPier in unlined steel (except stainless steel) containers or spray tanks.

## Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Keep containers closed to avoid spills and contamination.

## GENERAL INFORMATION

**AquaPier** is an aqueous solution intended to be mixed with water and surfactant(s) for application to control floating and emergent undesirable vegetation (see the *AQUATIC SPECIES CONTROLLED BY AQUAPIER* and *ADDITIONAL WEEDS CONTROLLED BY AQUAPIER* sections) in or near bodies of water which may be flowing, non-flowing or transient. AquaPier may be applied to aquatic sites that include lakes, rivers, streams, ponds, seeps, drainage ditches, canals, reservoirs, swamps, bogs, marshes, estuaries, bays, brackish water,

transitional areas between terrestrial and aquatic sites and seasonal wet areas. See the *AQUATIC USES, PRECAUTIONS AND RESTRICTIONS* section of this label for precautions, restrictions, and instructions on aquatic uses. AquaPier may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites (see the *AERIAL APPLICATIONS* and *GROUND APPLICATIONS* sections for additional details).

**DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow aquatic organisms to move into untreated areas.

Read and observe the following directions if aquatic sites are present in terrestrial non-crop areas and are part of the intended treatment area:

AquaPier controls most annual and perennial grasses, broadleaf weeds and many brush and vine species. AquaPier will provide some residual control of undesirable species that germinate above the waterline. AquaPier controls vegetation by absorption through emergent leaves and stems, from which it is translocated throughout the plant, where it accumulates in rapidly-growing meristematic tissue. Treated plants stop growing soon after treatment. Chlorosis (yellowing of plant tissue) first appears in the newest leaves, and necrosis spreads from this point. In perennials, AquaPier is translocated into and kills underground or submerged storage organs to prevent regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two or more weeks after application. Complete kill of plants may not occur for several weeks. Applications of AquaPier are rainfast within one hour after treatment.

AquaPier will not control plants which are completely submerged or have a majority of their foliage under water.

AquaPier must be applied to the emergent foliage of the target vegetation, since it has little-to-no activity on submerged vegetation. Residual concentrations of AquaPier from direct application to water are not expected to be sufficient to provide control of undesirable vegetation. Apply AquaPier so as to maximize spray contact onto the target vegetation while minimizing overspray onto adjacent or underlying water. For optimum performance, include a surfactant (see the *ADJUVANTS* section for specific recommendations) in the spray solution and apply to actively growing weeds. AquaPier may be applied in spot treatments by using low-volume directed application techniques or broadcast by using ground equipment, water craft or helicopter.

Apply AquaPier with surface or helicopter application equipment in at least 5 gallons of water per acre. When applying by helicopter, follow directions under the *AERIAL APPLICATIONS* section of this label; otherwise, refer to the section on *GROUND APPLICATIONS* when using surface equipment.

Applications to moving bodies of water should be made while traveling upstream to prevent concentration of this herbicide in water. **DO NOT** apply to bodies of water or portions of bodies of water where emergent and/or floating weeds do not exist.

When target vegetation covers a large percentage of the surface area of impounded water, apply AquaPier to alternating strips (i.e., **DO NOT** apply to every other swath) to avoid oxygen depletion due to decaying vegetation, since oxygen depletion may result in the suffocation of sensitive aquatic organisms. **DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days before retreatment. Begin treatment along the shore and proceed outward in alternating swaths as described above to allow aquatic organisms to move into untreated areas.

Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash for one hour after application.

Apply AquaPier at 2 to 6 pints per acre, depending on weed species and weed density. Use higher rates for heavy weed pressure. Refer to the *AQUATIC SPECIES CONTROLLED BY AQUAPIER* section and the *ADDITIONAL WEEDS CONTROLLED BY AQUAPIER* section of this label for specifics. **DO NOT** exceed 6 pints per acre (1.5 lb ai/A) per year.

AquaPier may also be applied as a draw down treatment in areas described above. Apply AquaPier to weeds after water has been drained and allow 14 days before reintroduction of water.

## PRECAUTIONS FOR AVOIDING INJURY TO NON-TARGET PLANTS

AquaPier can occasionally affect non-target or desirable vegetation by root uptake of the herbicide from treated soil. Injury or loss of non-target plants may result if AquaPier is applied onto or near desirable plants, or to areas where their roots extend, or in areas where treated soil may be washed or moved into contact with their root zone.

When making applications along shorelines where desirable plants may be present, exercise caution to avoid spray contact onto their foliage or spray application to the soil in which they are rooted. Shoreline plants with roots that extend into the waters where AquaPier has been applied will generally not be adversely affected.

If treated vegetation is to be removed from the application site, **DO NOT** use the vegetative matter as mulch or compost on or around desirable species.

**IMPORTANT: DO NOT** use on food crops. **DO NOT** apply this product within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water, such as a lake, pond or reservoir. **DO NOT** apply to water used for irrigation except as described in *APPLICATION TO WATERS USED FOR IRRIGATION* section of this label. Keep away from fertilizers, insecticides, fungicides, and seeds.

**DO NOT** drain or flush equipment on or near desirable plants, or onto areas where their roots may extend, or in locations where the chemical or treated soil may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, drive-ways, tennis courts or similar areas where roots of desirable vegetation may extend and be exposed to potential injury and/or mortality from root uptake of AquaPier. **DO NOT** side trim desirable vegetation with this product, unless severe injury or plant death is acceptable. Prevent drift of spray to desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-crop weed control is not within the scope of the Worker Protection Standard. See the *GENERAL INFORMATION* section of this label for a description of non-crop sites.

**DO NOT** enter treated areas without protective clothing until sprays have dried.

## SPRAY DRIFT MANAGEMENT

**Spray Drift:** Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator and the entity authorizing spraying are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. **DO NOT** apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, and temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces the drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see *Wind, Temperature and Humidity, and Temperature Inversions*).

### Controlling Droplet Size:

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. **DO NOT** use nozzles producing a mist droplet spray.

**Application Height:** Making applications at the lowest possible height (helicopter, ground driven spray bloom) that is safe and practical reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the treatment area, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

**Wind:** Drift potential is lowest between wind speeds of 3 - 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential.

**NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during temperature inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and

moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Wind Erosion:** Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

### Aerial Application Methods and Equipment (Helicopters ONLY):

- **Water Volume:** Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to produce accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

### Managing Spray Drift from Aerial Applications:

Applicators must follow these requirements to avoid off-target drift movement:

- 1) **Boom length** – the distance of the outermost nozzles on the boom must not exceed 3/4 the length of the rotor;
- 2) **Nozzle orientation** – nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees; and
- 3) **Application height** – without compromising helicopter safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants. Applicators must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

### Ground Application (Broadcast):

- **Water Volume:** Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

## AQUATIC SPECIES CONTROLLED BY AQUAPIER

AquaPier will control the following target species when used as recommended in the table below. Rate recommendations are expressed as product volume for broadcast applications and as a percent (%) solution for directed applications including spot treatments. For % solution applications, **DO NOT** apply more than the equivalent of 3 quarts of AquaPier per acre.



## AQUATIC SPECIES

Common Name	Scientific Name	Recommendations
<b>Floating Species</b>		
Duckweed†	<i>Lemna minor</i>	2 - 3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Duckweed, Giant†	<i>Spirodela polyrriza</i>	2 - 3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Frogbit†	<i>Limnobium spongia</i>	1 - 2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Spatterdock†	<i>Nuphar luteum</i>	Apply a tank-mix of 2 - 4 pints/acre AquaPier plus 4 - 6 pints/acre glyphosate (0.5% AquaPier plus 1.5% glyphosate) in 100 GPA water for best control. Completely cover all actively-growing emergent foliage.
Water Hyacinth†	<i>Eichhornia crassipes</i>	1 - 2 pints/acre (0.5% solution) in 100 GPA water to actively growing foliage.
Water Lettuce†	<i>Pistia stratiotes</i>	1 - 2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.

## Emerged Species

Alligatorweed†	<i>Alternanthera philoxeroides</i>	1 - 4 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. <b>DO NOT</b> apply as a tank-mix with glyphosate, since doing so may reduce control.
Arrowhead†, Duck-potato†	<i>Sagittaria</i> spp.	1 - 2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Bacopa, lemon†	<i>Bacopa</i> spp.	1 - 2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Parrot feather†	<i>Myriophyllum aquaticum</i>	Apply 2 - 4 pints in 100 GPA water to actively growing emergent foliage. Foliage must be above water for sufficient uptake.
Pennywort†	<i>Hydrocotyle</i> spp.	1 - 2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Pickereelweed†	<i>Pontederia cordata</i>	2 - 3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.

## AQUATIC SPECIES

Common Name	Scientific Name	Recommendations
<b>Emerged Species</b>		
Taro, wild†; Dasheen†; Elephant's Ear†; Cocco Yam†	<i>Colocasia esculentum</i>	4 - 6 pints/acre (1.5% solution) in 100 GPA with a high quality 'sticker' adjuvant. Completely cover all actively-growing emergent foliage.
Water lily†	<i>Nymphaea odorata</i>	2 - 3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
Water Primrose†	<i>Ludwigia uruguayensis</i>	4 - 6 pints/acre (1.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. <b>DO NOT</b> apply as a tank-mix with glyphosate, since doing so may reduce control.

## Terrestrial/Marginal

Soda Apple, aquatic†; Nightshade†	<i>Solanum tampicense</i>	2 pints/acre applied to foliage.
Bamboo, Japanese†	<i>Phyllostachys</i> spp.	3 - 4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Brazilian Pepper; Christmasberry	<i>Schinus terebinthifolius</i>	2 - 4 pints/acre applied to foliage.
Cattail	<i>Typha</i> spp.	2 - 4 pints (1% solution) applied to actively-growing, green foliage after full leaf elongation. Lower rates will control cattail in the north; higher rates are needed in the south.
Chinese Tallow Tree	<i>Sapium sebiferum</i>	16 - 24 oz applied to foliage.
Cogon grass	<i>Imperata cylindrica</i>	Burn foliage, till area, then treat in fall at 2 quarts/acre AquaPier plus MSO applied to new growth.
Cordgrass, prairie	<i>Spartina</i> spp.	4 - 6 pints applied to actively growing foliage.
Cutgrass†	<i>Zizaniopsis miliacea</i>	4 - 6 pints applied to actively growing foliage.
Elephant Grass†; Napier Grass†	<i>Pennisetum purpureum</i>	3 pints/acre applied to actively growing foliage.
Flowering rush†	<i>Butumu typha</i>	2 - 3 pints applied to actively growing foliage.
Giant Reed, Wild Cane	<i>Arundo donax</i>	4 - 6 pints/acre applied in spring to actively growing foliage.

AQUATIC SPECIES		
Common Name	Scientific Name	Recommendations
<b>Terrestrial/Marginal</b>		
Golden Bamboo <sup>†</sup>	<i>Phyllostachys aurea</i>	3 - 4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Junglerice	<i>Echinochloa colonum</i>	3 - 4 pints applied to actively growing foliage.
Knapweeds	<i>Centaurea species</i>	Russian Knapweed - Apply 2 - 3 pints plus 1 quart/acre MSO in the fall after senescence begins.
Knotweed, Japanese (see Fallopia japonica)	<i>Polygonum cuspidatum</i>	3 - 4 pints/acre applied postemergence to actively growing foliage.
Melaleuca; Paperbark Tree	<i>Melaleuca quinquenervia</i>	For established stands, apply 6 pints/acre AquaPier plus 6 pints/acre glyphosate plus adjuvant. For best results, use 4 quarts/A methylated seed oil for adjuvant. For ground foliar application, uniformly apply to completely cover all actively-growing foliage. For aerial foliar application, apply in at least two passes in a cross pattern at 10 GPA. For spot treatment, apply a 25% solution of AquaPier plus 25% solution of glyphosate plus 1.25% MSO in water as a frill or stump treatment.
Nutgrass <sup>†</sup> ; Kili'p'opu <sup>†</sup>	<i>Cyperus rotundus</i>	2 pints AquaPier plus 1 quart/acre MSO applied early postemergence.
Nutsedge <sup>†</sup>	<i>Cyperus</i> spp.	2 - 3 pints postemergence to foliage or pre-emergence incorporated. Non-incorporated preemergence applications will not provide control.
Phragmites; Common Reed	<i>Phragmites australis</i>	4 - 6 pints/acre applied to actively-growing, green foliage after leaf elongation. Completely cover all actively-growing foliage. If stand has a substantial amount of old stem tissue, mow or burn, allow to regrow to approximately 5' tall before treatment. Lower rates will control phragmites in the north; higher rates are needed in the south.
Poison Hemlock <sup>†</sup>	<i>Conium maculatum</i>	2 pints AquaPier plus 1 quart/ acre MSO applied preemergence or early postemergence up to rosette stage before flowering.

AQUATIC SPECIES		
Common Name	Scientific Name	Recommendations
<b>Terrestrial/Marginal</b>		
Purple Loosestrife	<i>Lythrum salicaria</i>	1 pint/acre applied to actively growing foliage.
Reed canarygrass	<i>Phalaris arundinacea</i>	3 - 4 pints/acre applied to actively growing foliage.
Rose, swamp	<i>Rosa palustris</i>	2 - 3 pints/acre applied to actively growing foliage.
Russian-Olive	<i>Elaeagnus angustifolia</i>	2 - 4 pints/acre or 1% solution applied to foliage.
Saltcedar; Tamarisk	<i>Tamarix species</i>	Aerially apply 2 quarts AquaPier plus 0.25% v/v NIS to actively growing foliage during flowering. For spot spraying, use a 1% solution of AquaPier plus 0.25% v/v NIS and spray to wet foliage. After application wait at least two years before disturbing treated saltcedar. Earlier disturbance can reduce overall control.
Smartweed	<i>Polygonum</i> spp.	2 pints/acre applied early postemergence.
Sumac	<i>Rhus</i> spp.	2 - 3 pints/acre applied to foliage.
Swamp Morning Glory; Water Spinach; Kangkong	<i>Ipomoea aquatica</i>	1 - 2 pints/acre AquaPier plus 1 quart/acre MSO applied at early postemergence.
Torpedo grass	<i>Panicum repens</i>	4 pints/acre (1 - 1.5% solution). Completely cover all actively growing foliage.
White Top <sup>†</sup> ; Hoary Cress <sup>†</sup>	<i>Cardaria draba</i>	1 - 2 pints/acre applied in spring to foliage during flowering.
Willow	<i>Salix</i> spp.	2 - 3 pints/acre AquaPier. Completely cover all actively growing foliage.

† Not approved for use in California.

## ADDITIONAL WEEDS CONTROLLED BY AQUAPIER

When used as recommended on terrestrial sites, AquaPier will provide preemergence or postemergence control with residual control of the following vegetation species. Annual weeds may be controlled by preemergence or postemergence applications of AquaPier. For established biennials and perennial vegetation control, postemergence treatments of AquaPier are recommended.

The tables below show broadcast rates and indicate relative weed sensitivity. It is important to consider relative weed sensitivity when preparing low volume spray solutions (see *Low Volume* section of *GROUND APPLICATIONS* section), since low volume treatments apply less AquaPier per acre than is shown for the broadcast treatments.

The relative sensitivity of the species listed below can also be used to determine the relative risk of causing non-target plant injury if any of those species are considered to be desirable within the treatment area.

**Resistant Biotypes:** Some weeds listed below may have naturally-occurring biotypes (plants within a given species that have a slightly different but distinct genetic makeup from other plants of that species) that are not effectively controlled by this and/or other herbicides (Oust®) with the ALS/AHAS enzyme-inhibiting mode of action. If naturally-occurring ALS/AHAS resistant biotypes are present in area, AquaPier should be tank-mixed or applied sequentially with a registered herbicide that depends on a different mode of action to ensure control.

GRASSES		
Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 2 - 3 pints per acre <sup>1</sup></b>		
Annual bluegrass	<i>Poa annua</i>	A
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	A
Canada bluegrass	<i>Poa compressa</i>	P
Downy brome	<i>Bromus tectorum</i>	A
Fescue	<i>Festuca</i> spp.	A/P
Foxtail	<i>Setaria</i> spp.	A
Italian ryegrass	<i>Lolium multiflorum</i>	A
Johnsongrass	<i>Sorghum halepense</i>	P
Kentucky bluegrass	<i>Poa pratensis</i>	P
Lovegrass	<i>Eragrostis</i> spp.	A/P
Napier grass <sup>†</sup>	<i>Pennisetum purpureum</i>	P
Orchardgrass	<i>Dactylis glomerata</i>	P
Paragrass	<i>Brachiaria mutica</i>	P
Quackgrass	<i>Agropyron repens</i>	P
Sandbur	<i>Cenchrus</i> spp.	A
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Smooth brome	<i>Bromus inermis</i>	P
Vaseygrass	<i>Paspalum urvillei</i>	P
Wild Oats	<i>Avena fatua</i>	A
Witchgrass	<i>Panicum capillare</i>	A
<b>Apply 3 - 4 pints per acre <sup>1</sup></b>		
Barnyardgrass	<i>Echinochloa crus-gali</i>	A
Beardgrass	<i>Andropogon</i> spp.	P
Bluegrass, Annual	<i>Poa annua</i>	A
Bulrush <sup>†</sup>	<i>Scirpus validus</i>	P
Cheat	<i>Bromus secalinus</i>	A
Crabgrass	<i>Digitaria</i> spp.	A
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	A
Fall Panicum	<i>Panicum dichotomiflorum</i>	A
Goosegrass	<i>Eleusine indica</i>	A
Itchgrass	<i>Rottboellia exaltata</i>	A
Lovegrass	<i>Eragrostis</i> spp.	A
Maidencane <sup>†</sup>	<i>Panicum hemitomon</i>	A
Panicum, Browntop	<i>Panicum fasciculatum</i>	A
Panicum, Texas	<i>Panicum texanum</i>	A
Prairie threeawn	<i>Aristida oligantha</i>	P
Sandbur, Field	<i>Cenchrus incertus</i>	A
Signalgrass	<i>Brachiaria platyphylla</i>	A
Wild barley	<i>Hordeum</i> spp.	A
Wooly Cupgrass	<i>Eriochloa villosa</i>	A

GRASSES		
Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 4 - 6 pints per acre <sup>1</sup></b>		
Bahiagrass	<i>Paspalum notatum</i>	P
Bermudagrass <sup>3</sup>	<i>Cynodon dactylon</i>	P
Big bluestem	<i>Andropogon gerardii</i>	P
Dallisgrass	<i>Paspalum dilatatum</i>	P
Feathertop	<i>Pennisetum villosum</i>	P
Guineagrass	<i>Panicum maximum</i>	P
Saltgrass <sup>3</sup>	<i>Distichlis stricta</i>	P
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Sprangletop	<i>Leptochloa</i> spp.	A
Timothy	<i>Phleum pretense</i>	P
Wirestem muhly	<i>Muhlenbergia frondosa</i>	P

<sup>1</sup> The higher rates should be used where heavy or well-established infestations occur.

<sup>2</sup> Growth Habit – A = Annual, B = Biennial, P = Perennial

<sup>3</sup> Use a minimum of 75 GPA - Control of established stands may require repeat applications.

† Not approved for use in California.

BROADLEAF WEEDS		
Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 2 - 3 pints per acre <sup>1</sup></b>		
Burdock	<i>Arctium</i> spp.	B
Carpetweed	<i>Mollugo verticillata</i>	A
Carolina geranium	<i>Geranium carolinianum</i>	A
Clover	<i>Trifolium</i> spp.	A/P
Common chickweed	<i>Stellaria media</i>	A
Common ragweed	<i>Ambrosia artemisiifolia</i>	A
Dandelion	<i>Taraxacum officinale</i>	P
Dog fennel	<i>Eupatorium capillifolium</i>	A
Filaree	<i>Erodium</i> spp.	A
Fleabane	<i>Erigeron</i> spp.	A
Hoary vervain	<i>Verbena stricta</i>	P
Indian mustard	<i>Brassica juncea</i>	A
Kochia	<i>Kochia scoparia</i>	A
Lambsquarters	<i>Chenopodium album</i>	A
Lespedeza <sup>†</sup>	<i>Lespedeza</i> spp.	P
Miners lettuce	<i>Montia perfoliata</i>	A
Mullein	<i>Verbascum</i> spp.	B
Nettleleaf goosefoot	<i>Chenopodium murale</i>	A
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>	P
Pepperweed	<i>Lepidium</i> spp.	A
Pigweed	<i>Amaranthus</i> spp.	A
Puncturevine	<i>Tribulus terrestris</i>	A
Russian thistle	<i>Salsola kali</i>	A
Smartweed	<i>Polygonum</i> spp.	A/P
Sorrell	<i>Rumex</i> spp.	P
Sunflower	<i>Helianthus</i> spp.	A
Sweet clover	<i>Melilotus</i> spp.	A/B
Tansymustard	<i>Descurainia pinnata</i>	A
Western ragweed	<i>Ambrosia psilostachya</i>	P
Wild carrot	<i>Daucus carota</i>	B
Wild lettuce	<i>Lactuca</i> spp.	A/B
Wild parsnip	<i>Pastinaca sativa</i>	B
Wild turnip	<i>Brassica campestris</i>	B
Woollyleaf bursage	<i>Franseria tomentosa</i>	P
Yellow woodsorrel	<i>Oxalis stricta</i>	P

## BROADLEAF WEEDS

Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 3 - 4 pints per acre <sup>1</sup></b>		
Broom snakeweed <sup>4</sup>	<i>Gutierrezia sarothrae</i>	P
Bull thistle	<i>Cirsium vulgare</i>	B
Burclover	<i>Medicago</i> spp.	A
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	A
Clover, Hop	<i>Trifolium procumbens</i>	A
Cocklebur	<i>Xanthium strumarium</i>	A
Cudweed	<i>Gnaphalium</i> spp.	A
Desert Camelthorn	<i>Alhagi pseudalhagi</i>	P
Dock	<i>Rumex</i> spp.	P
Fiddleneck	<i>Amsinckia intermedia</i>	A
Goldenrod	<i>Solidago</i> spp.	P
Henbit	<i>Lamium alexiclaude</i>	A
Knotweed, prostrate	<i>Polygonum aviculare</i>	A/P
Pokeweed	<i>Phytolacca americana</i>	P
Purslane	<i>Portulaca</i> spp.	A
Pusley, Florida	<i>Richardia scabra</i>	A
Rocket, London	<i>Sisymbrium irio</i>	A
Rush skeletonweed <sup>4</sup>	<i>Chondrilla juncea</i>	B
Saltbush	<i>Atriplex</i> spp.	A
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	A
Spurge, Annual	<i>Euphorbia</i> spp.	A
Stinging nettle <sup>4</sup>	<i>Urtica dioica</i>	P
Velvetleaf	<i>Abutilon theophrasti</i>	A
Yellow starthistle	<i>Centaurea solstitialis</i>	A

### Apply 4 - 6 pints per acre <sup>1</sup>

Arrowwood	<i>Pluchea sericea</i>	A
Canada thistle	<i>Cirsium arvense</i>	P
Giant ragweed	<i>Ambrosia trifida</i>	A
Grey rabbitbrush	<i>Chrysothamus nauseosus</i>	P
Little mallow	<i>Malva parviflora</i>	B
Milkweed	<i>Asclepias</i> spp.	P
Primrose	<i>Oenothera kunthiana</i>	P
Silverleaf nightshade	<i>Solanum eleagnifolium</i>	P
Sowthistle	<i>Sonchus</i> spp.	A
Texas thistle	<i>Cirsium texanum</i>	P

1 The higher rates should be used where heavy or well-established infestations occur.

2 Growth Habit – A = Annual, B = Biennial, P = Perennial

3 Use a minimum of 75 GPA - Control of established stands may require repeat applications.

4 For best results early postemergence applications are required.

† Not approved for use in California.

## VINES AND BRAMBLES

Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 1 pint per acre</b>		
Field bindweed	<i>Convolvulus arvensis</i>	P
Hedge bindweed	<i>Calystegia sepium</i>	A
<b>Apply 2 - 3 pints per acre <sup>1</sup></b>		
Wild buckwheat	<i>Polygonum convolvulus</i>	P
<b>Apply 3 - 4 pints per acre <sup>1</sup></b>		
Greenbriar	<i>Smilax</i> spp.	P
Honeysuckle	<i>Lonicera</i> spp.	P
Morningglory	<i>Ipomoea</i> spp.	A/P
Poison ivy	<i>Rhus radicans</i>	P
Redvine	<i>Brunnichia cirrhosa</i>	P
Wild rose	<i>Rosa</i> spp.	P
Including:		P
Multiflora rose	<i>Rosa multiflora</i>	
McCartney rose	<i>Rosa bracteata</i>	P

## VINES AND BRAMBLES

Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 4 - 6 pints per acre <sup>1</sup></b>		
Kudzu <sup>†</sup> , <sup>3</sup>	<i>Pueraria lobata</i>	P
Trumpetcreeper	<i>Campsis radicans</i>	P
Virginia creeper	<i>Parthenocissus quinquefolia</i>	P
Wild grape	<i>Vitis</i> spp.	P

1 The higher rates should be used where heavy or well-established infestations occur.

2 Growth Habit – A = Annual, B = Biennial, P = Perennial

3 Use a minimum of 75 GPA - Control of established stands may require repeat applications.

† Not approved for use in California.

## BRUSH SPECIES

Common Name	Scientific Name	Growth Habit <sup>2</sup>
<b>Apply 4 - 6 pints per acre <sup>1</sup></b>		
American beech	<i>Fagus grandifolia</i>	P
Ash	<i>Fraxinus</i> spp.	P
Bald cypress	<i>Taxodium distichum</i>	P
Bigleaf maple	<i>Acer macrophyllum</i>	P
Black locust <sup>5</sup>	<i>Robinia pseudoacacia</i>	P
Black gum	<i>Nyssa sylvatica</i>	P
Box elder	<i>Acer negundo</i>	P
Cherry	<i>Prunus</i> spp.	P
Chinaberry	<i>Melia azadarach</i>	P
Dogwood	<i>Cornus</i> spp.	P
Elm <sup>6</sup>	<i>Ulmus</i> spp.	P
Hawthorn	<i>Crataegus</i> spp.	P
Hickory	<i>Carya</i> spp.	P
Honeylocust <sup>5</sup>	<i>Gleditsia triacanthos</i>	P
Maple	<i>Acer</i> spp.	P
Mulberry	<i>Morus</i> spp.	P
Oak	<i>Quercus</i> spp.	P
Persimmon	<i>Diospyros virginiana</i>	P
Pine <sup>†</sup> , <sup>5</sup>	<i>Pinus</i> spp.	P
Poplar	<i>Populus</i> spp.	P
Privet	<i>Ligustrum vulgare</i>	P
Red Alder	<i>Alnus rubra</i>	P
Red Maple	<i>Acer rubrum</i>	P
Russian Olive	<i>Eleagnus angustifolia</i>	P
Sassafras	<i>Sassafras albidum</i>	P
Sourwood	<i>Oxydendrum arboreum</i>	P
Sweetgum	<i>Liquidambar styraciflua</i>	P
Water willow <sup>†</sup>	<i>Justica americana</i>	P
Willow	<i>Salix</i> spp.	P
Yellow poplar	<i>Liriodendron tulipifera</i>	P

1 The higher rates should be used where heavy or well-established infestations occur.

2 Growth Habit – A = Annual, B = Biennial, P = Perennial

5 Tank-mix with glyphosate or triclopyr.

6 Tank-mix with glyphosate.

† Not approved for use in California.



## ADJUVANTS

For optimal postemergence performance of AquaPier, the addition of an adjuvant to the spray solution is essential to aid in the deposition and uptake of the herbicide. For this purpose, **ONLY** use spray adjuvants that are approved or appropriate for aquatic use.

**Nonionic Surfactants:** Use a nonionic surfactant at 0.25% v/v or higher (depending on surfactant manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product. Alcohols, fatty acids, horticultural spray oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet these requirements.

**Methylated Seed Oils or Vegetable Oil Concentrates:**

Methylated seed oil or vegetable oil concentrate may be used at 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, mix methylated seed oil or vegetable oil concentrate at a rate of 1% of the total spray volume.

**Silicone Based Surfactants:** Silicone-based surfactants allow greater spreading of the spray droplet on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based surfactants may dry too quickly and limit herbicide uptake. Refer to the surfactant manufacturer's label for specific recommendations.

**Invert emulsions:** AquaPier can be applied as an invert emulsion (water-in-oil emulsion) designed to minimize spray drift and spray run-off, thereby delivering more herbicide to the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Refer to the invert chemical label for proper mixing directions.

**Other:** Anti-foam agents, spray pattern indicators or drift reduction agents may also be used if necessary or desired. Refer to the adjuvant manufacturer's label for specific recommendations.

## AQUATIC USES, PRECAUTIONS AND RESTRICTIONS

AquaPier may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. AquaPier may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water.

- **DO NOT** apply more than 6 pints of product (1.5 pounds acid equivalent) per acre per year.
- Aerial application is restricted to helicopter equipment only.
- AquaPier can only be applied by federal or state agencies, such as Water Management District personnel, municipal officials and the U.S. Army Corps of Engineers, or those applicators who are licensed or certified as aquatic pest control applicators and are authorized by the state or local government.

- Treatment of other-than-non-native, invasive species is limited to those that have been determined to be nuisance vegetation by a federal or state government entity.

- **Applications to private waters:** AquaPier may be applied to private non-flowing waters, such as ponds, lakes and drainage ditches where there is minimal or no outflow to public waters.

- **Applications to public waters:** AquaPier may be applied to public waters such as ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers and other slow-moving or non-moving waters for control of aquatic weeds or of riparian and wetland weed species.

- Consult local state fish and game agency and water control authorities before applying this product to any public waters. Permits may be required to treat such water.

- **Recreational Use of Water in Treated Areas:** There are no restrictions on the use of water in treated areas for recreational purposes, including swimming and fishing.

- **Livestock Use of Water in/from Treated Areas:** There are no restrictions on livestock consumption of water from treated areas.

- **Precautions for Potable Water Intakes: DO NOT** apply AquaPier directly to waters within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within one-half mile of active potable water intakes, the water intake must be turned off during application and for at least 48 hours after the treatment. Such aquatic applications may be made only where there are alternative water sources or holding ponds that can be used while active potable water intakes are turned off during the minimum 48 hours post-treatment period.

**Note:** Existing potable water intakes which are no longer in use, such as those replaced by connections to wells or municipal water systems, are not considered to be active potable water intakes. This restriction does not apply to intermittent, inadvertent overspray of water in terrestrial use sites.

## APPLICATION TO WATERS USED FOR IRRIGATION

**DO NOT** use water treated with AquaPier for irrigation purposes for 120 days after treatment or until AquaPier residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

**Seasonal Irrigation Waters:** AquaPier may be applied during the off-season to surface waters that are used for irrigation on a seasonable basis, provided at least 120 days elapses between AquaPier treatment and the first use of treated water for irrigation or until AquaPier residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

**Irrigation Canals/Ditches:** **DO NOT** apply AquaPier to irrigation canals/ditches unless the 120-day restriction on irrigation water usage can be observed or AquaPier residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less. **DO NOT** apply AquaPier to dry irrigation canals/ditches.

**Quiescent or Slow Moving Waters:** In lakes and reservoirs **DO NOT** apply AquaPier within one (1) mile of an active irrigation water intake during the irrigation season. Applications less than one (1) mile from an inactive irrigation water intake may be made during the off-season, provided the irrigation intake will remain inactive for at least 120 days after treatment or until AquaPier residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

**Moving Waters:** **DO NOT** apply AquaPier within one-half mile downstream of an active irrigation water intake. When making applications upstream from an active irrigation water intake, the intake must be turned off for a period of time sufficient to allow the upstream treated water to completely flow past the irrigation intake. Shut off time will be determined by the speed of water flow and the distance and length of water treated upstream from the intake. Consult local, state and/or federal authorities before making any applications upstream from an active irrigation water intake.

## TANK-MIXES

AquaPier may be tank-mixed with other aquatic-use herbicides to broaden the spectrum of control of emergent and floating aquatic vegetation provided that the tank-mix product label does not prohibit such mixing. Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank-mixes.

## AERIAL APPLICATIONS

Exercise all precautions to minimize or eliminate spray drift. Aerial applications may only be made by helicopter; applications may not be made by airplane. Helicopters can be used to apply AquaPier; however, **DO NOT** apply by helicopter unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or where damage to desirable vegetation can be tolerated. Helicopters equipped with a Microfoil™ boom, Thru-Valve™ boom or raindrop nozzles must be used and calibrated. Unless applying with a Microfoil boom, use a drift control agent at the recommended label rate. To avoid drift, **DO NOT** make applications during inversion conditions, when winds are gusty, or during any other conditions that promote spray drift. Side trimming is not recommended with AquaPier unless death of treated vegetation is acceptable.

Uniformly apply AquaPier in 5 to 30 gallons of water per acre. Use a nonionic surfactant, methylated seed oil or silicone-based surfactant (see the *ADJUVANTS* section of this label for specific recommendations). An anti-foam agent may be added, if needed.

Thoroughly clean application equipment, including landing gear, by thoroughly flushing with water immediately after using this product.

Prolonged exposure of uncoated/unpainted steel (except stainless steel) surfaces to this product may result in corrosion and failure of the exposed part. Maintaining painted surfaces may prevent corrosion.

## GROUND APPLICATIONS

### FOLIAR APPLICATIONS:

• **For Low Volume Foliar Applications:** Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. Thoroughly mix 0.5 to 5% (v/v) AquaPier in water plus surfactant (see the *ADJUVANTS* section of this label for recommendations). Use an anti-foam agent at the recommended rate, if needed. For difficult to control species (see *AQUATIC SPECIES CONTROLLED BY AQUAPIER* section and the *ADDITIONAL WEEDS CONTROLLED BY AQUAPIER* section for relative susceptibility of weed species), apply the higher concentrations of herbicide and/or spray volumes but **DO NOT** apply more than 6 pints of AquaPier per acre. Excessive wetting of foliage is not recommended. See the *SPRAY SOLUTION MIXING GUIDE* below for some suggested volumes of AquaPier and water.

Select appropriate nozzles to avoid over-application. Proper application is critical to ensure desirable results. Optimum results are achieved when the spray covers the crown and approximately 70 percent of the plant. The use of a flat fan nozzle tip with a spray angle of 40 degrees or less will aid in proper deposition.

Recommended nozzle tip sizes include 4004E or 1504E. For a straight stream and cone pattern, use adjustable cone nozzles such as 5500 X3 or 5500 X4. Attaching a rollover valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Moisten, but do not drench target vegetation. **DO NOT** spray to run off.

• **For Low Volume Foliar Applications by Backpacks:** For low-growing species, spray downward to cover approximately 70% of the plant and the crown.

For target species 4 to 8 feet tall, apply a directed spray in a smooth vertical motion from the crown upward on at least two sides of the target vegetation, making sure to cover the crown whenever possible.

For target species over 8 feet tall, apply a directed spray in a smooth zig-zag motion from the crown upward on at least two sides of the target brush.

• **For Low Volume Foliar Applications with Hydraulic Handgun Equipment:** Use same technique as described above for *Low Volume with Backpacks*.

For broadcast applications, simulate a gentle rain near the top of target vegetation, allowing spray to penetrate the target foliage and contact the crown without run-off onto understory vegetation.

**DO NOT** spray to run-off. Herbicide spray that contacts understory vegetation may result in severe injury or death of understory plants.

## Spray Solution Mixing Guide for Low-Volume Foliar Applications

Amount of spray solution	Desired Concentration (fluid volume)				
	0.5%	0.75%	1%	1.5%	5%
	(amount of AquaPier to use)				
1 gallon	0.6 oz	0.9 oz	1.3 oz	1.9 oz	6.5 oz
3 gallons	1.9 oz	2.8 oz	3.8 oz	5.8 oz	1.2 pint
4 gallons	2.5 oz	3.8 oz	5.1 oz	7.7 oz	1.6 pint
5 gallons	3.2 oz	4.8 oz	6.5 oz	9.6 oz	2 pints
50 gallons	2 pints	3 pints	4 pints	6 pints	10 pints
100 gallons	4 pints	6 pints	8 pints	6 quarts	5 gallons

2 tablespoons = 1 fluid ounce

- **For High Volume Foliar Applications:** For optimum performance when spraying medium to high-density vegetation, use equipment calibrated to deliver up to 100 gallons of finished spray per acre (GPA). Application volumes exceeding 100 gallons GPA may result in excessive spray run-off, causing injury to desirable ground cover species. Thoroughly mix AquaPier in water and include a surfactant (see *ADJUVANTS* section for surfactant recommendations). Use an anti-foam agent according to its label, if needed. For hard-to-control species (see *AQUATIC SPECIES CONTROLLED BY AQUAPIER* section and the *ADDITIONAL WEEDS CONTROLLED BY AQUAPIER* section for relative susceptibility of weeds), use the higher concentrations of the herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of AquaPier per acre. Uniformly cover the foliage of the target vegetation but **DO NOT** apply to run-off.
- **Side Trimming:** **DO NOT** side trim with AquaPier unless severe injury or death of the treated vegetation is acceptable. AquaPier is readily translocated and can result in the death of the vegetation.

### CUT SURFACE TREATMENTS

AquaPier will control undesirable woody vegetation when applied as a water solution to the cambium area of freshly-cut stump surfaces or to cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the spring. Tree injection and cut stem treatments are most effective in late summer and early fall. **DO NOT** over apply solution causing run-off from the cut surface.

**NOTE:** Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

#### Cut Surface Applications with Dilute and Concentrate Solutions:

Mix AquaPier as either a concentrated or dilute solution for cut surface treatments. Apply dilute solutions to the cut surface of the stump or to cuts on the stem of the target woody vegetation. Apply concentrate solutions to cuts on the stem. Use of concentrated solutions permits application to fewer cuts on the stem, especially for large diameter trees. Follow the application directions below to determine proper application techniques for each type of solution.

To prepare a dilute solution, mix 8 to 12 fluid ounces of AquaPier with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be added according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve herbicide uptake through partially callused cambium tissues.

To prepare a concentrated solution, mix 2 quarts of AquaPier with no more than 1 quart water.

- **For cut stump treatments:** Spray or brush the solution onto the cambium area of the freshly cut stump surface. Thoroughly wet the entire cambium area (the wood just inside the bark of the stump).
- **For cut stem (injection, hack and squirt) treatments:**  
**Dilute Solutions:** Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Insure that the injector completely penetrates the bark at each injection site.

**Concentrate Solutions:** Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut while a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site, place the injection cuts at approximately equal intervals around the tree.

- **Frill or girdle treatments:** Use a hatchet, machete or chainsaw to make cuts through the bark around the tree to expose the cambium. The cuts should angle downward into the cambium sufficiently deep to expose at least two growth rings. Use a spray applicator or brush to apply a 25% to 100% solution of AquaPier into each cut until thoroughly wet. Avoid applying so much herbicide that runoff to the ground or water occurs.

#### Storage and Disposal

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** **DO NOT** store below 10°F.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL FOR 2.5 GALLON AND 30 GALLON:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL FOR BULK:** When this container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase, or to a designated location. This container must only be refilled with the pesticide product. **DO NOT** reuse the container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of container must be in compliance with state and local regulations.

## Warranty

SePRO Corporation (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. The Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

## Terms of Sale

The Company's directions for use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. All such risks are assumed by the user.

## Limitation of Liability

To the extent permitted by law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent permitted by law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

**The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.**