



Active Ingredient: Garlic..... 20.00%
Other Ingredients*: 80.00%
 (*Water, Citric Acid, Potassium Sorbate)

All ingredients are exempt from tolerance requirements under 40 CFR 180.1001, and 180.1164. This product is FIFRA 25 (b) exempt.

USE RESTRICTIONS AND CAUTIONS

- Not warranted to kill insects. This product is an Insect Repellent.
- Do not apply **BioLink Insect Repellent** within 12 hours of crop harvest.
- Do not apply **BioLink Insect Repellent** if rain is forecasted or overhead watering occurs within 12 hours.
- **POLLINATION WARNING: MAY REPEL BEES.** BioLink Insect Repellent may repel bees. Do not apply less than three days prior to pollination.
- **CAUTION: KEEP OUT OF REACH OF CHILDREN.** Avoid contact with eyes. Wash with soap and water after handling.

Notice of Warranty: Westbridge warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. In no case shall Westbridge be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY
 **Westbridge**
 Agricultural Products

1260 Avenida Chelsea • Vista, CA 92081
 (800) 876-2767 • www.westbridge.com

NET CONTENTS

1 US Quart (0.9 L) / 2.1 LBS (1.0 KG)
 1 US Gallon (3.8 L) / 8.4 LBS (3.8 KG)
 2.5 US Gallons (9.5 L) / 21.0 LBS (9.5 KG)



For use on all crops including fruits and vegetables, fruit trees, vines, tree nuts, field and row crops, ornamental trees, shrubs and flowers, containerized plants, and turf

To repel aphids, beetles, borers, caterpillars, earwigs, fleas, flies, grubs, leafhoppers, leafrollers, lygus bugs, mealy bugs, mites, moths, scales, slugs, thrips, weevils, whiteflies, worms and other plant feeding insects.

Also for repelling deer, birds, field mice, rabbits, and other animal pests.

APPLICATION RATES

Foliar Application: ½ to 2 quarts per acre; use higher rates for heavier pest infestations. Do not spray to point of run-off.

Nursery & Container: ½ to 1 quart per 100 gallons of buffered water.

Home & Garden: 2 to 4 ounces per gallon of water; Spray plant foliage thoroughly including underside of leaves.

MIXING INSTRUCTIONS

General mixing instructions: Shake container well or stir contents thoroughly before adding **BioLink Insect Repellent** to the spray tank.

Buffering instructions for agricultural and large scale uses: spray tank pH must be adjusted to 5.5 to 6.0 before adding **BioLink Insect Repellent** to the spray tank. Recommended buffering agents are BioLink Buffer and Penetrant, citric acid, acetic acid, phosphoric acid or aliphatic polycarboxylate-type materials. Do not buffer with propionic acid or similar products that affect biologically active products. After mixing, agitate spray tank-mix before and during application to prevent settling.

Tank mixing with other products: when mixing BioLink Insect Repellent with other products, it is strongly recommended to test a small plot before applying the spray mix.

If tank-mixing **BioLink Insect Repellent** with other pesticides, use the recommended label rates of the other pesticides until exact tank-mixed product performance is determined.

APPLICATION TIMING AND PEST PRESSURE

When using **BioLink Insect Repellent**, the optimum pest pressure to start application is 0-5%. Repeat application timing should be 8-14 days, and determined by field scouting reports of pest pressure, plant nutrition and temperature.

SPRAY TEMPERATURE

The optimum temperature range to apply BioLink Insect Repellent is 70°-90°F (21°-32°C). Minimum recommended application temperature is 50°F (10°C). Maximum recommended application temperature is 100°F (38°C). For applications above 100°F higher application spray rates (more than 40 gals/acre) are recommended after there has been a minimum of a 15°F (9°C) drop in air temperature.

SPRAY ADJUVANTS

To maximize the effectiveness of BioLink Insect Repellent add 4 oz of BioLink Spreader-Sticker per 100 gallons water per acre. **BioLink Insect Repellent** should not be used with dormant spray oils, heavy oils, spreader-stickers or surfactants that will coat the surface of the plant foliage and prevent absorption of the product. Silicone-based or non-ionic surfactants are strongly recommended to increase product coverage and absorption. If **BioLink Insect Repellent** is not rapidly absorbed by the foliage, it will not be effective.