

WEED BEATER PLUS CRABGRASS & BROADLEAF WEED KILLER READY TO SPRAY

THIS PRODUCT CONTAINS:

0.456 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
0.128 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
0.043 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
Isomer Specific by AOAC Method

- **KILLS OVER 200 LAWN WEEDS PLUS CRABGRASS IN ONE EASY STEP**
- **ONE QUART COVERS UP TO 6,600 SQUARE FEET.**

CONTROLS: CRABGRASS, DANDELION, BLACK MEDIC, CHICKWEED, CLOVER, CREEPING CHARLIE, GROUND IVY, LAMBSQUARTERS, MORNINGGLORY, OXALIS, POISON IVY, POISON OAK, SPEEDWELL, SPURGE, WILD CARROT, YARROW AND OTHER BROADLEAF WEEDS.

EPA Est. No. 4-NY-1
EPA Reg. No. 4-466

Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING.

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grassy weeds including crabgrass, dandelion, foxtail, chickweed and clover. (see complete listing inside)

Where to Use: On Lawns: Kills the weeds, not the lawn when used as directed. Can be used on Kentucky bluegrass, Ryegrass (perennial and annual), Tall Fescue, Mixtures of cool season grasses, Bermudagrass* (common and hybrid), Zoysiagrass and Buffalograss.

When to Use: Apply when daytime temperatures are below 90°F.

Coverage: This container treats 6,600 sq. ft.

How it Works: This product contains three proven weed killers that target lawn weeds and crabgrass. It enters the lawn weeds through their leaves and moves throughout the plant to kill the weed, roots and all. Approved lawns listed on this label will not be harmed when used as directed.

*Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

HOW TO USE

- Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This 32 fl. oz. container will cover 6,600 square feet.
- Make sure sprayer nozzle is in "OFF" position then, connect sprayer to garden hose.
- Turn water on at faucet. Extend hose to the farthest area of lawn to be treated. To begin spraying, push lock tab down and turn knob control lever to "ON" position. The product mixes automatically with the water. Walk at a steady pace while spraying evenly and working back toward the faucet to avoid walking over wet, treated grass.
- To stop spraying, turn knob control lever to the "OFF" position. Turn off water at the faucet. To relieve pressure, turn knob control lever to the "ON" position and point the sprayer away from any desirable plants before disconnecting the spray nozzle from the garden hose.

Apply this product when weeds are small and actively growing.

Can be applied to the following turf grasses: Kentucky bluegrass, Ryegrass (perennial and annual), Tall Fescue, mixtures of cool season grasses, Bermudagrass* (common and hybrid), Zoysiagrass and Buffalograss.

*Do not apply to Bermudagrass lawns when daytime temperatures exceed 85°F. Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected.

The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

PRODUCT TIPS AND NOTES

When to apply:

- When weeds are small and actively growing.
- When temperature is below 90°F.
- For Newly Seeded Lawns, do not apply until after the third mowing (approximately 6 weeks after seeding emergence)
- For Newly Sodded, Sprigged or Plugged Lawns, apply 3 to 4 weeks after planting.

Watering:

- Adequate soil moisture and favorable growing conditions enhance the performance of this product.
- Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule watering, before and after the application. For best results, delay the second watering until approximately 24 hours after the application.

Mowing:

- Delay mowing 2 days before and until 2 days after the application.
- Do not use clippings from the treated areas as mulch or compost, around flowers, ornamentals, trees, or in vegetable gardens.

Non-Target Plants:

- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers or other nontarget plants.

GRASS AND BROADLEAF WEEDS CONTROLLED

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant		Signalgrass, broadleaf
BROADLEAF WEEDS			
Annual yellow sweetclover	Buckhorn	Clover, white	Ground ivy
Aster	Buckhorn plantain	Cockle	Gumweed
Austrian fieldcress	Bulbous buttercup	Cocklebur	Hairy fleabane
Bedstraw	Bull thistle	Common mullein	Hawkweed
Beggarticks	Bullnettle	Creeping Jenny	Healall
Betony, Florida	Burclover	(Creeping Charlie)	Heartleaf drymary
Bindweed, field	Burdock	Cudweed	Heathaster
Bird vetch	Burning nettle	Curly dock	Hedge bindweed
Bitter wintercress	Burweed	Daisy, English	Hedge mustard
Bittercress, hairy	Buttercup	Daisy fleabane	Hemp
Bitterweed	Buttonweed	Daisy, oxeye	Henbit
Black-eyed Susan	Canada thistle	Dandelion	Hoary cress
Black medic	Carolina geranium	Dichondra	Hoary plantain
Black mustard	Carpetweed	Dogbane	Hoary vervain
Blackseed plantain	Catchweed bedstraw	Dogfennel	Horsenettle
Blessed thistle	Catnip	Dollarweed	Jimsonweed
Bloodflower milkweed	Catsear	Elderberry	Knawel
Blue lettuce	Chickweed, common	False dandelion	Knotweed
Blue vervain	Chickweed, mouseear	Falseflax	Kochia
Bracted plantain	Chicory	False sunflower	Lambsquarters
Brassbuttons	Cinquefoil	Fiddleneck	Lespedeza
Bristly oxtongue	Clover, crimson	Florida pusley	Mallow
Broadleaf dock	Clover, hop	Frenchweed	Matchweed
Broadleaf plantain	Clover, red	Galinsoga	Mexicanweed
Broomweed	Clover, strawberry	Goathead	Milk vetch
	Clover, sweet	Goldenrod	Morningglory

BROADLEAF WEEDS Cont.

Mouseear hawkweed	Prostrate spurge	Stinging nettle	Wild onion
Mugwort	Prostrate vervain	Strawberry, India	Wild parsnip
Musk thistle	Puncturevine	mock	Wild radish
Mustard	Purslane, common	Tall nettle	Wild rape
Narrowleaf plantain	Ragweed	Tall vervain	Wild strawberry
Narrowleaf vetch	Red sorrel	Tansy ragwort	Wild sweet
Nettle	Redroot pigweed	Tansy mustard	potato
Orange hawkweed	Redstem filaree	Tanweed	Wild vetch
Oriental cocklebur	Rough cinquefoil	Thistle	Wild violet
Oxalis	Rough fleabane	Trailing crownvetch	Woodsorrel
Parsley-piert	Russian pigweed	Tumble mustard	Woolly croton
Parsnip	Russian thistle	Tumble pigweed	Woolly
Pearlwort	Scarlet pimpernel	Velvetleaf	morningglory
Pennycress	Scotch thistle	Venice mallow	Woolly plantain
Pennywort	Sheep sorrel	Virginia buttonweed	Wormseed
Peppergrass	Shepherdspurse	Virginia creeper	Yarrow
Pepperweed	Slender plantain	Virginia pepperweed	Yellow rocket
Pigweed	Smallflower	Water pennywort	Yellowflower
Pineywoods bedstraw	galinsoga	Wavyleaf bullthistle	pepperweed
Plains coreopsis	Smooth dock	Western Clematis	and other
(tickseed)	Smooth pigweed	Western salsify	broadleaf
Plantain	Sorrel	White mustard	weeds
Poison ivy	Sowthistle	Wild aster	
Poison oak	Spanishneedles	Wild buckwheat	
Pokeweed	Speedwell	Wild carrot	
Poorjoe	Spiny amaranth	Wild four-o'clock	
Prairie sunflower	Spiny cocklebur	Wild garlic	
Prickly lettuce	Spiny sowthistle	Wild geranium	
Prickly sida	Spotted catsear	Wild lettuce	
Prostrate knotweed	Spotted spurge	Wild marigold	
Prostrate pigweed	Spurweed	Wild mustard	

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Avoid contamination of food or feedstuffs.

PESTICIDE DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

If in eyes:	<ul style="list-style-type: none">•Hold eye open and rinse slowly and gently with water for 15-20 minutes.•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.•Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">•Call a poison control center or doctor immediately for treatment advice.•Have person sip a glass of water if able to swallow.•Do not induce vomiting unless told to by a poison control center or doctor.•Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1800-858-7378 for emergency medical treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

LIMITED WARRANTY AND DISCLAIMER

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, seller's liability for default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, seller shall have no liability for consequential damages.