



CYONARA™ PLUS

Pour-On Topical Insecticide



**Treats
94
Cattle
(<600 lbs.)**

- **Contains Lambda-Cyhalothrin and PBO**
- **For the control of Lice and Horn Flies on beef cattle, cows and calves**

ACTIVE INGREDIENT:

Lambda-Cyhalothrin	1.0%
Piperonyl Butoxide	5.0%
OTHER INGREDIENTS:	94.0%
TOTAL:	100.0%

EPA Reg. No.: 53883-248 EPA Est. No.: 53883-TX-002

Keep Out of Reach of Children

WARNING

(See back panel for additional Precautionary Statements)

Control Solutions, Inc. • 5903 Genoa-Red Bluff • Pasadena, TX 77507-1041

NET CONTENTS: 1 QUART

FIRST AID

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING - May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove contaminated clothing and wash before re-use. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Environmental Hazards: This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by disposal of rinse water. Apply this product only as specified on the label.

Physical or Chemical Hazards: Do not store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Ready to Use - dilution is not necessary. This product may be used with automatic dosing systems.

Restrictions: Do not apply to lactating or dry dairy cows, or to the face of beef cattle, cows or calves.

Cyonara™ Plus is not recommended for use on veal calves.

Apply to	Target Insects	Dosage
Beef Cattle, Cows and Calves	Lice*, Horn Flies	Apply product down the backline at the rate of 10 mL (1/3 fl oz) per head for cattle weighing less than 600 lb. or at a rate of 15 mL (1/2 fl oz) per head for cattle weighing more than 600 lb.

Repeat as necessary but, do not apply more than once every two weeks and do not apply more often than four times within any six month period.

* For reduction of sucking lice, two treatments at a 14-day interval are recommended.

Note: Cyonara™ Plus is not effective in controlling cattle gnubs. This product may be applied to cattle at any time during the year and not have host-parasite reactions, associated with grub treatment products, occur. Use **Cyonara™ Plus** in an IPM (integrated pest management) system which may involve repeat treatments and the use of other pest control practices. Continued exposure of horn flies to a single class of insecticide (e.g. pyrethroids or organophosphates) may lead to the development of resistance to that class of insecticide. In order to reduce the possibility of horn flies developing resistance, it is important to rotate, on a seasonal basis, the class of insecticide used and/or the method of horn fly control. For advice concerning current control practices with relation to specific local conditions, consult resources in resistance management programs and your Cooperative Agricultural Extension Service.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container and identify contents.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.





CYONARA™ PLUS

Pour-On Topical Insecticide



**Treats
378
Cattle
(<600 lbs.)**

- **Contains Lambda-Cyhalothrin and PBO**
- **For the control of Lice and Horn Flies on beef cattle, cows and calves**

ACTIVE INGREDIENT:	
Lambda-Cyhalothrin	1.0%
Piperonyl Butoxide	5.0%
OTHER INGREDIENTS:	94.0%
TOTAL:	100.0%

EPA Reg. No.: 53883-248

EPA Est. No.: 53883-TX-002

Keep Out of Reach of Children

WARNING

(See back panel for additional Precautionary Statements)

Control Solutions, Inc.

5903 Genoa-Red Bluff • Pasadena, TX 77507-1041

NET CONTENTS: 1 GALLON

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING - May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove contaminated clothing and wash before re-use. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Environmental Hazards: This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by disposal of rinse water. Apply this product only as specified on the label.

Physical or Chemical Hazards: Do not store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Ready to Use – dilution is not necessary. This product may be used with automatic dosing systems.

Restrictions: Do not apply to lactating or dry dairy cows, or to the face of beef cattle, cows or calves.

Cyonara™ Plus is not recommended for use on veal calves.

Apply to	Target Insects	Dosage
Beef Cattle, Cows and Calves	Lice*, Horn Flies	Apply product down the backline at the rate of 10 mL (1/3 fl oz) per head for cattle weighing less than 600 lb. or at a rate of 15 mL (1/2 fl oz) per head for cattle weighing more than 600 lb.
Repeat as necessary but, do not apply more than once every two weeks and do not apply more often than four times within any six month period.		
* For reduction of sucking lice, two treatments at a 14-day interval are recommended.		

Note: **Cyonara™ Plus** is not effective in controlling cattle grubs. This product may be applied to cattle at any time during the year and not have host-parasite reactions, associated with grub treatment products, occur. Use **Cyonara™ Plus** in an IPM (integrated pest management) system which may involve repeat treatments and the use of other pest control practices. Continued exposure of horn flies to a single class of insecticide (e.g. pyrethroids or organophosphates) may lead to the development of resistance to that class of insecticide. In order to reduce the possibility of horn flies developing resistance, it is important to rotate, on a seasonal basis, the class of insecticide used and/or the method of horn fly control. For advice concerning current control practices with relation to specific local conditions, consult resources in resistance management programs and your Cooperative Agricultural Extension Service.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container and identify contents.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.

