



Folo Spray 20-10-30

GUARANTEED ANALYSIS:

TOTAL NITROGEN (N)	20.00%
6.90% Nitrate Nitrogen	
13.10% Urea Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)	10.00%
SOLUBLE POTASH (K ₂ O)	30.00%
Boron (B)	0.02%
0.02% Water Soluble Boron	
Copper (Cu)	0.05%
0.05% Chelated Copper	
Iron (Fe)	0.10%
0.10% Chelated Iron	
Manganese (Mn)	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo)	0.0005%
0.0005% Water Soluble Molybdenum	
Zinc (Zn)	0.05%
0.05% Chelated Zinc	

Derived from Monopotassium Phosphate, Potassium Nitrate, Urea, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate, and Zinc EDTA.

**KEEP OUT OF REACH OF CHILDREN****WARNING**

CAUSES EYE IRRITATION. MAY INTENSIFY FIRE; OXIDIZER.

PRECAUTIONARY STATEMENTS: Wash hands thoroughly after handling. Keep away from heat. Keep/Store away from clothing and other combustible materials. Take any precaution to avoid mixing with combustible materials.

PERSONAL PROTECTIVE EQUIPMENT: Wear protective gloves, eye protection, face protection.

FIRST AID: IF IN EYES Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical attention.

IN CASE OF FIRE: Use chemical type foam, carbon dioxide, dry chemical, water fog or spray to extinguish.

DISPOSAL: Dispose of contents / container in accordance with Federal, state, and local regulations.

GENERAL USE DIRECTIONS

COMPATABILITY: Folo Spray 20-10-30 is compatible with most commonly used pesticides. However, a compatibility test is recommended before full scale mixing if the desired combination has not been used in the past. Do not combine with dormant oil, lime sulfur, copper fungicides, or other highly alkaline materials. Do not use with the amine form of 2,4-D or with Dicamba (e.g. Banvel).

DILUTION RATES: Unless otherwise specified, use the rate per acre and dilute as follows:

Air: 5-30 gallons of water per acre.

Dilute: 100 gallons of water per acre.

Concentrate: 10-100 gallons of water per acre.

Do not exceed 2 lbs. of product per gallon of water. Do not exceed recommended rates per acre.

MIXING INSTRUCTIONS

Fill tank half full of water and begin agitation. Add Folo Spray 20-10-30 and continue agitation. Do not premix as a slurry. Add pesticide last. Water temperature plays an important role in determining fertilizer solubility. The higher the water temperature the greater the solubility and quicker the dissolving time. As fertilizers are added to the solution, there is an immediate decrease in water temperature. It is important when preparing concentrated solutions to allow for this temperature drop to ensure that all the fertilizer is completely dissolved. This product will lower the pH of the spray solution.

APPLICATION IN IRRIGATION WATER

Solid Set, Overhead, Sprinkler or Pivot Irrigation Systems: Dissolve Folo Spray 20-10-30 in mix tank and apply 8-10 lbs. per acre in irrigation water. Use of an anti-siphon or back-check valve is recommended to prevent material from running back into primary water supply.

Do not use when temperatures are expected to exceed 90°F or when plants are under water stress conditions such as drought.

DECIDUOUS TREES (SUCH AS APPLES, PEARS, CHERRIES, PLUMS, PRUNES, PEACHES, NECTARINES, ALMONDS, WALNUTS, FILBERTS, PISTACHIOS AND PECANS): Apply 10-15 lbs. per acre in delayed dormant, pre-bloom and post-bloom sprays. Avoid combining with materials that may cause russetting. A post-harvest application of 10-40 lbs. per acre can enhance tree vigor and increase nutrient levels within spurs the following season.

CITRUS: Apply 8-10 lbs. per acre at 7-10 day intervals. Use the higher rate for full bearing trees.

VINE CROPS (SUCH AS MELONS, CUCUMBERS): Apply 8-10 lbs. per acre when plants are 12"-14" across and again at the first sign of bloom. Make a third application 2-3 weeks after bloom. Repeat at 2-3 week intervals as needed. For Air Application apply rate in no less than 10 gallons of water per acre.

HOPS: Apply 8-10 lbs. per acre when hops are halfway up the wire and again when they reach the top of the wire. For Air Application apply rate in no less than 10 gallons of water per acre.

GRAPES: Apply 5-10 lbs. per acre when leaves emerge, pre-bloom and post-bloom until grapes are ¼ size. For Air Application apply rate in no less than 10 gallons of water per acre.

POTATOES: Apply 10 lbs. per acre. Make the first application when plants are 4" to 6" high (hooking stage) and repeat at 2-3 week intervals.

CORN: (FIELD AND SWEET): Apply 8-10 lbs. per acre. Make the first application when corn is 6" to 8" high and again prior to tasseling.

BEANS, PEAS, LENTILS, RAPESEED (CANOLA): Apply 5-10 lbs. per acre at first sign of bloom. Repeat application 2 weeks later.

COLE CROPS, LETTUCE, LEAFY VEGETABLES: Apply 5-10 lbs. per acre after thinning or transplanting. Repeat at 10-14 day intervals as needed. Addition of a wetting agent is recommended.

SUGAR BEETS: Apply 8-10 lbs. per acre when leaves are 10" across and again 3-4 weeks later. Addition of a wetting agent is recommended.

FIELD CROPS (MINT, ALFALFA, CLOVER): Apply 8-10 lbs. per acre when plants have 6" to 8" of new growth. Repeat at pre-bloom and again 3 weeks later.

BERRIES (SUCH AS STRAWBERRIES, RASPBERRIES, BLACKBERRIES, BLUEBERRIES): Apply 5-10 lbs. per acre on new spring growth. Repeat at pre-bloom and again 3 weeks later.

CRANBERRIES: Apply 8-10 lbs. per acre at popcorn stage. Repeat at hooking and again 3 weeks later.

NOTICE: Before using this product, read the Directions for Use and Disclaimer of Warranty statements. If the Disclaimer of Warranty is not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

Warning: This product contains a chemical known to the State of California to cause cancer, birth defects or other reproductive harm. Proposition 65, the Safe Drinking Water and Toxic Enforcement Act of 1986, requires notification of potential exposure to substances identified by the State of California as causing cancer, birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-jr.com>

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

GHS_R2-15-15

NET WEIGHT: 25 POUNDS (11.34 Kg.)


Simplot

J. R. SIMPLOT COMPANY

P.O. Box 70013, Boise ID 83707-0113 • (208) 336-2110
GUARANTOR • P.O. Box 198, Lathrop, CA 95330-0198