



**For use in the SmartBox® system only, a closed handling and application system.**

**For control of various insects infesting certain field and vegetable crops.**

**Active Ingredient:**

Chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate..... 15.0%

**Inert Ingredients:**.....85.0%

**Total:**.....100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION / PRECAUCION**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.**

**(If you do not understand this label, find someone to explain it to you in detail).**

FIRST AID Organophosphate	
If in eyes:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed:	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
EMERGENCY INFORMATION	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. <b>FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:</b> Transportation: CHEMTREC..... 1-800-424-9300 Other: AMVAC..... 1-323-264-3910	
NOTE TO PHYSICIAN	
Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are barrier laminate or viton. For more information, follow instructions in Supplement Three of PR Notice 93-7. If you want more options, follow the instructions for category H on an EPA chemical resistance category selections chart.

**Loaders, applicators and all other handlers must wear:**

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant footwear plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS**

**IMPORTANT:** The SmartBox system, when used correctly, qualifies as a closed loading system under the WPS. Loaders using the SmartBox system must:

- Wear the PPE specified above for loaders
- In addition to wearing the required PPE, have immediately available for use in case of an emergency –
  - Chemical-resistant apron
  - Chemical-resistant footwear, and
  - NIOSH-approved respirator with an organic-vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

When applicators use closed cab equipment in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**Harmful If Swallowed • Causes Moderate Eye Irritation**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.**

**Do not attempt to open or tamper with the container. Do not attempt to repair the SmartBox container.** If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

This product is available only in the SmartBox, a closed handling and application system.

EPA REG. NO. 5481-525

EPA EST. NO. \_\_\_\_\_

NET CONTENTS: As Marked on Container

**AMVAC**  
 4100 E. Washington Blvd.  
 Los Angeles, CA 90023, USA  
 1-323-264-3910 • www.amvac-chemical.com

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Certified crop advisors or persons entering under their direct supervision under certain circumstances may be exempt from the early reentry requirement pursuant to 40 CFR Part 170.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made out of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### Specific Use Restrictions:

- Do not cut or graze treated alfalfa within 21 days after application.
- Maximum single application rate is 1 lb ai chlorpyrifos per acre.
- Do not make more than 1 application of Lorsban 15G per year.

### ASPARAGUS

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Postharvest Treatment:** Apply Lorsban 15G as a postharvest ground application for suppression of the target pest. Apply as a band over the entire crown area when the asparagus beds have been split (i.e., remove most of the soil from above the asparagus crowns). Cover the area with soil the day of application. **Note:** Control may be reduced in soils with high organic matter content.

Target Pests	Lorsban 15G (lb/acre)
Symphylans (centipede)	10

### Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 180 days of harvest.
- Do not apply more than a total of 3 lb active ingredient of chlorpyrifos per acre between harvests.
- For use only in the Midwest and Pacific Northwest states.

### CITRUS ORCHARD FLOORS

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Broadcast Treatment:** To control foraging ants and suppress mounds, apply Lorsban 15G in combination with a rotary granular applicator, such as a fertilizer spreader, that will uniformly broadcast the granules over the orchard floor. Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Foliar applications of Lorsban 4E may be made in addition to orchard floor treatments of Lorsban 15G (see Specific Use Restrictions).

Target Pests	Lorsban 15G (lb/acre)
Imported fire ant other ant species	6.7

### Specific Use Restrictions:

- **Preharvest Interval:** Do not apply last treatment within 28 days before harvest.
- Do not make more than 3 applications or more than 20.1 lb of Lorsban 15G (3.0 lb ai chlorpyrifos per acre) per year to the orchard floor.
- Do not make a second application within 10 days of any application of chlorpyrifos to the orchard.
- Do not apply more than 1 lb ai chlorpyrifos in a single application.
- Do not allow meat or dairy animals to graze in treated areas.

### CORN (FIELD CORN AND SWEET CORN, INCLUDING CORN GROWN FOR SEED)

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Lorsban 15G is compatible with ALS inhibiting herbicides when applied in accordance with label recommendations. Refer to the herbicide product labels for additional precautions and directions for use.

**At Plant T-Band Treatment:** Apply as a 4 to 5-inch T-band over the open seed furrow behind the planter shoe, and in front of the press wheel. In conventional and minimum-till corn, incorporate granules into the top 1/2 to 1 inch of soil using suitable equipment. Equivalent rates of Lorsban 15G per acre for various row spacings are given in Table 1.

**At Plant In-Furrow Treatment:** Apply Lorsban 15G at planting as an in-furrow treatment in conventional, minimum and no-till corn. Direct the granules into the planter shoe with the seed or place the applicator tube directly behind the planter shoe so that the granules drop into the open seed furrow and are completely covered with soil.

**Postplant Soil Treatment:** To control corn rootworm larvae, apply Lorsban 15G at cultivation by directing the granules at the base of the plant on both sides of the row just ahead of the cultivation shovels and cover the granules with soil. To control European and Southwestern corn borer larvae, apply Lorsban 15G in a band over the row so that the granules are directed into the whorl. Consult your state extension service specialist or agricultural consultant for application timing and local threshold information. Scouting for insect damage is strongly encouraged prior to making a treatment decision.

**Postplant Foliar Treatment:** Use Lorsban 15G for control of European and southwestern corn borers by directing the granules into the corn whorls using high-clearance ground application equipment.

## GENERAL INFORMATION

### SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

### APPLICATION RATES

#### Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings

Ounces of Lorsban 15G Per 1000 Feet of Row	Pounds of Lorsban 15G Required Per Acre from Various Row Spacing							
	40"	38"	36"	34"	32"	30"	22"	18"
3.7	3.0	3.2	3.4	3.6	3.8	4.0	5.5	6.7
4	3.3	3.4	3.6	3.8	4.1	4.4	5.9	7.3
4.5	3.7	3.9	4.1	4.3	4.6	4.9	6.7	8.2
6	4.9	5.2	5.4	5.8	6.1	6.5	8.9	10.9
6.5	5.3	5.6	5.9	6.2	6.6	7.1	9.7	11.8
7.5	6.1	6.4	6.8	7.2	7.7	8.2	11.1	13.6
8	6.5	6.9	7.3	7.7	8.2	8.7	11.9	14.5
9	7.4	7.7	8.2	8.6	9.2	9.8	13.4	16.3
12	9.8	10.3	10.9	11.5	12.3	13.1	17.8	21.8
15	12.3	12.9	13.9	14.4	15.3	16.3	22.3	27.2
16	13.1	13.8	14.5	15.4	16.3	17.4	23.8	29.0

## APPROVED USES

### ALFALFA

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant In-Furrow Treatment:** Apply Lorsban 15G at planting as an in-furrow treatment for suppression of the target pests during establishment. Direct the granules into the planter shoe with the seed or place the applicator tube directly behind the planter shoe so that the granules drop into the seed furrow or place the granular band applicator behind the planter shoe so that the granules fall on the soil surface and into the open seed furrow and are completely covered with soil.

Target Pests	Lorsban 15G (lb/acre)
Cutworms Grubs Wireworms	6.7

Apply at a rate of 5.0 lb per acre for low to moderate first generation infestations or at 6.5 lb per acre for severe first generation infestations and all second-generation infestations. Apply before larvae have entered corn stalks. Consult your state extension service specialist or agricultural consultant for application timing and local threshold information. Scouting for insect damage is strongly encouraged prior to making a treatment decision.

Pests Controlled	Banded/In furrow Applications (oz/1000 feet of row)		
	At Plant Treatments		Postplant Treatment
	T-Band	In-Furrow	
Northern and Western corn rootworm larvae	8	8	8
Southern corn rootworm larvae	8	8	8
Cutworms (1)	8	8	-
Wireworms (2)	8	8	-
Billbugs	8	-	-
Seed corn maggots	8	8	-
Grubs	8	8	-
Lesser cornstalk borer	8	-	-
Seed corn beetle	8	8	-
Symphylans	8	-	-
Armyworms	-	-	6 - 8
European and Southwestern corn borer (3)			
1st Generation	-	-	3.5 - 8
2nd Generation	-	-	6 - 8
Fire ants (1)	8	8	-
Chinch bug (1)	8	8	-

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**Pest-Specific Use Directions:**

- Cutworms, fire ants and chinch bugs:** The 8 oz rate per 1000 ft of row provides suppression only for in-furrow treatments.
- Wireworms:** For best control, apply as an in-furrow treatment. Consider using an insecticidal seed treatment with T-band applications.
- European corn borer:** When using postplant banded applications use rates of 3.5 to 4 oz of Lorsban 15G per 1000 feet of row for low to moderate first generation infestations before larvae have entered corn stalks. Use application rates of 6 to 8 oz of Lorsban 15G per 1000 feet of row for severe first generation infestations and all second-generation infestations before larvae have entered corn stalks.

**Specific Use Restrictions:**

- Preharvest Interval:** Do not apply within 35 days before harvest of grain.
- For foliar insect control, do not exceed the equivalent of 16 oz of Lorsban 15G per 1000 feet of row or 13 lb of Lorsban 15G per acre per crop season.
- Do not make more than 1 at-plant application and 1 foliar application of Lorsban 15G per season or make more than 3 applications per season of products containing chlorpyrifos. Re-treatment with a second soil application of Lorsban 15G is allowed under replant situations due to loss of crop during establishment.
- Do not apply more than a total of 3 lb active ingredient of chlorpyrifos per acre per season.
- Do not make a second application within 10 days of any application of chlorpyrifos to corn.
- Do not allow meat or dairy animals to graze in treated areas or harvest treated corn silage as feed for meat or dairy animals within 14 days after last treatment.
- Do not feed treated corn fodder to meat or dairy animals within 35 days after last treatment.

**GRAIN SORGHUM/MILO**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant T-Band Treatments:** Apply as a T-band in over the open seed furrow in front of the press wheel and incorporate into the top 1/2 to 1 inch of soil using suitable equipment. Equivalent rates of Lorsban 15G per acre for various row spacings are given in Table 1. Use the lowest rate for lesser cornstalk borer control when protection is desired for 2 to 3 weeks or a higher rate for longer residual activity. It is absolutely necessary to incorporate the granules into the soil.

Pests Controlled	At Plant T-Band Applications (oz/1000 feet of row)
Lesser cornstalk borer	4 - 8
Chinch bugs (1)	8
Corn rootworm	
Cutworms	
Fire ants (1)	

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**Pest-Specific Use Directions:**

- Chinch bugs and fire ants:** Suppression only.

**Specific Use Restrictions:**

- Do not make more than 1 application of Lorsban 15G per season.
- The maximum single application rate is 8 oz per 1000 feet of row (1.5 lb active chlorpyrifos per acre).
- Do not make a foliar application of Lorsban 4E or other product containing chlorpyrifos within 10 days of an at-plant application of Lorsban 15G.

**ONION (DRY BULB)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant In-Furrow Treatment:** Apply as an at plant in-furrow treatment so that granules are covered with soil.

Target Pest	Lorsban 15G (oz/1000 ft of row)
Onion maggot (1)	3.7

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**Pest-Specific Use Directions:**

- In Colorado, Idaho, Washington and Oregon,** to control **onion maggots** in onions planted in double rows with rows spaced 2 to 4 inches apart, apply Lorsban 15G at the rate of 3.7 oz per 1000 feet of double row. Place the granules in a 5 to 7 inch wide band over both rows behind the planter shoe and in front of the press wheel to achieve shallow incorporation.

**Specific Use Restrictions:**

- Do not apply more than 1.0 lb active chlorpyrifos per crop season.
- The maximum single application rate is 0.56 oz active chlorpyrifos per 1000 feet of row.
- Do not make more than 1 application per year of Lorsban 15G or other product containing chlorpyrifos.

**PEANUT**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant Preventive Treatment:** Apply Lorsban 15G in a 6 to 12 inch band over the row behind the planter shoe and in front of the press wheel. Incorporate granules to a depth of 1 inch into the soil with tines drag chains, or other suitable equipment. If the 7.5 oz rate per 1000 feet of row is used at planting time, then another application of 7.5 oz per 1000 feet of row may be needed postplant to extend control.

**Postplant Preventive Treatment:** Apply Lorsban 15G to peanuts at early flowering to pegging stage of growth in a 6 to 8 inch band over the row. For extended insect control and continued suppression of white mold (southern blight), a second application of Lorsban 15G may be made (see Specific Use Restrictions).

**Band Rescue Treatment:** Apply Lorsban 15G in a 10 to 18 inch band over the fruiting zone.

**Postplant Rescue Treatment:** Apply Lorsban 15G in a 10 to 18 inch band over the fruiting zone to control Lesser cornstalk borer.

Pests Controlled	Banded Applications (oz/1000 feet of row)	
	At Plant Treatment	Postplant Treatment
<b>Preventive treatments:</b>	7.5 - 15	7.5 - 15
Cutworms		
Lesser cornstalk borer		
Southern corn rootworm larvae		
Wireworms		
White mold (Southern blight) (1)		
Potato leafhopper	-	15
<b>Rescue treatments:</b>	-	7.5 - 15
Lesser cornstalk borer (2)		

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**Pest-Specific Use Directions:**

- Suppression of white mold:** Best suppression of white mold (southern blight) is obtained by applying the maximum rate of 15 oz per 1000 ft of row. Irrigation or rain following application is needed to enhance treatment effectiveness for suppression of white mold. Under conditions of heavy white mold pressure, a suitable fungicide may also be required and must be applied separately.
- Lesser cornstalk borer:** Use Lorsban 15G for the control of lesser cornstalk borer as a rescue treatment when the insect first appears, usually just prior to or at pegging.



**Specific Use Restrictions:**

- **Preharvest Interval:** Do not apply within 21 days before harvest of peanut.
- Do not apply more than 15 oz of Lorsban 15G per 1000 ft of row per crop season or apply more than 4 lb active chlorpyrifos per season.
- Do not make more than 2 applications per year.
- Do not make a second application of Lorsban 15G or other product containing chlorpyrifos within 10 days of the first application.
- Do not feed peanut forage or hay to meat or dairy animals.
- The combined total of preplant and postplant applications of Lorsban 4E and Lorsban 15G or other product containing chlorpyrifos must not exceed 4 lb of active ingredient per acre per crop season.

**SOYBEAN**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant and Postplant Treatments:** Apply as a T-band in a 4 to 10 inch band over the open seed furrow in front of the press wheel and incorporate into the top 1/2 to 1-inch of soil using suitable equipment. For postplant treatment, apply the granules in a 4 to 10 inch band over the row and incorporate into the top 1/2 to 1 inch of soil using a suitable cultivator. Apply Lorsban 15G with equipment that will provide uniform distribution of the granules over the row. Equivalent rates of Lorsban 15G per acre for various row spacing are given in Table 1.

Pests Controlled	Banded Applications (oz/1000 feet of row)	
	At Plant T-Band	Postplant Treatment
Cutworms	8	8
Lesser cornstalk borer		
Fire ants (suppression)	8	-

**Specific Use Precautions:**

- Do not apply as an in-furrow treatment to soybeans.

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not apply within 7 days before harvest of soybeans.
- Do not make more than 1 application of Lorsban 15G per season or apply more than 3 applications of other products containing chlorpyrifos.
- The maximum single application rate is 1.2 oz active chlorpyrifos per 1000 feet of row.
- Do not make a foliar application of Lorsban 4E or other product containing chlorpyrifos within 10 days of an at-plant application of Lorsban 15G.

**SUGARBEET**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant T-Band Treatment:** Apply Lorsban 15G in a band 4 to 5 inches wide behind the planter shoe, over the drill row, and in front of the press wheel. **Do not apply granules in direct contact with seeds.** Apply Lorsban 15G at the rate of 4.5 to 9 oz per 1000 feet of row (equivalent to 6.75 to 13.5 lb per acre based on 22-inch row spacing). Incorporate Lorsban 15G into the top 1/2 to 1-inch of soil using suitable equipment.

**Postplant Band Treatment:** Apply Lorsban 15G in a band 3 to 5 inches wide over the beet row (up to 2 to 4 true leaf stage of plant growth). Apply Lorsban 15G at the rate of 6.5 to 9 oz per 1000 feet of row (equivalent to 10 to 13.5 lb per acre based on 22-inch row spacing). Incorporate Lorsban 15G into the top 1/2 to 1 inch of soil using a suitable incorporation device.

Pests Controlled	Banded Applications (oz/1000 feet of row)	
	At Plant T-Band	Postplant Treatment
Sugar beet root maggot (1)	4.5 - 9.0	6.5 - 9.0
Cutworms	6.6 - 9.0	-
Wireworms (suppression)	6.5 - 9.0	-

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**Pest-Specific Use Directions:**

1. When **root maggot** populations are expected to be low, apply Lorsban 15G at a rate of 4.5 oz per 1000 feet of row (equivalent to 6.75 lb per acre based on 22-inch row spacing). If initial adult fly activity indicates higher than anticipated populations, apply Lorsban 4E at or near the time of peak adult emergence to augment control.

**Specific Use Precautions:** Granular insecticides, including Lorsban 15G, may contribute to the stress of the sugarbeet plant under certain environmental conditions. This stress may reduce plant stand or interfere with normal plant development. Herbicides used preplant incorporated may interact with insecticides and enhance this stress.

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not apply within 7 days before harvest of sugarbeets.
- Do not make more than 1 application of Lorsban 15G per season or apply more than 2 lb active chlorpyrifos per acre per season.
- Do not apply more than a total of 3 lb active chlorpyrifos per acre per year, or make more than 3 applications of products containing chlorpyrifos per season.
- Do not make a foliar application of Lorsban 4E or other product containing chlorpyrifos within 10 days of an at-plant application of Lorsban 15G.
- The maximum single application rate is 1.35 oz active chlorpyrifos per 1000 feet of row or 2 lb active chlorpyrifos per acre based on a 22-inch row spacing.

**SUNFLOWER**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At Plant T-Band Treatment:** Apply the granules in a 7-inch wide band over the row behind the planter shoe in front of the press wheel and incorporate into the top 1 inch of soil using suitable equipment.

Target Pests	Lorsban 15G (oz/1000 ft of row)
Cutworms	8

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not apply within 7 days before harvest of sunflowers.
- Do not make more than 1 application of Lorsban 15G per season or apply more than 3 applications of other product containing chlorpyrifos.
- Do not make a foliar application of Lorsban 4E or other product containing chlorpyrifos within 10 days of an at-plant application of Lorsban 15G.
- The maximum single application rate is 1.25 oz active per 1000 feet of row.

**SWEET POTATO**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Preplant Broadcast Treatment:** Apply Lorsban 15G in combination with a rotary applicator, such as a fertilizer spreader, to evenly broadcast the granules over the soil surface. Immediately following application, incorporate the granules uniformly into the soil to a depth of 4 to 6 inches using a rotary hoe or other incorporation device. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that Lorsban 15G will protect against feeding damage). Lorsban 15G will not control false wireworm or whitefringed beetle and other grubs that attack sweet potatoes.

Target Pests	Lorsban 15G (lb/acre)
<i>Conoderus</i> (wireworm)	13.5
<i>Systema</i> (flea beetle)	
Sweet potato flea beetle	

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not harvest within 125 days of treatment.
- Do not make more than 1 application of Lorsban 15G or other product containing chlorpyrifos per season.
- The maximum single application rate is 2 lb active chlorpyrifos per acre.

**TOBACCO**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Preplant Broadcast Treatment:** One week before transplanting, apply Lorsban 15G in combination with a rotary granular applicator, such as a fertilizer spreader, to evenly distribute the granules over the soil surface. Immediately following application, incorporate the granules uniformly into the soil to a depth of 2 to 4 inches using a rotary hoe or other incorporation device. The application of Lorsban 15G will also suppress movement of imported fire ant into a treated field.

Target Pests	Lorsban 15G (lb/acre)
Cutworms	13.5
Fire ant (suppression)	
Flea beetles	
Mole crickets	
Root maggots	
Wireworms	

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not harvest within 7 days of treatment.
- Do not make more than 1 application of Lorsban 15G or other product containing chlorpyrifos per season.
- The maximum single application rate is 2.025 lb active chlorpyrifos per acre.

**VEGETABLES (INCLUDING, BUT NOT LIMITED TO: BOK CHOY, BROCCOLI, BROCCOLI RAAB, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER, CHINESE BROCCOLI, CHINESE CABBAGE, COLLARDS, KALE, KOHLRABI, RADISH, RUTABAGA, AND TURNIP)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours (3 days for cauliflower) unless PPE required for early entry is worn.

**At Plant T-Band Treatment:** For direct seeded and transplanted crops, apply Lorsban 15G as a 4-inch wide band centered over the row. This application requires a banding device on the end of the applicator drop tube. Shallow incorporation is necessary. Application placement behind the planter shoe and in front of the press wheel is recommended.

Target Pests	Lorsban 15G (oz/1000 ft of row)
Root Maggots	4.6 - 9.2

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not harvest within 21 days of treatment for cauliflower and within 7 days for all other crops.
- Do not apply more than 7.5 lb of Lorsban 15G per acre to crops planted in 40-inch rows or more than 15 lb of Lorsban 15G per acre to crops planted in 20-inch rows (or two rows per bed). Use proportional amounts for other row spacings not to exceed 15 lb of Lorsban 15G per acre.
- Do not make more than 1 application of Lorsban 15G per season.
- Do not make a foliar application of Lorsban 4E or other product containing chlorpyrifos within 10 days of an at-plant application of Lorsban 15G.
- The maximum single application rate is 1.4 oz active chlorpyrifos per 1000 feet of row.
- Do not use rutabaga tops for food or feed purposes.

**RADISH**

**At Plant In-Furrow Treatment:** Apply the granules in the seed furrow at planting time so that the granules are covered with soil.

Target Pests	Lorsban 15G (oz/1000 ft of row)
Root maggots	3.3

**Specific Use Restrictions:**

- **Preharvest Interval:** Do not harvest within 7 days.
- The maximum single application rate is 0.5 oz active chlorpyrifos per 1000 feet of row (2.76 lb active chlorpyrifos per acre).
- Do not apply more than 18.3 lb of Lorsban 15G per acre or make more than 1 application per season.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Return:** DO NOT DISCARD THE CONTAINER. All SmartBox containers must be returned per instructions provided. Do not attempt to open or tamper with the container. Once empty, containers are not intended for reuse.

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE FULLEST EXTENT PERMITTED BY LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

Lorsban® is a registered trademark of Dow AgroSciences LLC  
SmartBox® is a registered trademark of Amvac Chemical Corporation

**AMVAC Chemical Corporation**

4100 E. Washington Boulevard  
Los Angeles, CA 90023 U.S.A.  
1-323-264-3910  
www.amvac-chemical.com