AGRI-TREND SOLUBLE

10-50-10

GUARANTEED ANALYSIS

Total Nitrogen (N)	10%
8.4% Ammoniacal Nitrogen	
1.6% Nitrate Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	50%
Soluble Potash (K ₂ O)	10%
	. 0 / 0
Trace Minerals:	
Boron (B)	0.02%
Boron (B) Copper (Cu)	0.05%
0.05% Chelated Copper	
Iron (Fe)	0.10%
0.10% Chelated Iron	
Manganese (Mn)	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo)	0.0005%
Molybdenum (Mo)Zinc (Zn)	0.05%
0.05% Chelated Zinc	

Derived From: Monoammonium Phosphate, Potassium Nitrate, Monopotassium Phosphate, Boric Acid, Sodium Molybdate and EDTA (Ethylenediaminetetraacetic Acid) Chelates of Copper, Iron, Manganese and Zinc.

Information regarding the contents and levels of metals in this product is available on the Internet at http://agr.wa.gov.

NOTE: The addition of this product will buffer the final solution. The pH will drop 1 to 2 digits depending on water hardness and rates.

NOTE: When mixing solubles with other chemicals, all chemical labels should be read in their entirety to ensure both chemical compatibility, and compatibility based on the pH.

CAUTION – KEEP OUT OF REACH OF CHILDREN

Warning: This product may contain chemicals known to the State of California to cause cancer, birth defects or other reproductive harm.

□ 25 lb. □ 50 lb.

LOT NUMBER



AGRI TREND SOLUBLE FORMULATION FEEDING SUGGESTION CHART

CROP	NO. OF TIMES TO APPLY	BEST TIME TO APPLY	(Each Applic.) AMOUNT PER ACRE
ALFALFA, CLOVER, ETC.	3	Apply desired formula when leaves appear after each cutting.	5 - 10 lbs.
SMALL GRAINS	2	When 3" to 5", again in boot stage.	7 - 10 lbs.
ONIONS, SUGAR BEETS, BEETS, CARROTS	2	When 4" to 6", again when tubers develop.	5 - 10 lbs.
POTATOES	4 to 6	Apply with each insecticide spraying.	5 - 10 lbs.
TOMATOES, PEPPERS, ETC.	4 to 6	Apply with each insecticide spraying.	3 - 5 lbs.
CORN, SWEET CORN, SORGHUM	2 to 3	When 6" to 8", then at 3 week intervals.	5 - 10 lbs.
BEANS, PEAS, SOYBEANS	2 to 3	When 5" to 6", just before flowering.	5 - 10 lbs.
RICE COTTON	2 to 4	Rice after growth is well established. Cotton with each insecticide spraying.	5 - 10 lbs.
TOBACCO	2 to 4	Use 1 lb. per 25 gal. water for transplant solution — then spray every 3 weeks.	5 - 7 lbs.
MISC. VEGETABLES		Each insecticide spraying.	5 - 10 lbs.
MELONS, SQUASH	2 to 4	With all insecticide spraying.	5 - 10 lbs.
BERRIES - GRAPES, STRAWBERRIES		Every 2 weeks after leaves appear the first season and once a month thereafter.	5 - 10 lbs.
FRUIT TREES	3 to 5	Add 1 lb. to each 30 gal. water at regular spraying.	
SHRUBBERY, FLOWERS, GARDENS		Sprinkle solution at base on ground also, spray foliage — 1 lb. to 25 gal. water.	
LAWNS	3 to 5	After raking in spring. Once every 2 or 3 weeks through early summer. 1 lb. to 25 gal. water (cover approx. 2,000 sq. ft.).	

NOTICE

Agri-Trend LLC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Agri-Trend LLC In no case shall Agri-Trend LLC be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Agri-Trend LLC makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.