

FERTI-MAG 4%

MICRONUTRIENT COMPLEX

GENERAL USE INFORMATION - READ ENTIRE LABEL BEFORE USING THIS PRODUCT

GENERAL INFORMATION

FERTI-MAG 4% is recommended for field, row, vegetable, fruit, tree, vine, ornamental, and turf crops.

FERTI-MAG 4%, alone or with pesticides or fertilizers, may be applied directly to soil or to the foliage of growing crops.

FERTI-MAG 4% may be applied via ground sprayers (high-volume or low-volume), aircraft or drip, trickle, center pivot, and lateral roll irrigation systems.

GENERAL DIRECTIONS FOR USE CONSULT YOUR LOCAL FERTIZONA REPRESENTATIVE OR OTHER AGRICULTURAL SPECIALIST FOR SPECIFIC CROP RECOMMENDATIONS.

Soil application of FERTI-MAG 4% is recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly-applied FERTI-MAG 4% most efficiently. Immediate post-irrigation or post-rainfall applications give best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Biuret Urea per 100 gals. of spray solution) will enhance foliar uptake of FERTI-MAG 4%.

AVOID foliar applications of FERTI-MAG 4% when plants are under moisture stress.

AVOID application to deciduous tree crops during bloom.

COMPATIBILITY: FERTI-MAG 4% is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

MIXING: ALWAYS add FERTI-MAG 4% to spray tank BEFORE adding pesticides.

NOTICE

READ ENTIRE LABEL BEFORE USE,
IF TERMS ARE UNACCEPTABLE RETURN AT ONCE – UNOPENED.

Fertilizer Company of Arizona, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Fertilizer Company of Arizona, Inc. In no case shall Fertilizer Company of Arizona, Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. The buyer shall assume all such risks. Fertilizer Company of Arizona, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



Distributed By:

FERTILIZER COMPANY OF ARIZONA, INC.

2850 S. Peart Road • Casa Grande, Arizona 85222 • (520) 836-7477

GUARANTEED ANALYSIS

Magnesium (Mg)4.0%	
Sulfur (S)	
Derived from magnesium sulfate and lignin sulfonate.	

NET WEIGHT: 10.2 LBS. PER GAL. AT 68° F.

1 Gal. (3.785 L), 10.2 lbs. (4.626 kg)

5 Gals. (18.93 L), 51 lbs. (23.128 kg)

_ Bulk

STORAGE AND DISPOSAL

Do not use aluminum or copper (including brass or bronze) components in any handling or storage equipment for this product. Mild steel components should not be used except for emergency, or temporary, short-term service. Use stainless steel (316 or better), polyolefin (polyethylene, polypropylene, etc.), fiberglass, or other high-quality acid-resistant plastic materials. Check with equipment manufacturer for compatibility of components prior to use with this product.

STORAGE INSTRUCTIONS: Protect from freezing. Store above 40°F. Freezing will not harm product's effectiveness. Store in original container tightly closed, do not allow water to be introduced into contents of this container.

CONTAINER DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Triple rinse (or equivalent), then offer container for recycling, or puncture and dispose in a sanitary landfill or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SHAKE, STIR OR SWIRL CONTENTS BEFORE USING SPECIFIC DIRECTIONS FOR USE

ALL CROPS:*

SOIL APPLICATIONS

Soil Fertility Maintenance	
Light Deficiency	
Severe Deficiency	1-3 gals, per acre

*The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, vine, ornamental, and turf crops.

FOLIAR APPLICATIONS

Apply 2-4 qts. per acre as needed on a 7-14 day schedule.

AVOID foliar applications of FERTI-MAG 4% when plants are under moisture stress.

DILUTE APPLICATIONS

Apply the suggested dose of FERTI-MAG 4% in enough water for good coverage.

CONCENTRATE APPLICATIONS

Use up to 2 qts. of FERTI-MAG 4% in a minimum of 25 gals. of water.

AERIAL APPLICATIONS

Up to 1 qt. of FERTI-MAG 4% per gal. of spray may be used, but more dilute solutions will improve coverage and results.

ALL IRRIGATION APPLICATIONS

Apply the recommended amount of FERTI-MAG 4% in split applications 1-2 weeks apart throughout the growing season.

CAUTIONKEEP OUT OF REACH OF CHILDREN

Fertizona 008 (8/05)
Yuma Printing Co.