

JIMMY SANDERS, INC.

BEFORE SEED · BEYOND HARVEST

**SANDERS**

**QUALITY**

**PRODUCT**

CLEVELAND, MS

**SILSURF**

**NONIONIC ORGANOSILICONE SURFACTANT**

**PRINCIPLE FUNCTIONING AGENTS:**

Polyalkylene Modified Heptamethyltrisiloxane and nonionic surfactants .....	99%
Constituents ineffective as a spray adjuvant.....	1%
Total.....	100%

All ingredients are exempt from tolerance requirements under 40 CFR 180.

**CAUTION**

**KEEP OUT OF THE REACH OF CHILDREN**

**NET CONTENTS: 1 GALLON**

**PEEL BACK BOOK HERE AND RESEAL AFTER OPENING**



## PRECAUTIONARY STATEMENTS

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

## FIRST AID

Call a poison control center or doctor immediately for treatment advice.

**If Swallowed:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. Have the product container with you when calling a poison control center or doctor, or going for treatment.

## GENERAL INFORMATION

SIL-SURF is a unique nonionic organosilicone wetting agent, that is highly effective with postemergent and non-selective herbicides. SIL-SURF's organosilicone chemistry, allows for enhanced wetting and absorption of herbicides into plant, leaf, and stem surfaces. SIL-SURF may be used with most pesticides and fertilizer products as directed on this label. Observe initial application to insure thorough coverage without excessive runoff of the spray.

## MIXING

Prior to pesticide application, clean mixing and application equipment according to pesticide label directions. To minimize foaming fill spray tank 2/3 or 3/4 full of water and begin agitation. If foaming is anticipated, the addition of an antifoaming agent should be added before pesticides, nutrients, or SIL-SURF.

Add pesticides and/or fertilizers as directed by label or in the following sequence: 1. Micronutrients and fertilizers, 2. Dry flowables and dispersible granules, 3. Flowables, 4. Water soluble pesticides, 5. Emulsifiable concentrates.

Continue agitation and add SIL-SURF, when completed, reduce agitation and allow to mix for 1 to 2 minutes. While under minimal agitation, continue filling tank and allow to mix for 10-15 minutes prior to application.

NOTE: For optimum results, spray mixes containing SIL-SURF should be applied within 36 hours.

## DIRECTIONS FOR USE

**SIL-SURF** when directed for use with pesticides and fertilizers, is effective when applied on crops such as alfalfa, corn, cotton, oats, peanuts, rice, sorghum, soybeans, sugar cane, wheat, lawns, turf, greens and tees.

### USE RATES

APPLICATION	RATE PER 100 GALLONS
Ground	12 to 64 ounces
Aerial	12 to 32 ounces
Aquatic	8 to 32 ounces
General Wetter/Spreader	6 to 12 ounces

For increased water penetration of hard to wet soils and the uniform distribution of applied moisture, follow recommended uses:

Lawns and Turf: .05% - .125% v/v concentration.

Greens and Tees: .05% - .125% v/v concentration.

**NOTE:** The application rates on this label are based on pesticide labels, which recommend the use of a nonionic surfactant. Rates of this product may be increased or decreased for optimum results, follow pesticide labeling for proper recommendations. Before using SIL-SURF where a nonionic surfactant may not be recommended, the user or applicator advisor must have experience with the combination, or must have conducted a phytotoxicity trial.

## STORAGE AND DISPOSAL

Do not contaminate food or feed by storage or disposal. Do not store near heat or open flame. Store in original container.

**Pesticide:** Wastes resulting from the use of this product may be disposed of on site, or at an approved waste disposal facility.

**Container:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration in accordance with local, state, or federal regulations.

## NOTICE

SELLER WARRANTS THAT THE PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE UNDER NORMAL CONDITIONS. NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESSED OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL DIRECTIONS, OR UNDER ABNORMAL CONDITIONS, OR UNDER CONDITIONS NOT FORESEEABLE TO SELLER. BUYER ASSUMES RISK UNDER ANY SUCH USE INSUFFICIENT CONTROL OF PESTS AND/OR CROP INJURY MAY RESULT FROM UNUSUAL ENVIRONMENTAL CONDITIONS, FAILURE TO FOLLOW PESTICIDE LABEL DIRECTIONS OR POOR APPLICATION PRACTICES, ALL OF WHICH ARE BEYOND THE CONTROL OF JIMMY SANDERS, INC.

JIMMY SANDERS, INC.

BEFORE SEED · BEYOND HARVEST

**SANDERS**

**QUALITY**

**PRODUCT**

CLEVELAND, MS

**SILSURF**

**NONIONIC ORGANOSILICONE SURFACTANT**

**PRINCIPLE FUNCTIONING AGENTS:**

Polyalkylene Modified Heptamethyltrisiloxane and nonionic surfactants .....	99%
Constituents ineffective as a spray adjuvant.....	1%
Total.....	100%

All ingredients are exempt from tolerance requirements under 40 CFR 180.

**CAUTION**

**KEEP OUT OF THE REACH OF CHILDREN**

**NET CONTENTS: 1 GALLON**