

*Hummert's*

**DYNA  
GREEN™**

Your Prescription for a Picture Perfect Lawn™

# Crabgrass Preventer Lawn Fertilizer With Dimension®

covers approximately 5,400 sq. ft.

**15-0-5**

- *Excellent pre-emergence and early-post-emergence control of crabgrass, therefore it can be used later in the season than other crabgrass preventers*
- *Can be applied on dormant or spring seeded lawns after the second mowing*
- *Spring nutrient package requires less mowing*

For information on our complete product line,  
go to **[www.dynagreeninfo.com](http://www.dynagreeninfo.com)**

Net Wt. 18 lbs. (8.20 kg)

© 2013 Hummert International

*Additional information on back of bag.  
Please read and follow the directions.*



With



*Hummert's*  
**DYNA  
GREEN™**

**Crabgrass  
Preventer  
Lawn Fertilizer  
With  
Dimension**







# Crabgrass Preventer

with 0.172% Dimension®  
15-0-5



## PREVENTS CRABGRASS

SELECTIVE HERBICIDE PLUS FERTILIZER FOR  
PREEMERGENCE CONTROL OF LISTED ANNUAL  
GRASSES AND BROADLEAF WEEDS IN  
ESTABLISHED LAWNS & ORNAMENTAL TURF  
ACTIVE INGREDIENT:  
Dithiopyr: 3,5-pyridinedicarbothioic acid,  
2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-  
S,S-dimethyl ester . . . . . 0.172%  
OTHER INGREDIENTS: . . . . . 99.828%  
TOTAL: . . . . . 100.000%

EPA Reg. No. 32802-69  
EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B),  
37351-CO-1 (C), 9404-FL-1 (E), 8378-IN-1 (D)

First letter of bag code shows establishment where manufactured.  
Read the entire "Directions for Use" and "Warranty  
Statement" before using this product. If terms are not  
acceptable, return the unopened product container at once  
to the seller.

## KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes eye irritation. Harmful if inhaled. Avoid contact  
with eyes or clothing. Avoid breathing dust. Wash thoroughly with  
soap and water after handling. Remove contaminated clothing and  
wash before reuse. Prolonged or frequently repeated skin contact may  
cause allergic reaction in some individuals.  
**ENVIRONMENTAL HAZARDS:** To protect the environment, do not  
allow this product to enter or run off into storm drains, drainage  
ditches, gutters or surface waters. Applying this product in calm  
weather will help to ensure that wind does not blow this product  
off the treated area. Sweeping any product that lands on a  
driveway, sidewalk, or street, back onto the treated area of the  
lawn or garden will help to prevent run off to water bodies or  
drainage systems..

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:** Store in original container in a locked  
storage area out of reach of children and domestic animals.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this  
product may be disposed of on site or at an approved waste  
disposal facility.  
**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or  
refill this container: IF EMPTY: Place in trash or offer for recycling if  
available. IF PARTLY FILLED: Call your local solid waste agency  
for disposal instructions. Never place any unused product down  
any indoor or outdoor drain.  
**NONREFILLABLE CONTAINER:** Do not use or refill this container.  
Offer for recycling, if available  
**IF PARTLY FILLED:** Call your local solid waste agency for disposal  
instructions. Never place any unused product down any indoor or  
outdoor drain.

Manufactured by:  
Howard Johnson's Enterprises, Inc.  
9675 S 60th Street  
Franklin, WI 53132  
Information regarding the contents and levels of metals in this product is  
available on the internet at <http://www.aapfco.org/metals.htm>  
\*Dyna Green® with fertilizer to help green up turf.



**GUARANTEED ANALYSIS**  
Total Nitrogen (N). . . . . 15.00%  
14.50% Urea Nitrogen  
0.50% Nitrogen from Organic Poultry Compost  
Soluble Potash (K2O). . . . . 5.00%  
Boron (B) . . . . . 0.05%  
Copper (Cu) . . . . . 0.13%  
0.03% Water Soluble Copper (Cu)  
Iron (Fe) . . . . . 0.25%  
0.03% Water Soluble Iron (Fe)  
Manganese (Mn) . . . . . 0.13%  
0.03% Water Soluble Manganese (Mn)  
Molybdenum (Mo) . . . . . 0.0013%  
Zinc (Zn) . . . . . 0.13%  
0.03% Water Soluble Zinc (Zn)  
Derived From: COPPER OXIDE, COPPER  
SULFATE, IRON OXIDE, IRON SULFATE,  
MANGANESE OXIDE, MANGANESE  
SULFATE, SODIUM BORATE, SODIUM  
MOLYBDATE, ZINC OXIDE, ZINC SULFATE,  
MURIATE OF POTASH, UREA  
Chlorine (Cl), not more than. . . . . 6.00%  
**Net Weight 18 lbs (8.20 kg)**  
**One bag covers up to 5,400 square feet**  
**FGB65101**

**WARNING**  
**Some crops may be injured by application of boron. This  
fertilizer is to be used only on soils which respond to molyb-  
denum. Crops high in molybdenum are toxic to ruminants.**

**First Aid**  

<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or physician immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or physician.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

**Hot line number:**  
Have the product container or label with you when calling a  
poison control center or doctor, or going for treatment.  
For information on this pesticide product (including health  
concerns, medical emergencies, or pesticide incidents),  
call the National Pesticide Telecommunications Network  
at 1-800-858-7378.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
In the State of New York: For use by professional turf applicators only, do not apply more than 6.67 lbs per 1000 square feet per  
year (0.5 pounds AI per acre per year). Not for use in Nassau or Suffolk Counties.  
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.  
**PRODUCT INFORMATION:** This product contains turf fertilizer plus a selective herbicide for control of crabgrass and other annual  
grasses and broadleaf weeds in established lawns and ornamental turf. Apply this product only to seeded, sodded, or sprigged turf  
that is well established. Turf must develop a good root system and uniform stand and have received at least two mowings following  
seeding, sodding or sprigging before making the first application of this product. Use of this product on turf that is not well established,  
or weakened by weather, pests or other stress, may increase the chance of turf injury.  
This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of  
crabgrass from the ground. In addition, this product provides "early post-emergence" control during the early stages of growth after  
crabgrass emerges from the ground. This early post-emergence crabgrass control is limited, and post-emergence applications are  
only effective if applied before crabgrass has displayed its fifth leaf or first tiler (which generally corresponds to time when you can  
first see the crabgrass plants in the lawn). The benefit of this post-emergence activity is that it gives the user a 2 to 4 week longer  
period of time to make applications for effective control of crabgrass.

**WHERE TO USE:** This product may be used on the following established turfgrasses: Creeping Bentgrass\* (*Agrostis palustris*),  
Kentucky Bluegrass (*Poa pratensis*), Fine Fescue\*\* (*Festuca rubra*), Tall Fescue (*Festuca arundinacea*), Perennial Ryegrass  
(*Lolium perenne*), Bahiagrass (*Paspalum notatum*), Bermudagrass (*Cynodon dactylon*), Buffalograss\*\*\* (*Buchloe dactyloides*),  
Carpetgrass (*Axonopus compressus*), Centipedegrass (*Eremochloa ophiuroides*), Kikuyugrass (*Pennisetum clandestinum*), St.  
Augustinegrass (*Stenotaphrum secundatum*), Zoysiagrass (*Zoysia japonica*).  
\*Turfgrass injury may result from use of this product on certain varieties of creeping bentgrass, such as Cohansey, Carmen, Seaside  
and Washington. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Colonial Bentgrass (*Agrostis  
tenuis*) varieties.\*\*Turfgrass injury may result from use of this product on the following Fescue varieties which are sensitive to this  
product: Atlanta, Banner, Beauty, Bilgart, CF-2, Enjoy, HF-93, Highlight, Ivalo, Koket, Jamestown, Majenta, Mary, Pennlawn,  
Tamara, Tatjana, Walfort and Waldina. Not all varieties of fescue have been tested.

\*\*\*Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and  
has established new roots.  
**SPRING APPLICATIONS:** For single applications in the spring or early summer, use the appropriate rate from "Application Rates"  
table (depending on user's location, mowing height, pre- or post-emergent use). The duration of residual weed control is directly  
related to the total rate applied, but varies depending on weather, weed pressure and location. Use "Program 1" for preemergence  
control on areas where turf is cut high (home lawns) for 3 to 5 months of crabgrass control. This program should not be used for  
postemergence crabgrass control. Use "Program 2" for preemergence control where turf is cut low (golf course fairways) or where weed  
control was not conducted the previous year. This program provides 4 to 6 months of preemergence weed control, and also for early  
postemergence control of crabgrass. This product may be applied as a single application or as a split application for crabgrass control  
in the spring, summer or fall.

APPLICATION RATES for Crabgrass Control		
REGION	PROGRAM 1 PREEMERGENCE USE (high cut turfgrass)	PROGRAM 2 PREEMERGENCE USE (low cut turfgrass) POSTEMERGENCE USE
North	1.66 lbs / 1000 sq ft (73 lbs / acre 0.125 lbs AI / acre)	3.33 lbs / 1000 sq ft (145 lbs / acre 0.25 lbs AI/acre)
Transition	2.45 lbs / 1000 sq ft (107 lbs / acre 0.183 lbs AI / acre)	4.9 lbs / 1000 sq ft (213 lbs / acre 0.366 lb AI / acre)
South	3.33 lbs / 1000 sq ft (146 lbs / acre 0.25 lbs AI / acre)	6.67 lbs / 1000 sq ft (291 lbs / acre 0.50 lbs / acre)
Coastal South	4.9 lbs / 1000 sq ft (213 lbs / acre 0.366 lb AI / acre)	6.67 lbs / 1000 sq ft* (291 lbs / acre* 0.50 lbs AI / acre*)
North: All other areas. Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY and RI South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN and TX Coastal South: HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, TX *May require split or sequential applications for full control.		

**RESEEDING, OVERSEEDING, OR SPRIGGING:** Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product  
may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best  
results, use "slit seeders" for good seed-to-soil contact.  
**CONTROL OF OTHER GRASS AND BROADLEAF WEEDS:**  
**APPLICATION LIMITS:** Certain states have limits on application rates and timing. Consult your local State Cooperative Extension Service for current  
information on application limits for lawn fertilizers in your area.  
**SPRING APPLICATIONS:** Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to  
their emergence: Barnyardgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green)\*, Goosegrass, Kikuyugrass, Lespedeza (common),  
Oxalis (buttercup, creeping & yellow woodsorrel), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (corn), Spurge (prostrate and  
spotted). \*Foxtail is also controlled at the program 2 rates.  
**FALL APPLICATIONS:** Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control  
or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Bittercress, Chickweed, Geranium (Carolina), Henbit,  
Parsley-piert, Pineappleweed, Shepherd's purse.

SPREADER SETTINGS	to apply at the listed pounds per 1000 sq. ft.				
	1.66 lbs	2.45 lbs	3.33 lbs	4.9 lbs	6.67 lbs
Dyna Green Broadcast	8	10	13.5	15.5	17.5
Ames Drop or Rotary	8	11	14	17	19
EarthWay Drop or Rotary	8	11	14	17	19
Scotts EasyGreen EG-3 (rotary)	25	26	27	28	29
Scotts SpeedyGreen (rotary)	3.5	4.5	6	8	11

**Warranty**  
The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated  
with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions,  
presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Hummert International, the manufactur-  
er, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Hummert International. The manufacturer  
and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this  
label Hummert International, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either  
express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to,  
merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable  
law, Buyer's or user's exclusive remedy, and Hummert International, the manufacturer's or the seller's total liability shall be limited to damages not  
exceeding the cost of the product. No agent or employee of Hummert International, or the seller is authorized to amend the terms of this warranty  
disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product. To the extent  
consistent with applicable law, Hummert International, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages  
resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user  
waive any right that they may have to such damages.

Settings shown are approximate for new  
equipment. Your spreader may need calibration,  
due to differences in terrain, walking speed and  
equipment condition. Calibrate your spreader  
to ensure proper coverage. Clean & lubricate  
spreader after use.