

**ACTIVE INGREDIENTS**n-OCTANOL (Fatty Alcohol C<sub>8</sub>) 35.5%n-DECANOL (Fatty Alcohol C<sub>10</sub>) 49.5%

INERT INGREDIENTS.....15.0%

TOTAL..... 100.0%

Weight of Active Ingredients: 6.01 lb. per gallon

Formulated from a fatty alcohol mixture that typically contains: C<sub>6</sub> - 1.0%; C<sub>8</sub> - 41.0%; C<sub>10</sub> - 57.0%; C<sub>12</sub> - 0.8%

Linear Fatty Alcohols

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See Side Panel For Additional  
Precautionary Statements.

Manufactured By:  
**Coastal AgroBusiness, Inc.**  
Greenville, North Carolina

E.P.A Reg. No. 5549-74

E.P.A. Est. No. 5549-NC-1

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING/AVISO:** Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid inhalation of mist.

### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed:** Call a physician or poison control center immediately. Drink promptly a large quantity of milk, eggwhites, gelatin solution, or if these are not available, large quantities of water.

Avoid alcohol. Do not give anything by mouth to an unconscious person.

**If in Eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. If irritation persists, seek medical attention immediately.

**If on Skin:** Wash with plenty of soap and water. If irritation develops, seek medical attention.

**If Inhaled:** Remove to fresh air. If not breathing, give artificial respiration and seek medical attention immediately.

### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, Neoprene Rubber, Polyvinyl Chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning, maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply this product through any type of irrigation system.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### STORAGE AND DISPOSAL

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by

storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide spray mixture or rinse that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**KLEEN-TAC '85'** is carefully balanced combination of active ingredients and wetting agents to be used for the control of sucker growth on Burley, Flue-Cured, Dark, Maryland, and Cigar Tobacco. The concentrated product is diluted with water to form a creamy emulsion which is then applied as a course spray. The emulsion is effective only when it comes in direct contact with suckers; therefore, the material is applied so that maximum contact is made with the suckers.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, Neoprene Rubber, Polyvinyl Chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

**WHEN TO APPLY**

**KLEEN-TAC '85'** can be applied before or after topping. Best results are usually obtained by spraying the tobacco with **KLEEN-TAC '85'** before topping, in early to late button stage, and then topping the tobacco within a few days, removing any suckers that were missed. If you top the tobacco plants before spraying, remove any suckers over one inch in length as you top and apply **KLEEN-TAC '85'** one or two days after topping. Because **KLEEN-TAC '85'** is a contact type agent, it is necessary to straighten any plants that are leaning so that the emulsion flows down the stalk and evenly contacts each sucker. **KLEEN-TAC '85'** can be applied any time during the day. However, for best results it is recommended that you wait until the dew dries before spraying. Do not spray after the leaves begin to close in the evening. Because the underside of the leaves may be injured by contact with **KLEEN-TAC '85'** do not apply when the wind is high enough to turn top leaves over. Do not apply during rain or when plants are wet. If, however, it rains after **KLEEN-TAC '85'** has been on the plants for over an hour, you should not have to apply **KLEEN-TAC '85'** again.

**HOW MUCH TO APPLY**

<b>FLUE-CURED:</b>	For Power Sprayer - use 1 1/2 - 2 gallons in 50 gallons of water. For Hand Sprayer - use 4-5 ounces per gallon of water.
<b>BURLEY:</b>	For Power Sprayer - use 1 3/4 - 2 gallons in 50 gallons of water. For Hand Sprayer - use 4-5 ounces per gallon of water.
<b>DARK:</b>	Use 6-8 ounces per gallon of water to apply with hand sprayer.
<b>CIGAR:</b>	Use 4-5 ounces per gallon of water to apply with hand sprayer.
<b>MARYLAND:</b>	Use 4 - 4 1/2 ounces per gallon of water to apply with hand sprayer.

When applied by hand, 1 gallon of diluted **KLEEN-TAC '85'** will treat approximately 190 plants. If a power sprayer is used, 40-60 gallons of diluted product should be applied per acre of tobacco.

**HOW TO APPLY**

The diluted emulsion is most easily prepared by adding the required amount of **KLEEN-TAC '85'** to your spray tank and then by adding the water. In order to obtain the best results, it is important that

the water be added to the **KLEEN-TAC '85'** rather than **KLEEN-TAC '85'** to the water. If you use a hand-held or backpack sprayer, the diluted emulsion should be applied at a rate of 2/3 to 1 ounce per plant (or enough to insure run down to the bottom of the plant). A course spray is recommended, directed downward at the top of the stalk from 6-8 inches above the top leaves. Very little tank pressure is required, and in no case should more than about 20 pounds be used. When applied with power equipment, three nozzles per row should be used (TG-3 full-cone tips, or equivalent is satisfactory). Replacing the center TG-3 nozzle with a TG-5 full-cone nozzle, which provides more spray solution down the stalk, is also recommended. One nozzle should be directed downward over the center of the row and one should be 9 inches on either side directed at or slightly above the top of the stalk. The diluted **KLEEN-TAC '85'** should be applied to the tobacco as a course spray from a height of 12-16 inches above the stalk. It is recommended that boom pressure be kept at 20-25 lbs. and should never exceed 25 lbs. By using the recommended spray tips, spraying at approximately 20 lbs. of pressure, and operating at a tractor speed of 2 1/2 - 3 mph, you will apply approximately 50 gallons of diluted emulsion per acre of tobacco.

**HOW OFTEN TO APPLY**

In many cases a single application of **KLEEN-TAC '85'** is satisfactory to control both primary and secondary suckers. If secondary sucker growth appears vigorous, a second treatment 5-7 days after the initial treatment may be advantageous. Some growers of Burley, Flue-Cured, and Dark tobacco may favor a dual treatment using **KLEEN-TAC '85'** first (at the early button stage) followed 1-2 weeks later with a systemic sucker control agent (maleic hydrazide) used in accordance with manufacturer's instruction. If this dual treatment is used on Burley or Flue-Cured tobacco, an application rate of 1 1/2 - 2 gallons of **KLEEN-TAC '85'** in 50 gallons of water is adequate. If this dual treatment is used on Dark tobacco, **KLEEN-TAC '85'** should be used at the rate of 6-8 ounces per gallon of water to apply by hand sprayer.

**NOTE:** If allowed to stand for extended periods of time, the diluted emulsion may separate. Always remember to mix well immediately prior to use. It is recommended that **KLEEN-TAC '85'** not be mixed with other chemicals or applied to tobacco plants which have recently been treated with other chemical. Be sure that

all spray equipment has been carefully rinsed out prior to using. In Burley tobacco, during prolonged periods of high heat and humidity or under other conditions which might favor the spread of hollow stalk or stem rot diseases, delay application of **KLEEN-TAC '85'** until these conditions pass.

**LATE SEASON APPLICATION**

To avoid application of more than the labeled rate or Maleic Hydrazide and/or to avoid reapplying Maleic Hydrazide late in the season when initial sucker control starts to break down, apply Kleen-Tac '85' 3-4 weeks after Maleic Hydrazide application. This late application should be made when suckers are less than one inch in length or when suckers more than one inch in length have been removed by hand.

Late application of Kleen-Tac '85' involves the use of 5% Kleen-Tac '85' applied 3-4 weeks after MH application.

Flue-Cured:	For Power Sprayer	Use 2-1/2 gallons in 47-1/2 gallons of water.
	For Hand Sprayer	Use 6-7 ounces per gallon of water.

When applied by hand, 1 gallon of diluted Kleen-Tac '85' will treat approximately 190 plants. If a power sprayer is used, 40-60 gallons of diluted product should be applied per acre of tobacco. In order for this late application treatment to be effective, Kleen-Tac '85' should be applied when suckers are less than one inch or when suckers longer than one inch have been removed by hand. Because Kleen-Tac '85' is a contact type agent, it is necessary to evenly apply the material so that maximum contact is made with the suckers.

**NOTICE:** The concentrations and application techniques recommended above have been found to control suckers with little or no leaf injury. Concentrations, application rates, sprayer pressures, and time of application other than those recommended by lead to inadequate sucker control and/or injury to the tender upper leaves of the plant. Neither **Coastal AgroBusiness** nor the seller, makes any representation of warranty, expressed or implied, as to this product other than that contained on the label thereof, and the buyer assumes all responsibility of personal injury, property damage, or other loss resulting from use, handling, or storage of the product, and accepts and uses the material on these conditions.