IPCO® Ag-Surf

LIQUID SPRAY ADJUVANT

AGRICULTURAL

FOR USE WITH ROUNDUP¹, ALLY², REGLONE³, and ROVRAL 50% W.P.⁴

READ THE LABEL BEFORE USING

GUARANTEE: Nonylphenoxy polyethoxyethanol 92%

REGISTRATION NO: 15881

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 10 litres

Interprovincial Cooperative Limited, P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9

®T.M. Reg'd. RC 636-0105

AG-SURF

GENERAL INFORMATION

Ag-Surf is a surfactant for use with Roundup¹, Ally² and Reglone³ herbicides, and Rovral 50% W.P.⁴ fungicide, in situations where a non-ionic surfactant is called for (minimum or zero tillage systems surfactant additive, etc.). It is important to follow label directions for Roundup, Ally, and Reglone herbicides and Rovral fungicide, using the recommended rates in combination with Ag-Surf.

PRECAUTIONS: KEEP OUT OF THE REACH OF CHILDREN.

Do not store or place near open flames or sources of heat. Do not apply more surfactant than recommended or excessive run-off may occur. Avoid contact with eyes.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- (1) Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- (2) Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID: If in eyes or if on skin, rinse with clean water for 15 minutes. If irritation persists, get medical attention. If swallowed, do not induce vomiting. If conscious, dilute by giving 2 glasses of water. Call a physician or a poison control centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO USER

This control product is to be used in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising out of the use or handling of the product and accepts the product on that condition.

DIRECTIONS

With Roundup: Minimum/Zero-Tillage/Summerfallow: To improve the effectiveness for post-emergent control of annual grasses and broadleaf weeds in Minimum/Zero-Till systems for wheat, oats, barley or on summerfallow, apply Ag-Surf at 350 mL/ha plus Roundup at 1.1 L/ha in 50 - 100 L/ha of clean water. See Minimum and Zero Tillage section of Roundup label for weeds controlled and other details of application. Quackgrass: When applying the "Quackgrass Control" rate of 2.5 L/ha of Roundup and water volumes of 150 - 300 L/ha, add Ag-Surf at 0.5% (0.5 litres/100 litres of water) to maintain adequate surfactant concentration in spray solution. See "Quackgrass Control" section of Roundup label for other details of application.

With Ally: For maximum weed killing effect – after the addition of Ally to water in the tank add 200 mL of Ag-Surf for each 100 litres of spray solution (0.2% total spray volume). An antifoaming agent may be needed to reduce foaming.

With Regione: To improve the effectiveness for crop desiccation, post-emergent weed control in non-crop land, chemical mowing and potato vine killing, add Ag-Surf at 1 L per 1000 Litres of spray where the addition of a wetting and spreading agent is recommended on the Reglone label.

With Rovral 50% W.P.: For use in rapeseed only. For the control of Sclerotinia Stem Rot add Ag-Surf at the rate of 0.5% of the final spray solution (500 mL in 100 litres).

®T.M. of Interprovincial Cooperative Ltd.

¹T.M. of Monsanto Co.

²T.M. of Dupont of Canada

³T.M. of Syngenta Crop Protection Canada, Inc. ⁴T.M. of Bayer CropScience Inc.

RC 641-0105