

BIOMIN[®] BOOSTER 235

1-0-0

Amino Acid Chelated Minerals for Soil and Foliar Application



Product Information

BIOMIN BOOSTER 235 contains amino acid chelated minerals for soil and foliar application. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. It is designed to prevent and correct mineral deficiency and to boost crops during critical stages or fast growing periods. It can be applied in combination with nitrogen fertilizer.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	1.0%
1.0% Water Soluble Nitrogen	
Magnesium (Mg).....	0.5%
0.5% Chelated Magnesium	
Boron (B).....	0.025%
Copper(Cu).....	0.5%
0.5% Chelated Copper	
Iron (Fe).....	1.0%
1.0% Chelated Iron	
Manganese(Mn)	1.5%
1.5% Chelated Manganese	
Zinc (Zn)	2.5%
2.5% Chelated Zinc	

Derived from magnesium sulfate, sodium borate, cupric sulfate, ferrous sulfate, manganese sulfate, zinc sulfate, and hydrolyzed vegetable protein.

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

**PREVENTS AND CORRECTS
MINERAL DEFICIENCIES**

NET CONTENTS: 2.5 gal. (9.45 l)

Weight per Gallon: 10.5 lb. (4.8 kg) @ 68°F

U.S. Pat.# 5,504,055

Lot #:



Directions and General Recommendations

SOIL APPLICATION

BIOMIN BOOSTER 235 can be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1 qt. to 1 gal./ac. (2.5-10 l/ha) depending on soil conditions and application methods. For maintenance, 1 qt./ac. (2.5 l/ha) is recommended. Dilute **BIOMIN BOOSTER 235** with enough water to cover the entire acre.

FOLIAR APPLICATION

FIELD CROPS: Apply 16-32 oz./ac. (1.25-2.5 l/ha) of **BIOMIN BOOSTER 235** when crops are 6-12 in. (15-30 cm) high or when deficiency symptoms appear. Repeat application 2 - 4 weeks later if necessary.

FRUIT TREES AND GRAPES: Apply 16-32 oz./ac. (1.25-2.5 l/ha) of **BIOMIN BOOSTER 235** starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS: Apply 16-32 oz./ac. (1.25-2.5 l/ha) of **BIOMIN BOOSTER 235** at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 1-3 weeks later.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

ALMONDS, WALNUTS, PISTACHIOS, AND OTHER NUT CROPS: Apply 16-32 oz./ac. (1.25-2.5 l/ha) before bloom. Repeat the application if necessary.

For best results, apply **BIOMIN BOOSTER 235** when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, dilute it with at least 100 times of water. **BIOMIN BOOSTER 235** will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN BOOSTER 235 can be applied to strawberry, lettuce, tomato, celery, stone fruits, vegetables, grapes, commercial turf, and ornamental gardens.

Precautionary statement: do not swallow. Wash hands thoroughly after handling. **First Aid:** if swallowed: Seek medical attention if feeling unwell. Rinse mouth. **Storage & Disposal:** store in a safe manner. Dispose product/container in accordance with federal, state, and local regulations.

JH Biotech Inc.

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006 U.S.A. Phone: 805-650-8933 Website: www.jhbiotech.com

Copyright © 2015 All rights reserved

0311-D-B-2