

# **BIOMIN CALCIUM** 2-0-0



## Amino Acid Chelated Mineral for Soil and Foliar Application

#### PRODUCT INFORMATION

BIOMIN CALCIUM contains amino acid chelated calcium designed for soil and foliar application. It is completely bioavailable and nonphytotoxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizer.

#### DIRECTIONS AND GENERAL RECOMMENDATIONS:

SHAKE WELL BEFORE USING SOIL APPLICATION: BIOMIN CALCIUM can be applied before or at planting as banding or broadcast. The application rate is 1 quart - 1 gallon per acre (2.5 to 10 liters/hectare) depending on soil conditions and application methods. For maintenance, 1 quart per acre (2.5 liters/hectare) is recommended, BIOMIN CALCIUM should be diluted with enough water to cover the entire acre.

#### FOLIAR APPLICATION:

FIELD CROPS: Apply 1-3 quarts per acre (2-8 liters/hectare), when crops are about 6-12 inches (15 - 30 cm) high or when deficiency symptoms appear. Repeat the application in 2-4 weeks if necessary.

FRUITTREES AND GRAPES: Apply 1-3 quarts per acre (2-8 liters/hectare) starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS: Apply 1-3 quarts per acre (2-8 liters/hectare) at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1-3 weeks if necessary.

#### **GUARANTEED ANALYSIS** Tatal Allerana (Al)

Total Nitrogen (N)	.0%
2.0% Water Soluble Organic Nitrogen	
Calcium (Ca) 5	.0%
5.0% Chelated Calcium	

Contains Calcium Carbonate, Citric Acid and Glycine as chelating agents. Nitrogen derived from Glycine.

**PREVENT** AND CORRECT MINERAL DEFICIENCIES

**KEEP OUT OF REACH** Of CHILDREN

Manufactured by: JH BIOTECH, INC. P.O. Box 3538

Ventura, CA 93006 USA

Information regarding the contents and levels of metals in this product is available on the internet at http://agr.wa.gov

For best results, apply when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water, BIOMIN CALCIUM will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN CALCIUM can be applied to STRAW-BERRY, LETTUCE, TOMATO, CELERY, GRAPES, STONE FRUITS, TREE FRUITS, VEGETABLES, COMMERCIAL TURF AND ORNAMENTAL GARDENS.

#### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neithermanufacturernorsellershall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

### **NET CONTENTS:**

- 1 Pint (0.47 liters) 1 Gallon (3.78 liters)
- 2.5 Gallons (9.45 liters)
  - 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters) 55 Gallons (207.9 liters)

Weight per Gallon: 10.8 lbs. (4.9 kg.)

Lot #:

U.S. Pat.# 5.504.055