

# NUTRI-AID

12-26-26

## Water Soluble Fertilizer With Micronutrients



### GUARANTEED ANALYSIS

Total Nitrogen (N)	12%
7.0% Nitrate Nitrogen	
5.0% Ammoniacal Nitrogen	
Available Phosphoric Acid ( $P_2O_5$ )	26%
Soluble Potash ( $K_2O$ )	26%
Boron (B)	0.02%
Copper (Cu)	0.05%
Iron (Fe)	0.10%
Manganese (Mn)	0.05%
Zinc (Zn)	0.05%

Derived from ammonium phosphate, potassium nitrate, solubor, copper sulfate, ferrous sulfate, manganese sulfate, zinc sulfate, and citric acid.

### DIRECTIONS

DILUTE APPLICATION:	3 to 5 lbs. per 100 gallons of water
SEMI-CONCENTRATE:	5 to 10 lbs. per acre in 40 to 100 gal of water
AERIAL APPLICATION:	3 to 10 lbs per acre in sufficient water for good coverage

Fill tank with water and slowly add **NUTRI-AID** while agitating. Continue to agitate until all **NUTRI-AID** is completely dissolved.

### APPLICATION IN IRRIGATION WATER

Solid Set, Overhead, Drip Sprinkler or Pivot Irrigation Systems: Dissolve **NUTRI-AID** in mix tank and apply 8-10 lbs. per acre in irrigation water. Use of anti-siphon or check valve is recommended to prevent solution from running back into primary water supply.

**NUTRI-AID** is generally compatible with pesticides and other fertilizers. Determine compatibility with a jar test before mixing products.

**BEANS, PEAS, LENTILS, RAPESEED** (Canola): Use 5 to 10 lbs. per acre. Apply at first sign of bloom. Repeat 2 weeks later.

**BERRIES:** (Strawberries, raspberries, blackberries, blueberries,) Use 5 to 10 lbs. per acre in spring on new growth. Repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

**CARROTS, BEETS, SWEET POTATOES, AND OTHER VEGETABLE ROOT CROPS:** Use 5 to 10 lbs. per acre per application.

**CITRUS INCLUDING ORANGE, LEMON, GRAPEFRUIT, TANGERINE:** Use 10 to 20 lbs. per acre per application. In concentrate spray, use 10 to 12 lbs. per acre per application in 20 to 100 gallons of water. In dilute sprays, use 3 to 4 lbs. per 100 gallons of water and wet foliage completely.

**COLE CROPS, LETTUCE, LEAFY VEGETABLES:** Use 5 to 10 lbs. per acre. Apply right after thinning or transplanting. Repeat every 10 to 14 days.

**CORN:** (Field and Sweet): Use 5 to 10 lbs. per acre. Apply when corn is 6" - 8" high and again prior to tasseling.

**COTTON:** Use 4 to 10 lbs. per acre per application in sufficient water to give even distribution of material. May be applied as general maintenance 4 to 5 times per season.

**CRANBERRIES:** Use 5 to 10 lbs. per acre. Apply at popcorn stage. Repeat at hooking and 3 weeks later.

**DECIDUOUS FRUITS** (Apples, pears, cherries, plums, prunes, peaches, almonds, walnuts, filberts, pistachios and pecans): Apply in delayed dormant, pre-bloom, and post-bloom sprays at 3 to 5 lbs. per 100 gal. in dilute sprays. Apply 10 to 15 lbs. per acre in concentrate sprays. Avoid combining with material that will cause russetting.

**FIELD CROPS** (Mint, alfalfa, clover): Use 5 to 10 lbs. per acre. Apply when plants have 6" to 8" new growth. Repeat application to new growth after each cutting.

**GRAPES:** Ground sprayers: Use 5 to 10 lbs. per acre in 50 to 100 gal. of water. Air Application: Use 5 to 8 lbs. per acre in 10 gal. of water. Apply when leaves emerge, pre-bloom and post-bloom until grapes are 1/4 size.

**HOPS:** Ground strayers: Use 5 to 10 lbs. per acre in 100-200 gallons of water. Air Application: Use 5 to 8 lbs. per acre in no less than 10 gal. of water. Apply when hops are halfway up wire and again when they reach the top of the wire.

**MELONS, CUCUMBERS AND OTHER CUCURBITS:** Use 5 to 10 lbs. per acre per application. Apply two or more applications every 2 to 3 weeks beginning with pre-bloom.

**ONIONS, GARLIC, TOMATOES, PEPPERS:** Use 5 to 10 lbs. per acre per application. Begin application prior to bloom. Repeat every 2-3 weeks thereafter.

**POTATOES:** Use 5 to 10 lbs. per acre. Apply when plants are 4" - 6" high (hooking stage). Subsequent applications should be at 2 to 3 week intervals.

**SAFFLOWER, SUNFLOWER, AND OTHER OILSEED CROPS:** Use 5 to 10 lbs. per acre per application.

**SMALL GRAINS** (Wheat, oats, barley, rye): Use 5 to 10 lbs. per acre in sufficient water to cover by air or ground. Apply at tillering and again at early boot stage.

**SUGAR BEETS:** Use 5 to 10 lbs. per acre. Apply when leaves are 10" across. Repeat application 2 to 4 weeks later.

### CONDITIONS OF SALE AND LIMITED WARRANTY:

The directions for use of this product reflect the opinions of experts based on field uses and laboratory testing. Always follow label instructions. Manufacturer warrants that this product conforms to the product description on this label. Manufacturer or seller makes no other expressed or implied warranty. Neither manufacturer nor seller shall be liable for any injury or any application not specifically described on the label.

**CAUTION: KEEP OUT OF REACH OF CHILDREN - HARMFUL IF SWALLOWED  
AVOID BREATHING DUST**

**MANUFACTURED BY:** JH BIOTECH, INC.  
P.O. Box 3538  
Ventura, CA 93006

**NET WEIGHT:** 50 LBS (22.5 Kg.)

SPECIMEN LABEL

**JH BIOTECH, INC.**