

# MINOR ELEMENT SOLUBLE

Soluble Trace Element Fertilizer for Nurseries, Turf, and Landscapes.

Use for the Correction and Prevention of Minor Element Deficiencies (yellowing foliage).

## DIRECTIONS FOR USE

**FOLIAR APPLICATION:** For full volume sprays use 1/2 to 2 pounds per 100 gallons. Use lower rates for prevention of deficiencies, and higher rates for correction of deficiencies on hardy plants. Only use lower rates on sensitive plants.

**CONSTANT FEED INJECTION SYSTEMS:** Use 1/2 pound for every 100 pounds of regular soluble fertilizer.

**DRENCH APPLICATION:** To established plants for the correction of deficiencies as a one time treatment, use 1/2 pound per 100 gallons. Drench to wet root area.

**WARM SEASON TURF GRASS:** Use 2 - 5 lb. in at least 100 gallons of water per acre. Repeat applications as needed or desired for color.

**COOL SEASON TURF GRASS:** Use 2 - 2 1/2 lb. in at least 100 gallons of water per acre. Repeat applications as needed.

### MINIMUM GUARANTEED ANALYSIS:

Soluble Manganese as Mn .....	8.15%
Soluble Iron as Fe .....	7.50%
Soluble Copper as Cu .....	3.20%
Soluble Zinc as Zn .....	4.50%
Boron as B .....	1.45%
Molybdenum as Mo .....	0.046%
Combined Sulfur as S .....	13.00%
Chlorine not more than .....	0.1%

**DERIVED FROM:** Copper Sulfate, Manganese Sulfate, Iron Sulfate, Zinc Sulfate, Sodium Borate, and Sodium Molybdate.

**GUARANTEED BY:** Southern Agricultural Insecticides, Inc.

**USE PRECAUTIONS:** Do not mix with other fertilizers or pesticides unless compatibility is known. When using on new crops or varieties, test rate and application techniques on a few plants before treating entire crop to check for phytotoxicity. **KEEP OFF SIDEWALKS, DRIVEWAYS, PORCHES AND ALL MASONRY SURFACES OR SERIOUS STAINING WILL RESULT.** Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material.

F 672      J2      Net Weight: 25 lbs.(11.3 kg)

## CAUTION:

**KEEP OUT OF REACH OF CHILDREN.**

**May be harmful if swallowed.** Avoid breathing dust, and contact with skin, eyes and clothing. If ingested, give two glasses of water, induce vomiting and call a physician. If in eyes or on skin flush with water. If on clothing, launder before reuse. Wash thoroughly after handling.

Contains more than .001% Molybdenum and should not be used on plant parts used for forage or feed for livestock.

This material does not contain Magnesium or Calcium, these elements should be added for healthy plants.

# Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998