11.05 LBS. PER GALLON @ 68°F 1235 GRAMS PER LITER

APPLICATION RATES

Use a total of 2-3 gallons perseason if needed:

ALFALFA CLOVER - 1 to 2 qts. per acre per application. For hay, apply at 6 to 8 inches of new growth and after each cutting. For seed, apply at bud stage and repeat at intervals of 7 to 10 days.

APPLES, APRICOTS, CHERRIES, NECTARINES, PEARS, PLUMS, AND PRUNES - 1 to 4 gts. per acre per application. Apply pre and /or post-bloom and repeat at intervals of 7 to 10 days.

HOPS - 1 to 2 qts. per acre.

NUT CROPS (ALMONDS, FILBERTS, PECANS, PISTACHIOS, AND WALNUTS) - 1 to 4 gts. per acre per application. Repeat every 7 to 10 davs.

ARTICHOKES - 1 to 2 qts. per acre per application. Apply bud stage, repeat every 7 to 10

ASPARAGUS - 1 to 2 gts. per acre per application. Apply just after start of new growth. Repeat every 7 to 10 days.

AVOCADOS - 1 to 2 qts. per acre perapplication. BEANS, SOYBEANS, PEAS, LENTILS, - 1 to 2 qts. per acre . First application at bud stage. Repeat every 7 to 10 days.

BLUEBERRIES, RASPBERRIES - 1 to 2 qts. per acre pre bloom. Repeat every 7 to 10 days.

CARROTS, BEETS SWEET POTATOES AND OTHER VEGETABLE ROOT CROPS - 1 to 3 ats. per acre when plants are three to four weeks old. Repeat every 7 to 10 days.

CITRUS - 2 to 4 gts. per acre per application prebloom. Then every 7 to 10 days.

CORN - 1 to 4 qts. per acre.

COTTON - 1 to 2 gts. per acre

CRANBERRIES - 1 to 2 qts. per acre. First application at hook stage, second application after fruit set. WARRANTY: Western Nutrients Corp. makes no warranty, express or implied, including the warranties of merchantability and/ FIGS - 1 to 3 qts. per acre

GRAPES - 1 to 4 qts. per acre. First application at the 4 to 6 leaf stage, repeat every 7 to 10 days as needed. Use enough dilution for full coverage.

Nutraplex® is a Registered Trademark of Western Nutrients Corporation.



NUTRAPLEX_® **CAL MAG** 5-0-0 **Plus Boron**

GUARANTEED ANALYSIS

Total Nitrogen (N)	5.6%
5.6% Nitrate Nitrogen	
Calcium (Ca)	6.5%
Magnesium (Mg)	0.6%
Boron (B)	0.5%
Copper (Cu)	.05%
.05% Chelated Copper	
Iron (Fe)	.10%
.10% Chelated Iron	
Manganese (Mn)	.10%
.10% Chelated Manganese	
Zinc (Zn)	.20%
.20% Chelated Zinc	

Derived from Calcium Nitrate, Boric Acid, Magnesium Nitrate. Copper. Iron, Manganese, and Zinc Edta. ALSO CONTAINS NON - PLANT FOOD INGREDIENTS

KEEP OUT OF REACH OF CHILDREN

or fitness for any particular purpose, concerning this material, except those which are contained on the Western Nutrients Corp. Label attached to the product container.

WARNING: EXCESS AMOUNTS OF BORON (B) CAN CAUSE PLANT INJURY.

MANUFACTURED BY WESTERN NUTRIENTS CORPORATION

245 Industrial Street, Bakersfield, Ca. 93307 Phone (661) 327-9604/Fax (661) 327-1740 E-mail: westernnutrients@lightspeed.net Website: http://www/westernnutrients.net

NET CONTENTS 5 GALLONS 18.93 LITERS

APPLICATION RATES

LETTUCE, CELERY, AND LEAFY VEGETABLE CROPS - 1 to 2 qts. per acre. Repeat every 15 days. **GRAPES -** 2 - 6 quarts per acre five times each season starting at first full leaf, then past bloom, early bunch set, berry size and at berry color and just before harvest.

MELONS AND CUCUMBERS - 1 to 2 gts. per acre. Make applications at intervals of 7 to 10 days starting iust prior to bloom.

MINT - 1 to 2 qts. per acre.

OLIVES - 1 TO 2 gts. per acre.

ONIONS AND GARLIC - 1 to 2 qts. per are.

PEANUTS - 1 to 2 gts. per acre. First application at early bloom stage, repeat at intervals of 7 to 10 days. **POTATOES** - 1 to 2 gts. per acre per application. First application when plants are 4 to 12 inches high. Second application 3 weeks later, repeat every 10 days.

SAFFLOWER AND SUNFLOWER - 1 to 2 gts. per acre per application.

STRAWBERIES - 1 to 2 ats, per acre per applica-

SUGAR BEETS - 1 to 3 qts. per application. First application when plants are 6 to 8 inches across. Second application 3 to 4 weeks later, repeat as needed.

TOMATOES AND PEPPERS - 1 to 2 qts. per appli-

VEGETABLES - 1 to 3 qts. per acre. First application at bud stage, repeat applications at intervals of 7 to 10 days.

WHEAT, BARLEY, RICE, AND OTHER SMALL **GRAINS** - 1 to 2 qts. per acre per application. First application at boot stage, repeat 7 to 10 days.

SOIL - 2 to 5 gallons per acre on all crops.

DILUTION RATE: Use sufficient water for thorough coverage. Maximum dilution 1 gt. per 10 gallons of

GENERAL USE - Nutraplex is compatible with most herbicide, pesticides, fungicides, and UAN solution. Use ample water for coverage. Can be used with all types of application equipment.