



PGR 113 MC

For Turf Growth Management

A microemulsion concentrate for managing growth, improving quality, and stress tolerance and edging of warm- and cool- season turfgrasses

ACTIVE INGREDIENT: % BY WT
Trinexapac-ethyl
(CAS No.: 95266-40-3)11.3%
OTHER INGREDIENTS: ...88.7%
TOTAL:100.0%

EPA Reg. No. 86064-3

EPA Est. No. 37429-GA 001^{PT};
37429-GA-002^{BO}

Letter(s) in lot number correspond(s)
to superscript in EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

Manufactured by:

United Turf Alliance, LLC
8014 Cumming Highway,
Suite 403-282
Canton, GA 30115

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **For emergency information call Poison Control at 1-800-222-1222 or the National Pesticide Information Center (NPIC) at 1-800-858-7378.**

NOTE TO PHYSICIAN

If a large amount has been ingested, lavage stomach. An aqueous suspension of activated charcoal can be given to absorb remaining toxicant. Treat symptomatically.

See inside label booklet for additional Precautionary Statements and Directions for Use including Storage and Disposal instructions.

Net Contents: 1 gallon (3.78 liters)



United Turf Alliance

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as nitrile, butyl, neoprene, or barrier laminate)
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

The restricted-entry interval (REI) for this product is 0 days.

Do not enter treated areas without footwear until sprays have dried.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without footwear until sprays have dried.

Observe all precautions and limitations on this label and on the labels of each product used in tank mixture with this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN UNEVEN GROWTH REDUCTION OR SEVERELY STUNTED TURF.

ArmorTech PGR 113 MC is a plant growth regulator and is an odorless micro emulsion concentrate (MEC). ArmorTech PGR 113 MC mixes completely with water and may be tank mixed with many commonly used pesticides and liquid fertilizers.

ArmorTech PGR 113 MC is effective only by absorption through the grass foliage and not by plant uptake of the product through the soil. Within one hour after application, ArmorTech PGR 113 MC will not wash off the grass foliage. A number of factors affect the activity and performance of ArmorTech PGR 113 MC when used for growth management of turfgrasses including but not limited to the following:

- Environmental conditions
- Turfgrass management and cultural practices that affect grass growth and vigor
- Level of plant fertility
- Availability of moisture
- Plant vigor
- State of grass growth

Watering-in is not necessary for activation of ArmorTech PGR 113 MC.

When an adjuvant is to be used with this product, United Turf Alliance, LLC suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

TURFGRASS: ArmorTech PGR 113 MC reduces the growth of warm- and cool-season turfgrasses. When used according to directions for turf growth management, ArmorTech PGR 113 MC reduces the frequency of mowing and the amount of grass clippings generated by mowing. ArmorTech PGR 113 MC applications may also increase the color, quality, and density of turf.

ArmorTech PGR 113 MC can be applied to the following sites that have well-maintained, quality turfgrass areas:

- lawns (residential and commercial)
- golf courses
- sod farms
- sports fields, cemeteries, and similar areas
- difficult-to-mow areas

ArmorTech PGR 113 MC minimizes the need for edging of turfgrass along sidewalks, curbs, parking lots, driveways, flower beds, fences and around posts, storage sheds, and trees.

USE DIRECTIONS FOR APPLICATION TO TURFGRASS

MOWING

ArmorTech PGR 113 MC generally provides greater suppression when turfgrasses are maintained at lower mowing heights and the application rates in this label have been selected for typical mowing heights; be sure to adjust the application rate based on the actual mowing conditions under which ArmorTech PGR 113 MC will be applied. To minimize possible turf injury, apply ArmorTech PGR 113 MC at least 4 hours before mowing or 1 hour after mowing.

EQUIPMENT

ArmorTech PGR 113 MC may be applied using backpack sprayers, hand sprayers, boom sprayers, and with spray gun application devices. Be sure to clean the spray equipment thoroughly before use and make sure the sprayer is calibrated and capable of accurate and uniform application. Rinse sprayer with clean water after use and dispose of rinsate according to label directions.

MIXING INSTRUCTIONS

1. Add ArmorTech PGR 113 MC to a sufficient amount of water (ex., 0.5 to 4.0 gallons of water per 1,000 sq. ft.)
2. Mix only the volume of spray mixture required for the square footage to be treated.
3. Ensure adequate agitation during application.
4. When ArmorTech PGR 113 MC is used as a tank mix with other products, agitation is required.
5. Apply ArmorTech PGR 113 MC in a uniform manner to ensure thorough coverage of the crop.

For tank mixtures, use only if the mixing partner is registered for this use. Do not mix ArmorTech PGR 113 MC with any product which prohibits such mixing. Do not exceed any label application rate and follow the most restrictive label precautions and limitations. Refer to the tank mix partner label(s) for further information.

Backpack and Hand Sprayers (0.5-4.0 gal. capacity)

ArmorTech PGR 113 MC Alone: Add all of the required water to the mix tank followed by the appropriate amount of ArmorTech PGR 113 MC. Close the sprayer and vigorously shake. Begin application.

Boom and Hand Gun Sprayers

ArmorTech PGR 113 MC Alone: Add all the required amount of water to the spray tank. Then, while agitating, add ArmorTech PGR 113 MC.

ArmorTech PGR 113 MC Plus Tank Mixtures: Before mixing ArmorTech PGR 113 MC with other components, determine the compatibility of the tank mix using the **TANK MIX COMPATIBILITY TEST** below. Once the compatibility of the mixture is assured, add the products (such as carrier or other pesticide products) to the spray tank in the order noted below:

1. Add ½ of the required amount of water to the spray tank.
2. Keep the tank mix agitated to ensure formation of a uniform suspension.
3. Add any products that are packaged in water-soluble bags first. Keep the tank mix agitated to ensure that the water-soluble bags dissolve and the contents dispersed in the tank mix. Continue agitation and then add water-dispersible granules (WG), followed by wettable powders (WP). Before adding other products, ensure complete dispersal of the solid products.
4. Add emulsifiable concentrates (EC).
5. Add flowable liquids (FL) or suspension concentrates (SC).
6. Add ArmorTech PGR 113 MC.
7. Add spray adjuvant and spray markers. Use surfactants approved for application to turf. Check surfactant label before use.
8. Add the remainder of the water.
9. Agitate tank mix combinations. Do not let mixtures sit in the spray tank without agitation. Only prepare the amount of spray mix required for use on one day and apply it all that day.

TANK MIX COMPATIBILITY TEST

Add relative proportionate amounts of ArmorTech PGR 113 MC, other products to be used, and the water in a clear glass quart jar with lid. Close the lid and invert the jar several times and observe the mixture for approximately 30 minutes. **The mixture is not compatible if the following is observed:** formation of balls, flakes, sludges, gels, oily films or layers; or precipitates.

APPLICATION NEAR AND AROUND MONUMENTS AND HARDSCAPE MATERIALS

At normal dilution rates, ArmorTech PGR 113 MC will not stain brass, bronze, concrete, marble, granite, or other types of stone. Prior to using ArmorTech PGR 113 MC around other materials, test on a small scale basis first.

PRE-STRESS CONDITIONING OF TURFGRASS

Multiple ArmorTech PGR 113 MC applications, in conjunction with cultural practices such as fertilization, irrigation, drainage and mowing height will delay the onset of stress, improve stress survival, and enhance turf's recovery from stress. Apply ArmorTech PGR 113 MC to healthy, actively growing turf before the onset of stress and continue to apply ArmorTech PGR 113 MC throughout the growing season as long as the turf remains healthy. Do NOT apply more than 7.0 fl. oz./ 1,000 sq. ft. per year.

When turf top growth slows, energy is redirected to below-ground plant parts, and root and rhizome production increases resulting in root-mass that is often greater in ArmorTech PGR 113 MC -treated turf than in similar, untreated turf.

Research indicates that ArmorTech PGR 113 MC may also enhance the performance of fungicides. Monthly applications of ArmorTech PGR 113 MC at the label rate or biweekly application at ½ the label rate may strengthen the turfgrass to help it resist disease. Additionally, if mowing is less frequent, and leaf material removal is reduced, contact and systemic products will remain at effective concentrations on or in the turf longer.

Multiple ArmorTech PGR 113 MC applications will result in smaller, more compact turfgrass and reduced leaf area, resulting in reduced transpiration, water use and improved drought tolerance. Increased turf density will also reduce moisture loss from soil evaporation, and additional soil moisture will be available due to increased root depth and mass. ArmorTech PGR 113 MC may also enhance cold and heat tolerance by increasing carbohydrate levels.

SUPPRESSION OF ANTHRACNOSE

ArmorTech PGR 113 MC can be applied at rates of 0.1 to 0.2 fl. oz. per 1,000 sq. ft. every 7 to 14 days to suppress basal rot anthracnose. Use the lower rate when seedheads are present. For optimal disease control, tank mix with products containing propiconazole, chlorothalonil, Medallion®, or other fungicides registered for the control of anthracnose.

POA ANNUA CONVERSION/RENOVATION

As part of an overseeding / renovation program allowing better germination and seedling growth of more desirable turf, ArmorTech PGR 113 MC may be applied to existing turf infested with stands of *Poa annua*. By combining ArmorTech PGR 113 MC with appropriate cultural practices that help to ensure the vigor and growth of new seedlings, fewer clippings and reduced maintenance traffic on new seedlings will result. Additionally, seed germination is not affected because ArmorTech PGR 113 MC is absorbed foliarly. NOTE: Aggressive application rates of ArmorTech PGR 113 MC may result in temporary initial discoloration.

Apply ArmorTech PGR 113 MC 1-5 days before seeding and before verticutting, scalping, spiking or other similar operations. The following spring, apply the recommended upper end rate of ArmorTech PGR 113 MC for the turf type and setting listed in the Application Rate Table. Actual conversion success will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions.

BERMUDAGRASS OVERSEEDING

Establishment of cool-season turfgrasses can be enhanced by applying ArmorTech PGR 113 MC to bermudagrass. By combining ArmorTech PGR 113 MC with appropriate cultural practices that help to ensure the vigor and growth of new seedlings, fewer clippings and reduced maintenance traffic on new seedlings will result. Additionally, seed germination is not affected because ArmorTech PGR 113 MC is absorbed foliarly. NOTE: Aggressive application rates of ArmorTech PGR 113 MC may result in temporary initial discoloration.

Apply ArmorTech PGR 113 MC to the bermudagrass 1-5 days before seeding or before verticutting, scalping, spiking, or other similar operations. For maintenance applications, see the Application Rate Table. Use normal seeding rates for your area and turf setting (lawn, fairway, etc.). Actual overseeding results will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions.

APPLICATION WITH TURF MARKING PAINT

When applied before or with marking agents, ArmorTech PGR 113 MC can extend the duration of marking visibility. When combining with latex-based marking agents, mix ArmorTech PGR 113 MC with water first and refer to the marking agent's products label for further instructions.

To treat approximately 1,000 sq. ft. of line surface area, apply ArmorTech PGR 113 MC at 1 oz./gal of marking paint mix. Refer to the ArmorTech PGR 113 MC plus Tank Mixtures section of this label for additional instructions.

PLANT TOLERANCE

The grasses listed on this label have been shown to be tolerant of ArmorTech PGR 113 MC. It is impossible to test every species and cultivars of grasses for tolerance to ArmorTech PGR 113 MC. Before commercial use, the professional user should determine if ArmorTech PGR 113 MC can be used safely on grasses not listed in the Application Rate Table by conducting small-scale tests. The lower rate for turf setting (lawn, fairway, etc.) should be tested and evaluated for phytotoxicity and growth inhibition prior to use on a large scale.

USE RESTRICTIONS

- Do NOT apply ArmorTech PGR 113 MC through any type of irrigation system.
- Do NOT graze treated areas or feed treated clippings to livestock.

USE PRECAUTIONS

- Areas treated with ArmorTech PGR 113 MC should continue to receive regular maintenance practices, including irrigation; fertilization; and weed, disease, and insect control when necessary and as recommended for quality turf.
- Because some herbicides can injure turf, tank mixes with ArmorTech PGR 113 MC should be tested on a small scale before widespread use.
- ArmorTech PGR 113 MC may cause temporary yellowing that usually disappears approximately one week after application. To minimize yellowing and to enhance the green color of turf, apply readily available nitrogen at a rate of 0.2-0.5 lb. of actual nitrogen per 1,000 sq. ft. If desired, recommended rates of iron per 1,000 sq. ft. can also be used.

APPLICATION INSTRUCTIONS

Excessive turf growth during spring flushes or as a result of fertilization may require higher rates of ArmorTech PGR 113 MC (up to 50% greater) to provide an adequate length of control.

For extended growth suppression (up to 8 weeks) when temporary discoloration can be tolerated, a maximum of twice the ArmorTech PGR 113 MC rate from the Application Rate Table may be applied.

ArmorTech PGR 113 MC use rates may need to be reduced up to 50% for turfgrass grown under conditions of low fertility, compaction, or other factors, which stress the turf.

Multiple applications of ArmorTech PGR 113 MC can be made each growing season, but do not exceed a total of 7.0 fl. oz./1,000 sq. ft. (305 fl. oz./A; 19.0 pts./A) per year.

Application Timing

Apply ArmorTech PGR 113 MC to actively-growing turf. A lower rate of ArmorTech PGR 113 MC should be used if turf is going into dormancy because of high or low temperatures or lack of moisture.

As soon as the turf resumes growth or if more suppression is desired, repeat applications of ArmorTech PGR 113 MC may be made, but do NOT apply more than 7.0 fl. oz./1000 sq. ft. per year.

Table 1. ArmorTech PGR 113 MC Application Rates¹

NOTE: The rates listed below should provide 50% suppression of turf growth with minimal yellowing for 4-weeks under good growing conditions.

Turf Type	Residential and Commercial turf ²		Golf Course Fairways (cut at 1/2" or less)		Golf Course Greens & Tees		Edging/Banding ³	
	fl oz/ 1000 sq ft	fl oz/A	fl oz / 1000 sq ft	fl oz/A	fl oz/ 1000 sq ft	fl oz/A	fl oz/ 1000 sq ft	fl oz/A
Cool Season¹								
Bentgrass	0.75	33	0.25	11	0.125	6	1.0	44
Fescue, red	0.75	33					1.0	44
Fescue, tall (Ky-31)	1.0	44					1.0	44
Fescue, tall (Turf Types)	0.75	33					1.0	44
Kentucky Bluegrass	0.60	26	0.25	11			0.75	33
Mixture (Bentgrass/ <i>Poa annua</i>)			0.25 ⁴	11	0.125	7		
Mixture (K. Bluegrass/ Fescue/ Ryegrass)	0.75	33						
Mixture (K. Bluegrass/ Ryegrass/ <i>Poa annua</i>)			0.50 ⁴	22				
Ryegrass, Annual	1.0	44					1.0	44
Ryegrass, Perennial	1.0	44	0.50	22			1.0	44
Warm Season¹								
Bahiagrass	1.0	44					1-2	44-88
Bermudagrass⁵								
Bermudagrass Common	0.75	33	0.25	11			1-2	44-88
Bermudagrass Other Hybrids	0.25	11	0.20	9			0.50-0.75	22-33
Bermudagrass, Tifdwarf	0.20	9	0.20	9	0.062	3	0.50-0.75	22-33
Bermudagrass, Tifgreen (328)	0.25	11	0.20	9	0.125	6	0.75	33

(continued)

Turf Type	Residential and Commercial turf ²		Golf Course Fairways (cut at 1/2" or less)		Golf Course Greens & Tees		Edging/Banding ³	
	fl oz/ 1000 sq ft	fl oz/A	fl oz / 1000 sq ft	fl oz/A	fl oz/ 1000 sq ft	fl oz/A	fl oz/ 1000 sq ft	fl oz/A
Bermudagrass, Tifway (419)	0.38	16	0.25	11			0.75	33
Buffalograss	1.0	44					1.0	44
Carpetgrass	0.25-0.40	11-18					0.50	22
Centipedegrass	0.50	22					1.0	44
Kikuyugrass	0.30-0.50	13-22	0.3	13			1.0	44
Seashore Paspalum	0.375	16	0.125-0.25	6-11	0.08-0.20	4-9	0.50	22
St. Augustinegrass	0.10-0.15	4.50-6.50					0.40-0.80	18-36
St. Augustinegrass Texas Common	0.10	4.5					0.20	9
Zoysiagrass	0.25	11	0.125	6			0.75	33

1. Seedheads: At rates equal to or higher than the rates in Table 1, ArmorTech PGR 113 MC provides seedhead suppression of hybrid bermudagrass, and partial seedhead suppression of annual bluegrass, bahiagrass, buffalograss, carpetgrass, common bermudagrass, Kentucky bluegrass, and tall fescue. ArmorTech PGR 113 MC must be applied prior to seedhead formation. Do not apply more than 7.0 fl. oz./ 1000 sq. ft. per year.
2. Including, but not limited to: home lawns, parks, recreation areas, golf course roughs, cemeteries, business sites, sports fields, and sod farms.
3. ArmorTech PGR 113 MC can be applied along the perimeter of lawns, sidewalks, curbs, parking lots, driveways, posts, storage buildings, pet pens, fences, or other areas. ArmorTech PGR 113 MC can be used around trees, shrubs, flower beds, and other border plants or similar areas with no injury. Apply ArmorTech PGR 113 MC in an 8-12 inch band with a single nozzle sprayer. Use the higher concentration to reduce the growth of turfgrass into adjacent areas.
4. If yellowing of *Poa annua* is a concern, use half this rate and apply more frequently. For *Poa annua* conversion/renovation, where temporary discoloration can be tolerated, twice this rate may be applied.
5. Bermudagrass overseeding preparation: 0.5 oz/ 1000 sq. ft. for golf fairways and tees; 0.25 oz./1000 sq. ft. for golf greens.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and avoid excess heat. Do not store below 0°F. If crystals form, store above 70°F, shaking periodically until crystals are dissolved.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Turf Alliance, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Turf Alliance, LLC and Seller harmless for any claims relating to such factors.

United Turf Alliance, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or United Turf Alliance, LLC, and Buyer and User assume the risk of any such use. **UNITED TURF ALLIANCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither United Turf Alliance, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED TURF ALLIANCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED TURF ALLIANCE OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Turf Alliance, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of United Turf Alliance, LLC.

ArmorTech is a trademark of United Turf Alliance, LLC.

Medallion is a registered trademark of a Syngenta Group Company.