

# CH2<sup>®</sup>

Ornamental Herbicide  
Provides preemergence control of weeds in  
container and field grown ornamentals

**This specimen label is for information only.**  
Everris makes no representations as to the accuracy of this label. It is the responsibility of the user to read and follow the label attached to the pesticide product container.

**Net Weight: 50 lbs (22.68 kg)**  
**Stock # E95715**

GROUP 3 14 HERBICIDES

**ACTIVE INGREDIENTS:**

Oxyfluorfen <sup>†</sup> .....	2.00%
Pendimethalin <sup>††</sup> .....	1.00%
<b>OTHER INGREDIENTS:</b> .....	<b>97.00%</b>

Total 100.00%

<sup>†</sup>CAS # 42874-03-3

<sup>††</sup>CAS # 40487-42-1

**EPA Reg. No. 58185-178**

**EPA Est. 82757-OH-1<sup>MF</sup>, EPA Est. 8378-IN-1<sup>K</sup>**

Superscript used is the first letter of run code on end of container.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**FIRST AID (continued)**

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment. You may also call 1-888-241-6409 for emergency medical assistance.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes eye injury. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid inhalation.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Mixers, loaders, applicators and other handlers must wear:**

- coveralls over long-sleeved shirt and long pants
- chemical-resistant footwear plus socks
- chemical-resistant gloves
- chemical-resistant apron for mixers and loaders

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**If you have questions regarding the use of this product, call TOLL FREE 1-800-492-8255**

Distributed by Everris NA Inc., PO Box 3310 • Dublin, OH 43016

## USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Use with care when applying in areas adjacent to any body of water. Runoff from treated areas may be hazardous to organisms in neighboring areas. See Directions for Use for additional restrictions. Apply this product only as specified on the label. Do not contaminate water when disposing of wash water. In case of spills, either collect for use or dispose of properly.

## General Information

OH2<sup>®</sup> Ornamental Herbicide is a pre-emergent herbicide for container and field grown ornamentals in nurseries and does not control established weeds. OH2 Ornamental Herbicide combines two selective herbicides on one granule, giving the user a broad spectrum of weed control and ornamental plant tolerance. OH2 Ornamental Herbicide is not registered for use on vegetable or edible fruit crops, turf, bedding plants or flowering crops except as described in the plant tolerance list. For season-long weed control, repeat herbicide applications are required at approximately 3-month intervals during the growing season.

## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep all persons, children and pets out of the treated areas until dusts have settled.

For use only on container, field grown, and landscape ornamentals. This herbicide product has been extensively researched; however, testing has not been conducted on all species or cultivars, or under all growing conditions. In some instances performance characteristics may vary.

## RESISTANCE MANAGEMENT

OH2 Ornamental Herbicide is both a Group 3 and a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to Group 3 and/or Group 14 herbicides. Weed species with acquired resistance to Group 3 and/or Group 14 herbicides may eventually dominate the weed population if Group 3 and/or Group 14 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by OH2 Ornamental Herbicide or other Group 3 and/or Group 14 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of OH2 Ornamental Herbicide or other target site of action Group 3 or Group 14 herbicides that have a similar target site of action, on the same weed species.
- Using tank-mixtures or premixes with herbicides from different target site of action groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.

- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.
- Contacting Everris at 800-492-8255 for further information or to report suspected resistance.

**THE SPECIES LISTED BELOW HAVE BEEN TESTED UNDER ACTUAL GROWING CONDITIONS AND FOUND TOLERANT AT THE RECOMMENDED APPLICATION RATE.**

**Abelia**

x grandiflora (Glossy Abelia)

**Acacia**

latifolia longifolia (Bush Acacia)

**Acer**

palmatum 'Atropurpureum' (Red Cutleaf Japanese Maple)

**Achillea**

millefolium 'Cerise Queen' (Yarrow)

**Araucaria**

heterophylla (Norfolk Island Pine, Star Pine)

**Azalea Kurume, Glen Dale hybrid, et al.**

'Coral Bells', 'Fashion', 'Hardigzer Beauty', 'Hershey's Red', 'Mother's Day', 'Mrs. G.G. Gerbing', 'Twenty Grand' (Azalea)

**Berberis**

spp. 'Crimson Pygmy' (Barberry)  
thunbergii 'Atropurpurea' (Red-Leaf Japanese Barberry)

**Betula**

papyrifera (Paper Birch, Canoe Birch)

**Brassaia (see Schefflera)\***

**Buxus**

microphylla 'Green Velvet'  
microphylla japonica (Japanese Boxwood)  
microphylla koreana (New Large Leaf Asiatic Boxwood)  
sempervirens 'Suffruticosa' (True Dwarf Boxwood)

**Callistemon**

citrinus (C. lanceolatus) (Lemon Bottlebrush)

**Camellia**

japonica 'Lady Laura'  
sasanqua (Sasanqua Camellia)

**Campanula**

persicifolia 'Blue'

**Catharanthus**

roseus 'Vinca Blue Pearl'

**Ceanothus\***

(Lilac)

**Cercis**

canadensis (Eastern Redbud)

**Chamaecyparis**

lawsoniana 'Ellwoodii' (Ellwood Cypress)  
obtusata (Hinoki False Cypress)  
pisifera 'Filifera Aurea' (Cypress)

**Chamaedorea**

elegans (Chamaedorea, Parlor Palm)

**Coreopsis**

verticillata 'Moonbeam'

**Cornus**

florida (White Flowering Dogwood)

**Cortaderia**

selloana (Pampas Grass)

**Cotoneaster**

apiculatus (Cranberry Cotoneaster)

dammeri 'Coral Beauty' (Bearberry Cotoneaster)

**Crataegus\***

(Hawthorn)

**Cupressus**

sempervirens (Italian Cypress)

**Cytisus**

scoparius 'Burkwoodii' (Scotch Broom)

**Deutzia**

x 'Magician'

**Dianthus**

spp. 'Baths Pink'  
deltooides 'Arctic Fire'

**Escallonia**

spp. 'Fradesii'  
langleyensis 'Apple Blossom'

**Euonymus**

fortunei 'Emerald 'N Gold'  
fortunei 'Gracilis'  
fortunei 'Sarcoxie'  
japonica 'Aureo-variegata' (Gold Spot Euonymus)  
japonica 'Grandifolia' (Big Leaf Euonymus)  
x kiautschovicus 'Manhattan'

**Forsythia**

x intermedia 'Lynwood Gold'

**Fothergillia**

gardenii 'Mt Airy'

**Fragaria**

x 'Pink Panda' (Sand Strawberry)

**Gardenia**

jasminoides 'August Beauty' (Cape-jasmine)  
'Radicans'

**Hemerocallis**

spp. 'Happy Returns' (Daylily)

**Hydrangea**

macrophylla 'Nikko Blue' (Bigleaf Hydrangea)

**Hypericum**

patulum 'Hidcote' (St. Johnswort)

**Ilex**

'Blue Girl' (Dwarf Holly)  
opaca (American Holly)  
cornuta 'Burfordii' (Burford Holly)  
cornuta 'Burfordii Nana' (Dwarf Burford Holly)  
crenata 'Convexa' (Japanese Holly)  
crenata 'Helleri' (Heller's Japanese Holly)  
hybrid 'Nellie R. Stevens'  
vomitoria 'Nana' (Dwarf Yaupon Holly)

**Juniperus**

chinesensis 'San Jose' (San Jose Juniper)  
chinesensis 'Sea Green' (Sea Green Juniper)  
conferta (Shore Juniper)  
horizontalis 'Plumosa' (Andorra Juniper)  
horizontalis 'Wiltonii' (Blue Rug Juniper)  
sabina 'Tamariscifolia' (Tam Juniper)

**Ligustrum**

ibolium 'Variegatum' (Ibodium Privet)  
japonicum (Japanese Privet)  
japonicum 'Rotundifolium' (Japanese Privet)  
japonicum 'Texanum' (Wax Leaf Privet, Texas Privet)  
senensis (Chinese Silver Privet)  
vicaryi (Vicary Golden Privet)

**Lonicera**

fragrantissima (Winter Honeysuckle)

**Loropetalum**

chinensis 'Burgundy'

**Lupinus**

spp. 'RH My Castle' (Lupine)

**Magnolia**  
soulangiana (Saucer Mahonia)

**Mahonia**  
bealei (Leatherleaf Mahonia)  
repens (Creeping Mahonia)

**Monarda**  
didyma 'Claire Grace'

**Nandina**  
spp. 'Compacta'

**Nerium**  
oleander (Oleander)

**Pachysandra**  
terminalis (Japanese Spurge)

**Philodendron**  
selloum (Split Leaf Philodendron)

**Photinia**  
fraseri (Fraser's Photinia)

**Picea**  
glauca 'Conica' (Dwarf Alberta Spruce)

**Pieris**  
japonica (Lily-of-the-Valley Shrub)

**Pinus**  
halepensis (Aleppo Pine)  
thunbergiana (Japanese Black Pine)

**Pittosporum**  
tobira 'Wheeler's Dwarf' (Tobira, Mock Orange)

**Podocarpus** spp.

**Potentilla**  
fruticosa 'Katherine Dykes' (Bush Cinquefoil)

**Prunus**  
cistena 'Dwarf Red-Leaf' (Dwarf Red-Leaf Plum)

**Pyracantha**  
spp. 'Mohave'  
coccinea 'Lalandei Monrovia' (Scarlet Firethorn)

**Quercus**  
acutissima (Sawtooth Oak)  
shumardii (Shumard Red Oak)

**Raphiolepis**  
indica 'Pink Lady' (India Hawthorn)

**Rhododendron**  
'Nova Zembla'

**Salvia**  
spp. splendens 'Lavender Vista'  
'May Knight'

**Santolina**  
virens (Green Lavender Cotton)

**Scabiosa**  
columbaria 'Butterfly Blue'  
columbaria 'Pink Mist'

**Schefflera\***  
actinophylla (Schefflera, Queensland Umbrella Tree)  
arboricola (Arboricola, Dwarf Schefflera)

**Spiraea**  
bumalda 'Goldflame'

**Syngia**  
meyeri 'Pallbin' (Lilac)  
vulgaris (Common Lilac)

**Taxus**  
cuspidata 'Intermedia' (Taxus, Yew)

**Thuja**  
occidentalis 'Pyramidalis' (American Arborvitae)  
occidentalis 'Globosa' (Globe Arborvitae)

**Trachelospermum\***  
jasminoides (Star Jasmine)

**Tsuga**

canadensis (Canada Hemlock)

**Viburnum**  
davidii (David Viburnum)

**Vinca**  
spp. 'Bowles Variety' (Periwinkle)

**Weigela**  
florida 'Pink Princess'

**Xylosma**  
congestum (spp. senticosum) (Shiny Xylosma)

*\*Species not yet approved by the state of California.*

**THE SPECIES LISTED BELOW HAVE BEEN TESTED AND RESULTS DO NOT WARRANT USE AT THIS TIME:**

**Achillea**  
spp. 'Coronation Gold' (Yarrow)

**Ageratum**  
spp. 'Hawaii Blue'

**Azalea**  
'Hino-crimson'  
'Snow'

**Calluna**  
spp. 'Forsy' (Heather)

**Celosia**  
spp. 'Castle Yellow'

**Chrysanthemum**  
spp. 'Legends'

**Clematis**  
x 'Duchess of Edinburgh'

**Dianthus**  
caryophyllus

**Digitalis**  
spp. 'Excelsior'

**Echinacea**  
spp. 'Magnus'  
pallida  
purpurea 'Leuchstern'

**Erica**  
mediterranea

**Euonymus**  
alata compacta

**Geranium**  
spp. 'Pinto Red'

**Hydrangea**  
arborescens 'Annabelle'

**Lantana**  
spp. 'Patriot Desert Sunset'

**Leucanthemum**  
maximum 'White Knight' (Shasta Daisy)

**Lilium**  
orientalis 'Mr. Ruud'

**Magnolia**  
stellata 'Pinkie'

**Papaver**  
orientale 'Pizzicato Mix' (Poppy)

**Penstemon**  
spp. 'Thorn'

**Petunia** spp.

**Potentilla**  
fruticosa 'Goldfinger'

**Prunus**  
glandulosa (Dwarf Flowering Almond)

### **Rhododendron**

spp. 'English Roseum'  
'Hino-Crsimson'  
'Snow'

### **Rudbeckia**

spp. (Coneflower)  
'Goldsturm'

### **Sarcococca**

hookeriana humilis

### **Scabiosa**

spp. 'Pink Carnival'

### **Sedum**

spectabile 'Autumn Joy'  
brevifolium (Stonecrop)

### **Tagetes**

erecta (Marigold)

### **Vinca**

minor (Dwarf Periwinkle)

### **Zinnia spp.**

## **GIVES PREEMERGENCE CONTROL OF THE FOLLOWING WEEDS:**

### **GRASSY WEEDS**

Annual bluegrass	( <i>Poa annua</i> )
Barnyardgrass	( <i>Echinochloa crusgalli</i> )
Crabgrass	( <i>Digitaria sanguinalis</i> )
Sprangletop	( <i>Leptochloa uninervia</i> )

### **BROADLEAVED WEEDS**

Bittercress	( <i>Cardamine pensylvanica</i> )
Common Chickweed	( <i>Stellaria media</i> )
Cudweed	( <i>Gnaphalium obtusifolium</i> )
Dandelion	( <i>Taraxacum officinale</i> )
Eclipta	( <i>Eclipta prostrata</i> )
Fireweed	( <i>Epilobium angustifolium</i> )
Fleabane	( <i>Erigeron strigosus</i> )
Groundsel	( <i>Senecio vulgaris</i> )
Marestail	( <i>Gonyza canadensis</i> )
Oxalis	( <i>Oxalis stricta</i> )
Pearlwort	( <i>Sagina procumbens</i> )
Pepperweed	( <i>Lepidium virginicum</i> )
Pigweed	( <i>Amaranthus retroflexus</i> )
Shepherdspurse	( <i>Capsella bursa-pastoris</i> )
Sowthistle	( <i>Sonchus arvensis</i> )
Spurge, Garden	( <i>Euphorbia hirta</i> )
Spurge, Prostrate	( <i>Euphorbia supina</i> )
Spurge, Spotted	( <i>Euphorbia maculata</i> )

## **APPLICATION RECOMMENDATIONS**

- Apply to dry foliage only
- Water in immediately to wash the particles off the plant foliage
- Apply 2.3 lbs. product per 1,000 sq. ft.
- Repeat application at 3-month intervals

Before each application, remove any existing weeds. Apply product with a properly calibrated rotary spreader that will assure accurate, even particle distribution. Product can also be applied by aerial equipment.

Apply to **dry foliage** only. To test for moisture, rub hands over plant foliage. **Water in** with 1/2-1 inch immediately after application to wash the particles off the plant foliage and to activate the herbicide.

For season-long weed control, repeat applications are required at approximately 3-month intervals during the growing season.

**The maximum application is 6 lbs/season at 2 lbs ai/acre/ application.**

## **CONTAINER GROWN ORNAMENTALS**

**POTTED LINERS:** Remove established weeds from liners prior to transplanting into containers. Following planting and prior to chemical application, irrigate with approximately 1/2 inch of water to firm potting media around root systems. After two regular waterings and foliage has dried, broadcast product evenly over the container growing area. Water in with 1/2 to 1 inch immediately after application.

**BARE-ROOT LINERS:** Remove established weeds before herbicide application. Broadcast product evenly over the container growing area 3-4 weeks after potting. Water in with 1/2 to 1 inch immediately after application.

## **INDIVIDUAL CONTAINER APPLICATIONS**

For situations where broadcast application of the product is not appropriate, apply the product uniformly over the entire container surface at the rates in the following table (1 gm product/sq. ft):

Container Size	Surface Area (Sq. Ft.)	Grams of Product/ Container	Volume of Product/ Container
1 gal. (6" diam.)	0.20	0.20	1/8 tsp.
2 gal. (9" diam.)	0.44	0.44	1/4 tsp.
3 gal. (10" diam.)	0.55	0.55	1/4 tsp.*
5 gal. (12" diam.)	0.79	0.79	1/4 tsp.**
15 gal. (17" diam.)	1.60	1.60	1/2 tsp.

\*rounded  
\*\*heaping

## **FIELD GROWN ORNAMENTALS**

Apply immediately after planting or prior to weed seed germination following complete weed removal. Before application the soil should be loosened by light cultivation.

Water the product in with 1/2 to 1 inch immediately after application.

## **PRECAUTIONS/LIMITATIONS**

- Do not apply to moist foliage.
- Do not apply while plants are producing a new flush of spring growth. Tender, newly forming leaves are especially sensitive and may be injured by the herbicide.
- Do not apply when extreme cold stress (35°F or lower) is expected.
- Do not apply product in enclosed greenhouse or polyhouse structures as ornamental injury may result. Product should not be applied within two (2) weeks prior to enclosure in greenhouse type structure or within two (2) weeks prior to shipping.
- Do not apply to plant types whose leaves channel the herbicide granules to the leaf base (such as Yucca spp. and Dracaena spp.).
- Do not apply in wind conditions that affect distribution of the product.
- If metallic surfaces (nursery equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 mph) due to its staining potential.

- *Not* for use on vegetables. Do not use on fruit trees that will bear fruit within one year of application.
- Users who wish to use OH2® Ornamental Herbicide on ornamental species or non-bearing fruit trees *not listed on this label may determine suitability for such uses by making trial application of this product at a recommended rate to a small number of plants.* Prior to using this product on a large number of plants, the trial treatment should be observed for signs of herbicidal injury for 3-6 months to determine if the treatment is safe to the target plant species.
- Non-bearing fruit trees such as citrus (oranges, grapefruit, etc.), pome fruits (apples, pears, etc.) and stone fruits (cherries, peaches, etc.) are plants which will not bear fruit for at least one year following application.
- The user assumes responsibility for any crop damage or other liability from the use of this product on ornamental plant species not recommended on this label.
- Not recommended for use on bedding plants, liner production beds, flowers, ground covers, vegetables, and turf.
- Do *not* incorporate product into soil or potting media nor apply product directly to bare roots.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## DISCLAIMER AND LIMITATION OF LIABILITY

### IMPORTANT NOTICE FROM EVERRIS NA INC. ("Everris").

#### PLEASE READ BEFORE USE.

To the extent consistent with applicable law, user or buyer accepts the conditions, disclaimer of warranties and limitations of liability. Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once for full refund.

**CONDITIONS:** The directions for use of this product are believed to be adequate and the user or buyer must always follow the label directions carefully and exercise judgment and caution when using this product under their growing conditions. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness, unsatisfactory or substandard results or other unintended consequences may result because of such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Everris. All such risks shall be assumed by the user or buyer.

**WARRANTY:** This product corresponds to all claims and descriptions set forth on the label and is reasonably fit for the purposes set forth in the directions for use on the label when used in accordance with those directions. This warranty is subject to the conditions and limitations stated herein. Everris recognizes that the rights and remedies of the user or buyer are subject to the provisions of the applicable law, but makes no other warranties or representations, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Everris is authorized to make any warranties beyond those contained herein or to modify the warranties contained therein. Subject to the user's or buyer's rights and remedies under the applicable law, Everris disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** Subject to the user's or buyer's rights and remedies under the applicable law, the exclusive remedy of the user or buyer and the liability of Everris or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid by the user or buyer for the quantity of this product involved or, at Everris' election, the replacement of the product. To the extent consistent with applicable law, Everris must have prompt notice of any claim so that a timely investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify Everris of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

Manufactured for:  
Everris NA Inc.  
PO Box 3310  
Dublin, OH 43016  
1-800-492-8255

OH2® is a Registered Trademark of Everris International BV for its brand of Herbicide products.

Product of U.S.A.