



ferti·lome





# Triple Action Plus Insecticide • Fungicide • Miticide Ready To Use Ready To Use

 For Control Of Insects, Diseases & Mites On Vegetables, Fruits. Nuts, Herbs, Spices, Roses, Flowers & Shrubs

# **KEEP OUT OF REACH** OF CHILDREN

SEE BACK PANEL AND BOOKLET FOR FIRST AID AND ADDITIONAL

PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENTS:

Pyrethrins 0.02% Clarified Hydrophobic Extract of Neem Oil 0.90%

Net Contents 32 FL. OZ. (946ML)

	FIRST AID
If On Skin:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If In Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes then continue rinsing the eye.     Call a poison control center or doctor for treatment advice.
if Swallowed:	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center odctor. Do not give anything by mo
If Inhaled:	Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth.     Call a Poison Control Center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Chem Tel Service 1 and the vent of a medical or chemical emergency, contact North America 1-800-255-3924 or Worldwide Intl. + 01-813-248-0585	

Manufactured By:

230 FM 87 • BONHAM, TEXAS 75418 EPA Reg. No. 70051-74-7401 • EPA Est. No. 7401-TX-001 Visit Us At: www.fertilome.com 25M-2-0113 RTUH





# PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacc. Remove and wash contaminated clothing before reuse. Wear: Long-sleeved shirt and long pants, socks, shoes, gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

**BEE HAZARD:** This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

#### PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### GENERAL INFORMATION

SHAKE WELL BEFORE AND DURING USE. Multi-purpose insecticide/ fungicide/niticide. Kills eggs, laval and adult stages of insects. Prevents Fungal attack of plants' tissues. Not for use in food handling areas. Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment. When using in conjunction with beneficial Insects, conduct a small trial to assure compatibility before using on a large scale. As with other products that contain oils, take care in timing applications to early morning/late afternoon/early evening to minimize the potential for leaf burn. Use with care on plants with tender tissue. Do not apply to known spray sensitive plant species such as: Impatiens Flowers, Fuchsia Flowers, Hibiscus Flowers, some Rose Flowers, Ornamental Olive Trees, Red Maples, some Carnation varieties or similar plants without prior testing. Keep unprotected persons out of treated areas until soravs have dried.

#### **FUNGICIDE USE**

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES. HOUSEPLANTS, FLOWERS AND SHRUBS, Application Instructions: fertilome® TRIPLE ACTION PLUS RTU is an effective Fungicide for the prevention and control of various Fungal Diseases including Powdery Mildew, Black Spot, Downy Mildew, Anthracnose, Rust, Leaf Spot, Botrytis, Needle Rust, Scab and Flower, Twig and Tip Blight, As a preventative, apply fertilome® TRIPLE ACTION PLUS RTU on a 7 to 14 day schedule until the potential for Disease development is no longer present. To control Disease already present. apply fertilome® TRIPLE ACTION PLUS RTU on a 7 day schedule until Disease pressure is eliminated. Then continue spraying on a 14 day schedule to prevent the Disease from reoccurring. To prevent Rust Leaf Spot Diseases, Anthracnose and Scab, begin applications at the first sign of Spring budding. To prevent Powdery Mildew, apply in mid-Summer or when Disease is first detected. Continue applications until Disease pressure no longer exists. Thoroughly spray all plant leaf surfaces (including undersides of leaves) until foliage is uniformly wet, but not dripping. Excessive application is unnecessarv and should be avoided.

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS AND SHRUBS. Application Instructions:
Apply fertilome® TRIPLE ACTION PLUS RTU at the first sign of Insects/
Mites. For control of the following: Aphids, Spider Mites, Scale,
Mealybugs, Whiteflies, Beetles (including Japanese), Loopers,
Leafminers, Leafrollers, Armyworms, Webworms, Weevils, Tent
Caterpillars, Fungus Gnats, Leafhoppers, Cabbageworms, Squash Bugs,
Squash Vine Borer, Stink Bugs, Thrips, Tomato Hornworm, Lace Bugs,
Boxelder Bugs, Grasshoppers, Crickets, Ants, Fleas, Ticks, Earwigs,
Cockroaches, Mosquitos, Chinch Bugs, Chiggers and many other Insect
pests. fertilome® TRIPLE ACTION PLUS RTU is most effective when
applied on a 7 to 14 day schedule. For heavy Insect populations, spray
on a 7 day schedule. Thoroughly spray all plant surfaces (including
undersides of leaves) until completely wet, but not dripping. Excessive
application is unnecessary and should be avoided.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticide Storage: Store upright at room temperature. Avoid exposure to extreme temperatures. Pesticide Disposal and Container Handling: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Voluntary Purchasing Groups, Inc., warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the Directions For Use. Timing and method of application, weather, watering practices, nature of soil, the Insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.





230 FM 87 • BONHAM, TEXAS 75418 EPA Reg. No. 70051-74-7401 EPA Est. No 7401-TX-001 Visit Us At: www.fertilome.com



Get the free mobile app at http://gettag.mobi

