

### **General Information**

**NUE Copper 5%** is manufactured specifically as a liquid foliar fertilizer to be used to help prevent and correct copper deficiencies that may limit crop growth and yields. **NUE Copper 5%** is a supplement to an adequate and balanced soil fertility program.

It is important to monitor crop needs and responses by maintaining an adequate tissue-testing program. When unfamiliar with the effects of foliar applications of minerals it is recommended to perform small-scale field trials.

### **Directions for Use**

#### **Mixing Directions**

- 1. Be sure spray tank is thoroughly cleaned.
- 2. Fill 1/2 full with clean water and start agitation.
- 3. Add other nutrients if compatible.
- 4. Add NUE Copper 5% and continue agitation.
- 5. Adjust pH if necessary.
- 6. Add pesticides if compatible.
- 7. Fill tank and agitate thoroughly.

Compatibility - NUE Copper 5% is compatible with most pesticides.

For maximum nutritional results with tank mixes containing NUE Copper 5% adjust the final pH to 5.0 -7.0. Use NUE Cal-8 raise to pH of the final tank mix, if necessary. When tank mixing NUE Copper 5% with pesticides requiring specific pH ranges, always follow pesticide label directions.

Always check compatibility with a jar test before mixing with other fertilizers (i.e. phosphorus). If any precipitation occurs, it may be necessary to add a compatibility agent. Consult your Bio-Gro dealer for more information.

Dilution and Spray Volumes - Always use enough water for uniform application and to prevent possible plant tissue damage to the intended crop from concentrated droplets. It is recommended to use a dilution of at least 20 parts of water to 1 part of NUE Copper 5%.

BG Rev 03-01-2007

Rorn

# 5-0-0

# Guaranteed Analysis

Derived from: Copper Nitrate

# Keep out of Reach of Children CAUTION

Harmful if swallowed, inhaled or absorbed through skin. May cause eye and skin irritation. In case of contact, immediately flush eyes and/or wash skin with plenty of fresh water for at least 15 minutes.

Do not contaminate feed or food items.

Use caution in application to assure coverage on intended treatment areas only.

#### **CONDITIONS OF SALE**

Seller and/or manufacturer warrants only that this product consists of the ingredients specified on this label. No one is hereby authorized to make any warranty and/or guarantee concerning the use of this product. Being that the use, timing, placement, rate, application and other conditions are beyond the control of the seller and/or manufacturer, the seller's and/or manufacturer's liability for handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

### Manufactured by:

Bio-Gro, Inc. 681 Glade Road Mabton, WA 98935 Tel: 509-894-4110



## **Suggestions for Use**

Optimum rate of application will vary between fields depending on soil and/or climatic conditions, stage of crop growth and size of plants.

Application rates are best determined by soil and plant analysis recommendations.

### **Foliar Application Rates:**

**Field Crops:** Apply 16 to 32 ounces/acre as a directed foliar spray during periods of rapid growth or nutritional demand once plants have reached 4-8 inches in height. NUE Copper 5% may be applied at 10 to 14 day intervals.

**Vegetable Crops:** Foliar rate is 16 to 32 ounces/acre. Apply NUE Copper 5% as needed throughout vegetative development and/or during rapid fruit growth.

**Tree and Vine Crops**: Apply 16 to 32 ounces/acre during growing season. Applications during verasion should be limited to 16 ounces /acre and used with appropriate adjuvant unless experience dictates otherwise. Repeat applications may be made in 10 to 21 day intervals.

**Berry Crops:** Apply 8 to 32 ounces/acre of NUE Copper 5% after active vegetative growth begins. Applications may be repeated in 7 to 14 day intervals.

Foliar fertilization can be an effective remedy for diagnosed plant deficiencies of secondary nutrients and micronutrients, by may cause plant damage if applied at more then recommended rates or concentrations. Use of this product is recommended only as a supplement to a regular fertilization program and only on plants with confirmed secondary nutrient or micronutrient deficiencies.

250 gallons (946 liter)
9.74 lbs/gallon (1.167 kg/liter)

**MADE IN USA** 

NUE Copper 5%