TECH-SPRAY®

Post Harvest Blend

0-12-5

TECH-SPRAY Post-Harvest Blend supplies the guaranteed nutrients in a water soluble form for use as a nutritional spray. It is designed to be combined with TECH-FLO® products but also may be applied alone.

GUARANTEED ANALYSIS	
Available Phosphate (P ₂ O ₅)	. 12.0%
Soluble Potash (K ₂ O)	5.0%
Sulfur (S)	0.6%
Cobalt (Co)	0.18%
Copper (Cu)	0.68%
Molybdenum (Mo)	0.1%
Zinc (Zn)	0.58%
Derived from: Phosphoric Acid, Po	tassium
Phosphate, Cobalt Sulfate, Copper	Sulfate,
Ammonium Molybdate, Zinc Sulfate.	

CAUTION

KEEP OUT OF REACH OF CHILDREN HARMFUL IF SWALLOWED

CAUTION: Causes skin/irritation. Avoid contact with skin, eyes, open cuts or sores. In case of contact, flush with water. Get medical attention for eyes.

WARNING: This product contains Molybdenum (Mo); crops high in molybdenum are toxic to grazing animals (ruminants). Use only according to manufacturer's directions.

NOTICE: Do not store this product at temperatures below 40° F as precipitates may form that are difficult to redissolve. Do not expose to temperatures below 32° F.

UTRIENT TECHNOLOGIES, INC. DINUBA. CA 93618

MADE IN THE U.S.A.

NET CONTENTS: 2.5 GAL. (9.46 LITERS) 10.40 LBS./GAL.

TECH-SPRAY® Post Harvest Blend - SPECIMEN LABEL (cont.)

GENERAL RECOMMENDATIONS

AS A NUTRITIONAL SPRAY:

DILUTION: Apply the suggested rates of TECH-SPRAY® in the following amounts of water per acre. Minimum dilution rate: 1 qt. per 5 gallons water per suggested rate.

 COMPATIBILITY: This product has been designed to be compatible with most commonly used insecticides, fungicides and acaricides. However, it is not feasible to test all possible combinations of pesticides with which it may be mixed. Do not mix with lime, fixed copper fungicides, or TPTH. A compatibility test is recommended for combinations that have not been used previously. Consult your NUTRI-TECH representative for detailed compatibility information.

SUGGESTED USES (RATES PER ACRE):

- Alfalfa, Hay, & Other Forage Crops Apply 1 pt.- 2 qts. over spring growth, repeat after each cutting.
- Almonds Apply 1 pt.- 4 qts. at 2-3 week intervals starting prebloom
- **Apples, Pears** Apply 1 pt.- 4 qts. at 2-4 week intervals starting pre-bloom.
- **Asparagus** Apply 1 pt.- 2 qts. after start of new fern growth, repeat every 2-4 weeks as necessary.
- Barley, Oats, Rice, Wheat & Other Small Grains Apply 1 pt.-2 qts. at tillering, repeat at early boot if necessary.
- Beans, Peas, & related Legumes Apply 1 pt.- 2 qts. at early bud, repeat after bloom if necessary.
- Cabbage, Cauliflower, Broccoli & Other Cole Crops Apply 1 pt.- 2 qts. as needed after thinning or transplant.
- Carrots, Onions, Radishes, Beets & Other Vegetable Root Crops - Apply 1 pt. - 2 qts. when top growth is 4-8" high, repeat as necessary.
- Citrus, Avocados Apply 1 pt.- 4 qts. in spring, summer, and fall sprays.
- **Coffee** Apply 1 pt.- 2 qts. starting pre-bud stage, repeat as needed throughout growing cycle.
- Corn, Field & Sweet Apply 1 pt.- 2 qts. when 10-20" high, repeat in 2-4 weeks as necessary.
- Cotton Apply 1 pt.- 2 qts. at first square, repeat in 2-4 weeks as needed.
- **Cranberries** Apply 1 pt.- 2 qts. as needed, repeat 3-4 weeks as needed.
- Cucumbers, Melons, Squash & Other Cucurbits Apply 1 pt.-2 qts. pre-bloom, repeat in 2-4 weeks as necessary.
- Figs Apply 1 pt.- 2 qts. as needed, repeat 3-4 weeks as necessary.
- **Grapes** Apply 1 pt.- 2 qts. dormant or as needed pre-bloom to post-bloom.
- **Hops** Apply 1 pt. 2 qts. as needed in cover sprays.
- **Kiwi** Apply 1 pt.-2 qts. as needed, repeat in 3-4 weeks as necessary.
- **Lettuce, Celery, Spinach & Other Green Leafy Vegetables -** Apply 1 pt. 2 qts. as needed after thinning and transplant.
- Mangos, Guavas, & Other Tropical Fruits Apply 1 pt. 4 qts. as needed; repeat every 3-4 weeks as necessary.

Mint - Apply 1 pt. - 3 qts. over new growth, repeat as needed.

- Olives Apply 1 pt.- 2 qts. pre-bloom or post-bloom, repeat 3-4 weeks as necessary. For crops that require thinning, apply 1-2 weeks after thinning, repeat 3-4 weeks as necessary.
- **Ornamentals, Trees & Shrubs** Apply 1 pt.- 2 qts. at 4-6 week intervals as needed.
- Peaches, Plums, Cherries, & Other Stone Fruits Apply 1 pt.-2 qts. pre-bloom and at 3-4 week intervals as needed post-bloom
- **Peanuts** Apply 1 pt.- 2 qts. at early bloom and just prior to pegging.
- Pistachios, Pecans, & Other Deciduous Nut Trees Apply 1 pt.-4 qts. at 3-4 week intervals starting at bud break.
- **Pomegranates** Apply 1 pt.- 2 qts. as needed, repeat in 2-4 weeks as necessary.
- **Potatoes** Apply 1 pt.- 2 qts. at tuber initiation, repeat in 2-4 weeks as necessary.
- Raspberries, Blackberries, Blueberries & Other Small Fruit Apply 1 pt.- 2 qts. pre-bloom, repeat in 2-4 weeks as necessary.
- **Soybeans, Seed Crops** Apply 1 pt.- 2 qts. pre-bloom, repeat in 2-4 weeks as necessary.
- Sugarbeets Apply 1 pt.- 2 qts. at 4th-6th leaf, repeat in 2-4 weeks as necessary.
- **Strawberries** Apply 1 pt.- 2 qts. pre-bloom, repeat in 2-4 weeks as necessary.
- **Tomatoes, Peppers, Eggplant & Okra** Apply 1 pt.- 2 qts. prebloom, repeat in 2-4 weeks as necessary.
- **Tropical Flowers & Foliage Plants -** Apply 1 pt.-2 quarts as needed, repeat in 3-4 weeks as necessary.
- Turf (Farms) Apply 1 pt. 2 qts. at 3-6 week intervals as needed.
- Walnuts Apply 1 pt. 4 qts. in blight sprays or as necessary.

NOTE: For crops not listed on this label, consult your NUTRITECH representative.



NOTICE: Information regarding the contents and levels of metals in this product is available on the internet at: http://aapfco.org/metals.htm