TruPhos Platinum 5-18-2

Guaranteed Analysis

Total Nitrogen (N) Copper (Cu) Available Phosphoric Acid (P205) 18% Iron (Fe) 0.4% Soluble Potash (K2O) 2% 0.04% Manganese (Mn) Magnesium (Mg) 0.4% Molybdenum (Mo) 0.05% Boron (B) 0.1% 0.8% Zinc (Zn) Cobalt (Co) 0.05%

An OrthoPhosphate True Solution with Nitrogen, Potassium and **Micronutrients**

Derived from orthophosphate, potassium carbonate, urea, sulfates of manganese, copper, iron and magnesium, zinc oxide, sodium molybdate, cobalt chloride and boric

For use as a fertilizer in foliar and fertigation applications to supplement phosphorus, potassium, nitrogen, zinc, magnesium, iron, magganese, molybdenum, boron and copper levels on agricultural crops.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground sprays minimum of 5 gallons of water/acre; Dilute sprays minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- TruPhos Platinum is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

Crop	Foliar Rates*	Fertigation Rates*
Forage Crops: Alfalfa, Clover, Hay Crops		The number and frequency of
Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane		applications determine
Legumes: Dry bean, Lentil, Peanut, Pea, Soybean		application rates.
Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga,		
Sugar Beet, Sweet Potato, Turnip		Apply no more than
Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops,		1 – 2 quarts/acre/application.
Lettuce, Parsley, Spinach		
Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing),	2 – 3 pints/acre/	
Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato	application	Always apply toward the end of
Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry		an irrigation cycle.
Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach,		
Pear, Persimmon, Plum, Pomegranate, Prune, Quince		
Other Fruit: Citrus (all), Fig, Kiwi, Melons		
Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut		
Grapes: Currant, Raisin, Table, Wine		
Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower		

*Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental Net Contents: conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe 2x2.5 Gal. (2x9.46 L) deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit use or nannung or this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the 52.90 lbs. (24.00 kg) use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: This product contains Boron (B). The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

CAUTION: This product contains Molybdenum (Mo). The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

Caution: KEEP OUT OF REACH OF CHILDREN



Net Weight:

Net Weight/Gallon:

10.58 lbs. (4.80 kg)

Batch No.: 180215

Rev: 011216





May cause an allergic skin reaction. Causes serious eye. irritation. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause cancer. May damage fertility or the unborn child. Contains 21.5-25.9% of ingredients of unknown toxicity (inhalation). Contains 3.5-6.5% of ingredients of unknown toxicity (dermal).

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid breathing mist, spray, Contaminated work clothing must not be allowed out of the workplace. Wear eye protection/face protection. In case of inadequate ventilation wear respiratory protection (NIOSH approved air-purifying respirator with an appropriate cartridge.

If on skin: Wash with plenty of water. If inhaled: If breathing is difficult, remove person to fresh air and keep comfortable for breathing. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If exposed or concerned: Get medical advice/attention. Specific treatment (see supplemental first aid instruction on this label). If experiencing respiratory symptoms: Call a poison center/doctor. Wash contaminated clothing before reuse

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations.

