



Moletox[®] Mole Killer

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only for the sites, pests and application methods specified on this label.

Read this Label:

Read this entire label and follow all use directions and precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Follow all application directions and **Use Restrictions** on this label. Apply bait only in moles' underground runways. Keep pets out of treated areas and isolate treated areas using existing fencing, if present.
3. Dispose of product container and unused or spoiled bait as specified on this label.

USE RESTRICTIONS: This product may be used only to control eastern moles (*Scalopus aquaticus*), star-nosed moles (*Condylura cristata*), hairy-tailed moles (*Parascalops breweri*), coast moles (*Scapanus orarius*), broad-footed moles (*S. latimanus*) and/or Townsend moles (*S. townsendii*) on lawns, turf areas, golf courses and other non-food grassy areas. This product may only be applied at the use sites and by the application methods indicated on this label. Bait must be applied directly into moles' burrow systems (main underground tunnel or subsurface runways). Do not place bait on the ground surface. *Not for use against the star-nosed mole in North Carolina.

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of surface ridges in the turf or by a series of conical mounds of earth pushed up from deep burrows. Conical mounds may betray the location of main underground runways.

BAITING: Prior to treatment, determine which burrow systems are active. Using a round-ended wood rod, such as a broomstick, probe ground in vicinity of conical mounds or surface ridges until main underground runway is detected by a sudden decrease in resistance against the probe. Mark open burrow systems and revisit them 2-3 days later. For deeper tunnels, open system with a shovel and leave open. Burrow systems that are active are likely to be plugged by moles.

Treat only active burrow systems. To treat a burrow system, locate main runways by probing. Remove the cap from the tip of the applicator syringe and turn dosing ring to mark indicating half ounce. Inject half an ounce of bait (one-sixth of the contents of one syringe) through probe or shoveled opened hole at each bait placement location. Make six (6) such placements per burrow system. Cover holes made to enter tunnel with sod and soil, but do not allow dirt to sift in to foul or cover bait.

FOLLOW-UP: Check effects of treatment 4-5 days after bait application by opening and marking burrows in the manner used to check for initial activity. Return 2-3 days later and retreat all active burrows. Maximum application rate is 1 syringe/10,000 ft.²

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets. Do not store used or partially used syringes. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Non-refillable container; do not reuse or refill this container. Offer for recycling, if available; otherwise, dispose of empty container in a sanitary landfill; or by incineration.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. Anyone handling baits or retrieving carcasses must wear gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If Swallowed: • Call a poison center, doctor, or National Pesticide Information Center (1-800-858-7378) immediately for treatment advice.
• Have a person sip a glass of water if able to swallow.
• Do not induce vomiting unless told to do so by the poison center or doctor.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-800-222-1222 at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the anticoagulant Warfarin. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁ orally or intramuscularly.

Environmental Hazards

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

WARRANTY: To the extent consistent with applicable law, Manufacturer and Seller make no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Notice: This product contains a chemical known to the state of California to cause birth defects or other reproductive harm.

Distributed by **BONIDE PRODUCTS, INC.**
6301 SUTLIFF ROAD • ORISKANY, NY 13424



0 37321 00693 0



Moletox[®] Mole Killer

- Treats an entire burrow system
- Poison Bait with syringe applicator
- Use on lawns, turf areas, golf courses & other non-food grassy areas



Moletox[®] Mole Killer

This product may only be used to control moles in manual, below-ground applications

Gel bait designed to mimic moles natural food source



poison bait with syringe applicator



Keep Out Of Reach Of Children
CAUTION (See Back Panel for Additional Precautionary Statements)
Net Weight 3 FL OZ

ACTIVE INGREDIENT:	
Warfarin . . . (CAS No. 81-81-2)	0.025%
OTHER INGREDIENTS	99.975%
TOTAL	100.000%
EPA Est. No. 4-NY-1	EPA Reg. No. 72500-2-4

FIRST AID: HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If Swallowed: Call a poison control center, doctor, or National Pesticide Information Center (1-800-858-7378) immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian or 1-800-222-1222 at once.

NOTE TO PHYSICIAN OR VETERINARIAN: Contains the anticoagulant Warfarin. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁ orally or intramuscularly.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Do not store used or partially used syringes. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Non-refillable container; do not reuse or refill this container. Offer for recycling, if available; otherwise, dispose of empty container in a sanitary landfill; or by incineration.



This product may only be used to control moles in manual, below-ground applications.

ACTIVE INGREDIENT:

Warfarin (CAS No. 81-81-2) 0.025%

INERT INGREDIENTS: 99.975%

TOTAL: 100.000%

Dist. by Bonide Products, Inc., 6301 Sutliff Road, Oriskany, NY 13424

EPA Reg. No. 72500-2-4

EPA Est. No. 4-NY-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Read complete label and instructions on packaging before using this product.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. Anyone handling baits or retrieving carcasses must wear gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.