

19.000"

30.000"

LESCO® T-Storm™ 2G Fungicide

LESCO®

T-Storm™ 2G Fungicide

LESCO® T-Storm™ 2G Fungicide contains thiophanate-methyl, the active ingredient used in Cleary 3336™.

Systemic Fungicide for Turf and Ornamentals

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

Table with 2 columns: Ingredient Name and Percentage. Includes Thiophanate-methyl (2.08%), Other Ingredients (97.92%), and TOTAL (100.00%).

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

See the back of this bag for additional PRECAUTIONARY STATEMENTS

FIRST AID

- If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (877) 325-1840 for emergency medical treatment information.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. For Medical Emergencies Only, Call (877) 325-1840.



#10163032 NET WEIGHT: 30 Lbs. (13.60 Kg)

Distributed By: LESCO, Inc. 1385 East 36th Street Cleveland, OH 44114-4114

EPA Reg. No. 228-631-10404 EPA Est. No. 8378-IN-001



LESCO® T-Storm™ 2G Fungicide

8 Thomasville, NC

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION/PRECAUCIÓN Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. PERSONAL PROTECTIVE EQUIPMENT (PPE) Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart. Applicators and other handlers must wear: Long-sleeved shirt and long pants Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber User Safety Requirements Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements specific to your State or Tribe, consult the agency responsible for pesticides.

AGRICULTURAL USE REQUIREMENTS Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants Chemical-resistant gloves made of any waterproof material Chemical-resistant footwear plus socks Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Turf and Landscape Uses: Do not enter or allow entry into treated areas until granules and dust have settled unless wearing personal protective equipment listed on this label.

COMMERCIAL APPLICATIONS

GENERAL INFORMATION This product is a broad-spectrum fungicide for use on a wide variety of turf and ornamental diseases. Apply this product using ground-based, hand-operated (push) or mechanical (power) equipment.

USE PRECAUTIONS For use only by certified applicators or those under their immediate supervision. Not for use on turf being grown for sale or other commercial use as seed. Do not apply this product through any type of irrigation system. Do not use on turf being grown for sale or other commercial use as seed. Do not apply this product by hand or with hand-held rotary sprayer (e.g., belly grinder). Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Use only properly calibrated application equipment when applying this product. Thoroughly clean all application equipment after each use. For application schedules appropriate to your area, contact your local State Extension Service for instructions.

RESISTANCE MANAGEMENT In order to prevent the development of tolerant strains of fungi, use this product with other fungicides that use different modes of action. Do not use this product with products containing the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can contribute to the development of disease tolerance. This product can remain useful for disease control as long as the specified precautions are followed. A tolerant strain of fungi is most likely present if a treatment of this product is found to be ineffective and you should promptly contact your State Agricultural Experiment Station or State Cooperative Extension Service for disease identification and advice on other suitable fungicide or disease control strategies.

MAXIMUM APPLICATION RATES AND MINIMUM RETREATMENT INTERVALS Refer to the following table for the maximum treatment and seasonal application rates allowed based on the use site being applied to.

Table with 5 columns: Use Site, Pre-Treatment Application Rate, Maximum Per-Season Application Rate, Maximum Per-Season lbs. per 1,000 sq. ft., Minimum Retreatment Interval. Includes Residential or Public Areas, Golf Course Agrons, Greens and Tees, Golf Course Fairways, and Florida Golf Courses.

1 Maximum of four (4) applications per year. 2 Maximum of one (1) application per year.

COMMERCIAL TURF APPLICATIONS Use this product on all cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures in the following sites:

- Commercial turf
- Residential turf
- Public turf (parks, athletic fields, schools, day care centers, etc.)
- Golf courses (greens, tees, fairways and aprons)

When used in accordance with the instructions in this label, this product is not phytotoxic to any of the above-mentioned grasses.

Application Instructions Apply under dry or moist leaf/soil conditions. For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 to 0.25 inch of water. Treat aprons when treating golf greens.

Refer to the following table for specific rates and timing depending on the disease being treated. When treating under severe disease pressure, use the highest specified rate listed below.

Table with 3 columns: Disease(s), Disease Specific Instructions, Application Rate (lbs./1,000 sq. ft.). Includes Dollar Spot, Large Brown Patch, Acaochyia Leaf Blight, Copper Spot, Fusarium Patch, Zoysia Patch, Pink Snow Mold, Anthracnose, Iolar, and Anthracnose, basal.

13218000 RV051414[1]

Table with 3 columns: Disease(s), Disease Specific Instructions, Application Rate (lbs./1,000 sq. ft.). Includes Bermudagrass Decline, Take-All-Patch, Cool Season Brown Patch, Necrotic Ring Spot, Spring Dead Spot, Coprinus Snow Mold, Fusarium Blight, Gray Leaf Spot (Blotch), Leaf Spot, Leaf, Crown, and Root Diseases, Stripe Smut, Summer Patch, Bentgrass Dead Spot.

COMMERCIAL HORTICULTURAL APPLICATIONS This product is a broad-spectrum systemic fungicide for control of soilborne diseases affecting bulbs, corms, rhizomes, roots, stems and stems caused by:

- Botrytis, Gloecladium, Ramularia, Thielaviopsis, Cylindrocladium, Myrothecium, Rhizoctonia, Fusarium, Penicillium, Sclerotinia

Use this product on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Intergreenhouse, and Landscape sites.

When using this product for commercial horticultural applications, be sure to follow the guidelines in the "General Information" section above.

NOTE: Because this product does not control Pythium or Phytophthora, rotating with metolachlor, etridiazole, propanil, or other fungicides is recommended.

USE PRECAUTIONS Do not use this product on Swedish Ivy (Nepenthes exaltata) and Easter Cactus (Hatiora gaertneri). Do not apply to home orchards/fruit trees after fruit set. Do not use on fruit or nuts from treated ornamentals for food or feed. For ground and drench applications, do not exceed 300 pounds active ingredient from all thiophanate-methyl containing products per acre per crop season. For cut flowers, the maximum application rate is 0.5 pound active ingredient per acre per application. Because it is impossible to test this product on all species and cultivars, test in a preliminary trial on a small scale before full treatment is applied to any plant type not shown on this label. When conducting the trial, wait 5 to 7 days after treatment to evaluate results.

Application Instructions Apply this product as a surface, fly-on, broadcast or over-the-top application or media incorporated.

Surface and Broadcast Applications to established plants: Apply 8 lbs. of this product/1,000 sq. ft. Apply under dry or moist leaf/soil conditions. For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 to 0.25 inch of water.

Media Incorporation: Mix at a rate of 2 to 3 lbs./cubic yard before seeding or transplanting of desired crop. If not activated by the application of moisture, stockpile media incorporated with this product until needed for later use.

For Preventive Control: Prior to planting, apply as a media incorporation using the rate listed above. After seeding or transplanting propagation beds, containers, pots, or nursery or landscape beds, apply as a broadcast, fly-on, surface or over-the-top application using the "Surface and Broadcast Applications" instructions above.

For Treatment: When disease first appears after seeding or transplanting, apply using the "Surface and Broadcast Applications" instructions above. To maintain protection, make repeat applications every 21 to 28 days.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. PESTICIDE STORAGE: Store in the original container in a dry, temperature-controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/encapsulate spillage and use in accordance with instructions below or dispose of in accordance with the "Pesticide Disposal Instructions" listed below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Nonrefillable container: DO NOT reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available, or place empty container in trash.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC. BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc. (RV051414)

LESCO® T-Storm™ 2G Fungicide is not manufactured or distributed by Cleary Chemical Corporation, seller of Cleary 3336™. Cleary 3336™ is a trademark of Cleary Chemical Corporation. LESCO® is a registered trademark and T-Storm™ is a trademark of LESCO Technologies, LLC.

FG-55709 02 MADE IN THOMASVILLE Customer: NUFARM Brand: LESCO T-STORM 2G Date: 10.07.2015

AMG 003 BLACK

Please Note: This proof is NOT accurate for color