

ORGANIC INPUT MATERIAL

SprayBor Lite

Guaranteed Analysis:
Boron (B) 16.5%
Derived from boric acid

A Soluble Boron Powder for use in Organic Production

For use in organic production as a fertilizer in foliar, fertigation and soil applications to supplement boron levels in agricultural crops.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- SprayBor Lite does not require any additional surfactants.
- Aerial application and ground sprays – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Dissolve a maximum of 1 lb in 3 gallons of water.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- SprayBor Lite is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- **Consult a NutriAg representative for application timing and necessity of its use.**

Suggested Application Rates:

Crop	Foliar Rates*	Fertigation Rates*	Soil Rates*
Forage Crops: Alfalfa, Clover, Hay Crops Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane Legumes: Dry bean, Lentil, Peanut, Pea, Soybean Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince Other Fruit: Citrus (all), Fig, Kiwi, Melons Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut Grapes: Currant, Raisin, Table, Wine Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	Typical rates are 0.5-2 lbs/ acre/ application Maximum rate of 4 lbs/acre/ application with no more than 6 lbs/season Caution: Crops vary in their sensitivity to boron.	The number and frequency of applications determine application rates. Apply no more than 3 lbs/acre/ application. Apply no more than 6 lbs/season. Always apply toward the end of an irrigation cycle.	Caution: Overapplication of boron to soil can cause crop injury. Broadcast evenly on soil surface or apply dissolved in irrigation water. Maximum rate of 10 lbs/acre.

*Crops vary in their sensitivity to boron. Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: The use of the fertilizing material on any crops other than those recommended may result in serious injury to crop(s).

Caution: KEEP OUT OF REACH OF CHILDREN.

OMRI
Listed

Listed by the Organic Materials Review Institute (OMRI) for use in organic production.



Warning

Causes serious eye irritation. May damage fertility of the unborn child if inhaled and/or swallowed. Contains 4-5% of ingredients of unknown toxicity (dermal).

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves/ protective clothing/eye protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations.

Net Weight:
50 lbs. (22.7 kg)

Batch No.: 120619



NutriAg Ltd.
39 Gail Grove, Toronto, ON, Canada, M9M 1M5
Tel: (416) 636-1555 www.nutriag.com