



## Sulfur Complete SP

Foliar Nutritional Soluble Powder

A formulated soluble powder mixture of micronutrients with magnesium and sulphur.

### GUARANTEED ANALYSIS

Magnesium (Mg) .....	8.0%
8.0% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	14.5%
14.5% Water Soluble Combined Sulfur (S)	
0.00% Free Sulfur (S)	
Boron (B) .....	5.0%
Manganese (Mn) .....	7.0%
7.0% Water Soluble Manganese (Mn)	
Molybdenum (Mo) .....	0.4%

Derived from Magnesium Sulfate, Manganese Sulfate, Sodium Borate, and Sodium Molybdate.

Chlorine (Cl), Maximum .....

### KEEP OUT OF REACH OF CHILDREN DANGER

Causes serious eye damage.  
May damage fertility. May damage the unborn child.  
May cause damage to organs through prolonged or repeated exposure if inhaled.  
Harmful to aquatic life with long lasting effects.



NET WEIGHT: 20.0 lbs. (9.07 kg) SN 0714/0315

Information about the components of this lot of fertilizer may be obtained by writing to Helena Chemical Company, 225 Schilling Boulevard, Suite 300, Collierville, TN 38017 and giving the lot number which is found on the container.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Contents Made In Great Britain

F224

MANUFACTURED FOR

HELENA CHEMICAL COMPANY

225 SCHILLING BOULEVARD, SUITE 300

COLLIERVILLE, TN 38017

901-761-0050

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### DANGER

BEFORE USING THIS PRODUCT, READ ALL PRECAUTIONS, DIRECTIONS FOR USE, CONDITIONS OF SALE—LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES.

Causes serious eye damage. May damage fertility. May damage the unborn child. May cause damage to organs through prolonged or repeated exposure if inhaled. Harmful to aquatic life with long lasting effects. Do not handle until all safety precautions have been read and understood. Do not breathe dust. Avoid release to the environment. Contains manganese compounds. Injurious to health by ingestion or inhalation of excessive quantities. Injurious to eyes and skin. In case of contact with eyes and/or skin, immediately flush with plenty of water and seek medical attention.

This fertilizer contains boron which may be injurious to certain crops. This fertilizer is to be used only on soils which respond to molybdenum (Mo). Crops high in molybdenum (Mo) are toxic to grazing animals (ruminants).

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Use personal protective equipment as required. Wear suitable gloves and faceshield when handling this product.

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.

### FIRST AID

IF IN EYES:	Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. <b>If exposed or concerned:</b> Get medical attention.
IF INHALED:	Remove to fresh air. Get medical attention immediately.
IF IN CONTACT WITH SKIN:	Wash with soap and water. Continue to rinse for at least 10 minutes. Get medical attention.
IF SWALLOWED:	Wash out mouth with water. If exposed person is conscious, give small quantities of water to drink. Do not induce vomiting unless directed to do so by medical personnel. Get medical attention.

### STORAGE AND DISPOSAL

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost (39°F) and other climatic extremes (temperatures exceeding 100°F). Keep container tightly closed. Do not allow water to be introduced to the contents of this container. Do not contaminate water sources by cleaning of equipment or disposal of spray waste. Dispose of empty containers by triple rinsing with detergent solution or puncture and discard empty containers in a landfill in accordance with current local, State and Federal regulations

### GENERAL INFORMATION

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis. Contact your county agent or field consultant for specific information.

### DIRECTIONS FOR USE

#### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Slowly add the product through the top filter into the tank three-quarters filled with water, while agitating. Top off with water and continue to agitate until spraying is completed.

### APPLICATION RATES AND TIMINGS

**Alfalfa:** 1-3 lbs/acre applied at flower bud formation and again at the start of flowering. Water rate: 20 gallons/acre

**Asparagus:** Two applications of 1-3 lbs/acre during vegetative growth at a 10-14 day interval. Water rate: 20 gallons/acre.

**Beans:** 1-3 lbs/acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4-6 leaf stage). Repeat applications can be made at 10-14 day intervals. Water rate: 5-20 gallons/acre.

**Blueberries: (High Bush Blueberries Only):** Apply 1-3 lbs/acre just before the onset of leaf drop. Repeat the application at bud separation the following fruiting season. Water rate: 20 gallons per acre.

**Broccoli:** 1-3 lb/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 20 gallons/acre.

**Brussel Sprouts:** 1-3 lbs/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 5-20 gallons/acre.

**Cabbage:** 1-3 lbs/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 5-20 gallons/acre.

**Calabrese:** 1-3 lbs/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 5-20 gallons/acre.

**Canola:** 1-3 lbs/acre at 4-9 true leaves, followed by a further application 10 days later. For fall-sown crops, apply prior to winter dormancy and in spring up to flower buds enclosed (G.S. 31). Water rate: 5-20 gallons/acre.

**Cauliflower:** 1-3 lbs/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 20 gallons/acre.

**Cereals (Wheat):** 1-3 lbs/acre at growth stage F3-F6. Water rate: 5-20 gallons/acre.

**Cole Crops:** 1-3 lbs/acre at 4-6 leaf stage, with 1 or 2 repeat applications at 10-14 day intervals if necessary. Water rate: 5-20 gallons/acre.

**Corn:** 1-3 lbs at the 4-8 leaf stage. Repeat 10-14 days later if necessary. Water rate: 5-20 gallons/acre.

**Cotton:** 1-3 lbs/acre applied 3-4 weeks after emergence. Repeat once or twice at 10-14 day intervals up to the start of flowering. Water rate: 5-20 gallons/acre.

**Courgette (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Cucumber (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Cucurbits (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Grapevines:** 1-3 lbs per acre. Two applications pre-flowering at 10-14 day intervals. Also, if necessary, one or two applications at no less than a 10-14 day interval from fruit set to start of ripening. Water rate: 20-50 gallons/acre.

**Groundnuts/Peanuts:** 1-3 lbs/acre as early in the growing season as possible, when there is sufficient leaf cover to intercept the spray. Repeat at 10-14 day intervals as necessary. Water rate: 5-20 gallons/acre.

**Melon (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Onions:** Two applications of 1-3 lbs/acre applied as soon as there is adequate leaf cover (6 leaves) and repeated 2 weeks later. Water rate: 20 gallons/acre.

**Peas:** 1-3 lbs/acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4-6 leaf stage). Repeat applications can be made at 10-14 day intervals. Water rate: 20 gallons/acre.

**Potatoes:** 1-3 lbs/acre applied one week after 100% emergence with a second application 10-14 days later. Water rate: 20 gallons per acre.

**Pumpkin (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Soybean:** 1-3 lbs/acre applied at the 4-6 leaf stage. Repeat as necessary at 10-14 day intervals. Water rate: 5-20 gallons per acre.

**Spinach:** 1-3 lbs/acre 10-14 days after transplanting or emergence. Repeat after 10-14 days if necessary. Water rate: 50 gallons/acre

**Strawberry (field grown):** 1-3 lbs/acre applied to new leaf growth in the spring, and again 10-14 days later (pre-flowering). N.B. Do not apply during flowering. Water rate: 20 gallons/acre.

**Tobacco:** 3 lbs/acre two to three weeks after transplanting (3-4 leaf stage). Repeat 10-14 days later if necessary. Water rate: 10-50 gallons/acre.

**Water Melon (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

**Zucchini (field grown):** Two applications of 1-3 lbs/acre sprayed at the male flowers stage and again 10-14 days later. Water rate: 50 gallons/acre.

---

#### **TANK MIXING/CO-APPLICATION**

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. Before co-application of products you, or your advisor, must visit [www.tankmix.com/elemax](http://www.tankmix.com/elemax) for important information or do a jar test.

---

#### **PRECAUTIONS**

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g., rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

---

#### **CONDITIONS OF SALE—LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES**

Read the Conditions of Sale—Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

This label is believed to be reliable and must be followed carefully. Injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to subject to factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2015

ELE-MAX® and PRECISION NUTRITION® are registered trademarks of Helena Holding Company.

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.