



Active Ingredient: Garlic 20.14%
 Other Ingredients (Water, Citric Acid):..... 79.86%
 Total.....100.00%

This product is FIFRA 25 (b) exempt. All ingredients are exempt from tolerance requirements under 40 CFR 180.

PRECAUTIONARY STATEMENT

Human and Animal Health Hazards

Signal Word: **WARNING**

Hazard Statement: Causes eye irritation.

Precautionary Statement: Wear protective eyewear, such as goggles. Avoid contact with eyes or clothing. Do not ingest. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If symptoms persist, seek medical attention.

USE RESTRICTIONS AND PRECAUTIONS

- Not warranted to kill insects. This product is a repellent.
- Do not apply **BioLink Insect & Bird Repellent** within 12 hours of crop harvest.
- Do not apply **BioLink Insect & Bird Repellent** if rain is forecasted or overhead watering occurs within 12 hours.
- **POLLINATION WARNING: MAY REPEL BEES.** BioLink Insect Repellent may repel bees. Do not apply less than three days prior to pollination.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Do not store near heat or open flame. Keep in original container. Store in cool, dry place. Protect from excessive heat. Do not store below 4°C as product will freeze. **Disposal:** Do not reuse container. Triple rinse and offer empty containers for recycling or reconditioning, or dispose of in a sanitary landfill.

Notice of Warranty: Westbridge warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. In no case shall Westbridge be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



MANUFACTURED BY
Westbridge

1260 Avenida Chelsea • Vista, CA 92081
 (800) 876-2767 • www.westbridge.com

For use on all crops including fruits and vegetables, fruit trees, vines, tree nuts, field and row crops, ornamental trees, shrubs and flowers, containerized plants, and turf

Repels insects such as aphids, beetles, borers, caterpillars, earwigs, fleas, flies, leafhoppers, leafrollers, lygus bugs, mealy bugs, mites, moths, scales, slugs, thrips, weevils, whiteflies, and other plant feeding insects.

Also great for repelling birds, deer, field mice, rabbits and other animal pests.

APPLICATION RATES

Foliar Application: ½ to 4 quarts per acre; use higher rates for heavier pest infestations. Do not spray to point of run-off.

Nursery & Container: ½ to 1 quart per 100 gallons of buffered water.

Home & Garden: 2 to 4 ounces per gallon of water; Spray plant foliage thoroughly including underside of leaves.

MIXING INSTRUCTIONS

General mixing instructions: Shake container well or stir contents thoroughly before adding **BioLink Insect & Bird Repellent** to the spray tank.

Buffering instructions for agricultural and large scale uses: spray tank pH must be adjusted to 5.5 to 6.0 before adding **BioLink Insect & Bird Repellent** to the spray tank. Recommended buffering agents are BioLink Buffer and Penetrant, citric acid, acetic acid, phosphoric acid or aliphatic polycarboxylate-type materials. Do not buffer with propionic acid or similar products that affect biologically active products. After mixing, agitate spray tank-mix before and during application to prevent settling.

Tank mixing with other products: when mixing **BioLink Insect & Bird Repellent** with other products, it is strongly recommended to test a small plot before applying the spray mix.

If tank-mixing **BioLink Insect & Bird Repellent** with other pesticides, use the recommended label rates of the other pesticides until exact tank-mixed product performance is determined.

APPLICATION TIMING AND PEST PRESSURE

When using **BioLink Insect & Bird Repellent**, the optimum pest pressure to start application is 0-5%. Repeat application timing should be 8-14 days, and determined by field scouting reports of pest pressure, plant nutrition and temperature.

SPRAY TEMPERATURE: The optimum temperature range to apply BioLink Insect & Bird Repellent is 70°-90°F (21°-32°C). Minimum recommended application temperature is 50°F (10°C). Maximum recommended application temperature is 100°F (38°C). For applications above 100°F higher application spray rates (more than 40 gals/acre) are recommended after there has been a minimum of a 15°F (9°C) drop in air temperature.

SPRAY ADJUVANTS: To maximize the effectiveness of BioLink Insect & Bird Repellent add 4 oz of BioLink Spreader-Sticker per 100 gallons water per acre. Silicone-based or non-ionic surfactants are strongly recommended to increase product coverage and absorption. If **BioLink Insect & Bird Repellent** is not rapidly absorbed by the foliage, it will not be effective.

NET CONTENTS:

2.5 Gallons (9.5 L) / 21.0 Lbs (9.5 KG)

WSDA Reg. No. 51517-13003