

PCNB 12.5% Plus Fertilizer

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
For the control of certain soil-borne turfgrass diseases.

Active Ingredient:

Pentachloronitrobenzene 12.50%

Inert Ingredients: 87.50%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. This material may cause skin irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt while handling product. Wash hands and face after use and prior to eating, drinking, or smoking. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CHEMIGATION

Do not apply this product through any type of irrigation system.

LAWNS AND TURF

Brown Patch and Dollar Spot: For southern or warm season lawns such as St. Augustine, Bermuda, and similar grasses: Make one application in the fall or spring at first indication of infection. Apply 6 lbs. per 1,000 sq. ft. Following application, the treated areas should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

For Dichondra: Treat as above, using 6 lbs. per 1,000 sq. ft. For Northern or cool season lawns such as Ryegrass, Fescue, Bluegrass, Bent, or similar grasses: apply 1.6 to 2.0 lbs. per 1,000 sq ft at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. Water treated area lightly to wash chemical down to soil. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

Helminthosporium Leaf Spot: Apply at rate of 4 to 6 lbs. per 1,000 sq. ft. Preventive applications should be made in Spring or Fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

Snow Mold (*Typhula incarnata*, *Fusarium nivale*): Apply at the rate of 4 to 6 lbs. per 1,000 sq. ft. using a granular lawn spreader or comparable equipment. Application should be made prior to first snowfall.

EPA Reg. No. 10404-37

EPA Est. No. 10404-OH-04 (M), 10404-FL-01 (S)

First Letter of Lot Code Indicates Manufacturing Site

#014804

F1406

Manufactured by: LESCO, Inc.

15885 Sprague Road
Strongsville, Ohio 44136-1772

Do not graze treated areas or feed clippings to livestock.

PRECAUTION: For northern or cool season grasses, such as Ryegrass, Fescue, Bluegrass, Bent, or similar grass, application is NOT recommended when temperatures exceed 75 °F.

SUGGESTED SPREADER SETTINGS

Spreader	Recommended Rates in lb per 1,000 sq ft			
	1.6 lb	2.0 lb	4.0 lb	6.0 lb
LESCO	#8	#9	#11	#14
SCOTTS® R8A	G	G ½	H ½	J
Cyclone® or Spyker®	1½	2	3	4

*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application using the LESCO Calibrating Gauge #009952.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage Instructions: Store in original container only, separate from other pesticides, fertilizer, food, and feed. Do not store near open flame or heat sources.

Pesticide Disposal Instructions: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

10-3-23

ELITE® FERTILIZER GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 10.00%

0.60% Ammoniacal Nitrogen

9.40% Urea Nitrogen

AVAILABLE PHOSPHATE (P₂O₅) 3.00%

SOLUBLE POTASH (K₂O) 23.00%

SULFUR (S) Total 11.20%

3.00% Free Sulfur (S)

8.20% Combined Sulfur (S)

DERIVED FROM: Polymer Coated Sulfur Coated Urea, Urea, Ammonium Phosphate, Sulfate of Potash.

*3.00% Slowly available Nitrogen from LESCO Poly Plus® Sulfur Coated Urea.

Brush off walks and painted surfaces following application to avoid discoloration.

WARRANTY

LESCO, Inc. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. Regulatory Affairs Department, 15885 Sprague Road, Strongsville, OH 44136-1772, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a1.com>

LESCO and Poly Plus are registered trademarks of LESCO Technologies, LLC. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.

G:\REGULWP\STRONGSV\Aprvd Lbl - Word\Control Products\PCNB\014804.doc

Created 8/28/03 Rev. 9/3/03 WBW (021003)

Net Weight: 40 lb (18.2 kg)



(01) 00758073148043