

**USE AND RATE RECOMMENDATIONS FOR TURF**

DISEASE	TURF TYPE	FL. OZS. PER 1000 SQ. FT.	APPLICATION NOTES
Brown Patch ( <i>Rhizoctonia solani</i> )	Warm climate grasses such as: St. Augustine Bermuda	16-24	Mix specified amount in 1-5 gals. of water and apply a single application in the spring or fall or at first indication of disease. If disease is severe or reappears, area should be retreated 3 - 4 weeks later. Mix specified amount in 5-10 gals. of water and treat as above.
	Dichondra	16-24	Mix specified amount in 1-5 gals. of water and apply at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85°F or higher.
	Cool climate grasses such as: Bluegrass Ryegrass Fescue Bent	4 1/2 to 6	Mix specified amount in 1-5 gals. of water and apply as a preventative application in the spring or fall of each year. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85°F or higher. For maximum spring disease protection of Melting Out/Leaf Spot on cool season grasses, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications of two week intervals. Do not make more than a total of 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.
Melting Out/ Leaf spot* ( <i>Drechslera</i> spp., <i>Biopolaris</i> spp., <i>Exserchilium</i> spp.)	All	10 1/2 to 15	as a preventative application in the spring or fall of each year. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85°F or higher. For maximum spring disease protection of Melting Out/Leaf Spot on cool season grasses, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications of two week intervals. Do not make more than a total of 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.
Dollar Spot* ( <i>Sclerotinia homoeocarpa</i> )	All	3	as a preventative application in the spring or fall of each year. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85°F or higher. For maximum spring disease protection of Melting Out/Leaf Spot on cool season grasses, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications of two week intervals. Do not make more than a total of 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.
Gray Snow Mold ( <i>Typhula</i> spp.)	Cool climate grasses	12-16	one application prior to first snowfall or when temperatures remain below 60°F and extended wet conditions are expected. In the absence of snowcover, supplemental applications of 3-6 fl. ozs. can be applied at 4 to 6 week intervals to maintain control when disease pressure is severe. *NOTE: This product must be moved into the soil/thatch layer to be effective. Applications made in less than 5 gals. of water must receive 0.25" of irrigation or rainfall prior to snow cover. For maximum spring disease protection of pink snow mold, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications at two week intervals. Do not make more than a total 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.
Pink Snow Mold ( <i>Microdochium nivale</i> )	Cool climate grasses	3	one application prior to first snowfall or when temperatures remain below 60°F and extended wet conditions are expected. In the absence of snowcover, supplemental applications of 3-6 fl. ozs. can be applied at 4 to 6 week intervals to maintain control when disease pressure is severe. *NOTE: This product must be moved into the soil/thatch layer to be effective. Applications made in less than 5 gals. of water must receive 0.25" of irrigation or rainfall prior to snow cover. For maximum spring disease protection of pink snow mold, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications at two week intervals. Do not make more than a total 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.

\*Use not registered by the California Department of Pesticide Regulation

**LESCO**®

**Revere™ 4000**  
Flowable Turf Fungicide

**COMPOSITION**

**Active Ingredient: (% by weight)**  
Pentachloronitrobenzene (PCNB) ..... 40%  
**Inert Ingredients:** ..... 60%  
**TOTAL** ..... 100%

Contains 4 lbs. of Pentachloronitrobenzene per U.S. gallon.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID	
<b>IF IN EYES:</b>	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or doctor for treatment advice.
<b>IF ON SKIN OR ON CLOTHING:</b>	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<b>IF INHALED:</b>	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<b>IF SWALLOWED:</b>	• Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>CROMPTON EMERGENCY PHONE</b>	<b>203-723-3670</b>
<b>SAFETY DATA AND INFORMATION</b>	<b>203-573-3303</b>
<b>TRANSPORTATION EMERGENCY (CHEMTREC)</b>	<b>800-424-9300</b>

See side panel for additional precautionary statements.

UCC  
(01)00758073608998

**#060899**  
**Net Contents: 2.5 gal. (9.46 l)**

EPA Reg. No. 400-454-10404

EPA Est. No. 37429-GA-01

Distributed by LESCO, Inc. • 1301 East 9th Street • Cleveland, OH 44114-1849

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS**  
**AND DOMESTIC ANIMALS**  
**CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

**Applicators and Other Handlers (other than mixers and loaders) Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks.

**Mixers and Loaders Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks. For exposures in enclosed areas: A respirator with an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter. For exposure outdoors: Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product to turf through any type of irrigation system.

Do not plant root crops in PCNB treated fields within 12 months of broadcast or banding applications unless PCNB is registered for use on those crops.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves such as barrier laminate or nitrile rubber, or neoprene rubber, or viton
- shoes plus socks

**GENERAL INFORMATION**

REVERE 4000 is a preventative fungicide recommended for control of certain soil borne diseases of turf. Best results are achieved when used in a preventative program following recommended rates and application directions. Carefully read, understand and follow label directions.

**COMPATIBILITY**

For broad spectrum disease control, this product can be tank mixed with TERRAZOLE® 35WP\* (EPA Reg. No. 400-416) for use on turf in accordance with the more restrictive label limitations and precautions. No label dosage rates should be

exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not tank mix with other chemicals unless prior use has proven compatibility.

**LAWNS AND TURF**

REVERE 4000 is a fungicide for the prevention and control of certain soil borne diseases on golf courses, sod farms, home lawns and institutional areas where turf is grown. Best results are obtained when REVERE 4000 is used as a preventative application before diseases appear, however, if you do not apply as a preventative treatment, REVERE 4000 should be applied at first sign of disease. Areas treated with REVERE 4000 should be watered following application to move material down to soil level. If treated area is subjected to flooding or unusually heavy rainfall, or if disease is severe or reappears, the area should be treated again. See use and rate recommendations for retreatment intervals. Do not graze treated areas or feed clippings to livestock. Under certain growing conditions, a temporary discoloration of the grass may occur. This causes no harm and will disappear in a short time.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not reuse empty containers. Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

LESCO is a registered trademark; Revere and the sweeping design are trademarks of LESCO Technologies LLC. Terrazole is a registered trademark of Crompton Corporation. 10/12/04 CV (021003)