

Weed & Feed

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.560%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.145%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	0.143%
INERT INGREDIENTS:	99.152%
Total	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.46%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.12%
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	0.12%

†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p
Patent No. 5,266,553

18-2-3

**MINI FERTILIZER
GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N)	18.00%
0.75% Ammoniacal Nitrogen	
17.25% Urea Nitrogen*	
Available Phosphate (P ₂ O ₅)	2.00%
Soluble Potash (K ₂ O)	3.00%
Sulfur (S) Total	2.70%
2.70% Free Sulfur (S)	
Derived From: Polymer Coated Sulfur Coated Urea, Urea, Ammonium Phosphate, Muriate of Potash.	
Chlorine (Cl) Max	2.50%
*4.85% Slowly Available Nitrogen from LESCO Poly Plus® Sulfur Coated Urea.	
Made in U.S.A.	F1406

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. LESCO Weed and Feed is a stable mixture of three herbicides. It is for use on Residential turf.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed, or other pesticides.

DISPOSAL: If Empty – Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled** – If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bident, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black eyed Susan, Black medic, Black Mustard, Blackseed Plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curley dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False Sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleaf marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherd's purse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiney Amaranth, Spiny cocklebur, Spotted catsear, Spotted knotweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperseed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results: 1) Avoid mowing 1 to 2 days before and after application; 2) Apply when grass is moist (after a rain, sprinkling or dew); 3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and 4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

SPREADER SETTINGS: The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 sq ft.

One pass Application:

MAKE	SETTINGS
LESCO Calibration Gauge	#13
SCOTTS® R8A	1 ½
Cyclone® or Spyker®	3 ½
LESCO Pendulum	20
Lely®	3 ½ II

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Warranty

LESCO, Inc. warrants that this product conforms to the analysis on its label. When used in accordance with label directions, under normal conditions, this product is reasonably fit for its intended purposes. Since time, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. RA Dept., 15885 Sprague Road, Strongsville, OH 44136-1772, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a-1.com>

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. SCOTTS is a registered trademark of the SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. Lely is a registered trademark of C Van Der Lely N. V.

G:\REGULWP\STRONGSV\Aprvd Lbl - Word\Control Products\Weed & Feed\026217.doc
Rev. 9/12/03 WBW (061803)

E.P.A. Reg. No. 228-281-10404
E.P.A. Est. No. 10404-OH-04 (M), 10404-FL-01 (S)
First Letter of Lot Code indicates manufacturing site.

Net Weight 50 lb (22.7 kg)

#026217

F1406

Distributed by: **LESCO, Inc.**
15885 Sprague Road
Strongsville, Ohio 44136-1772

