

Atrazine 1.05% Plus Fertilizer

For St. Augustine and Centipede Turfgrasses

A LESCO® TURF FOOD WITH ATRAZINE FOR CONTROL OF BROADLEAF WEEDS AND WEEDY GRASSES

50 Pounds covers up to 15,230 Sq.Ft. of Turfgrass
Prevents Crabgrass, Paspalums, and Spotted Spurge.
Controls Florida Pusley, Lippia, Pigweed.
Over Forty-Five More Weeds as Listed.

A WEED & FEED FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS LAWNS

ACTIVE INGREDIENTS:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine 1.01%
Related Compound 0.04%

INERT INGREDIENTS: 98.95%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid inhalation or contact with eyes, skin or clothing. Remove clothing and wash clothing before reuse. Keep away from feed or food products.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks and a Dust/Mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C or a NIOSH approved respirator with any N, R, P or HE filter. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. If product is spilled, while wearing a dust mask, sweep up product and use as this label instructs (as a Weed & Feed) or dispose of in accordance with the STORAGE AND DISPOSAL section on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes and socks.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (<i>Poa annua</i>)	Cutleaf Evening Primrose	Oxalis (<i>Common and Yellow Woodsorrel</i>)
Annual Sowthistle	Dichondra	Pigweed
Clover	Dollar Weed	Purslane
Black Medic	Florida Beggarweed	Ragweed
Blue Toadflax	Florida Betony	Red Sorrel
Bull Paspalum	(<i>Rattlesnake Weed</i>)	Rescuegrass
Buckhorn Plantain	Florida Pusley	Sandspur
Bur Clover	Fragrant Cudweed	Sedge*
Carolina Geranium (<i>Cranesbill</i>)	(<i>Rabbit Tobacco</i>)	Shepards Pursue
Chickweed (<i>Common and Mouse Ear</i>)	Henbit	Sicklepod
Cocklebur	Knotweed	Spotted Spurge
Common Dandelion	Lambsquarter	Spurweed
Common Purslane	Lespedeza	Stinging Nettle
Crabgrass	Lippia (<i>Matchweed</i>)	Swinecrest
Crowfoot Grass	Moneywort	Venus Looking Glass
Cudweed	Morning Glory	Virginia Creeper
	Narrowleaf Vetch	Wild Mustard
	Parsley-Piert	Wood Sorrel

NON-AGRICULTURAL REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

IMPORTANT: In addition to providing turfgrass with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from seeds".

FOR WEEDS THAT ARE PRESENT: Includes broadleaf weeds from the list such as; Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.

FOR WEEDS GERMINATING FROM SEEDS: Applied before weed seeds germinate; includes grassy weeds from the list such as; Crabgrass, Paspalum, Sedge* (an additional application may be required to control Sedge and Spotted Spurge. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

WHERE TO USE

This material can be used on St. Augustinegrass and Centipedegrass grown in home and corporate lawns, parks and recreational areas, golf courses, cemeteries, athletic fields, and sod farms. Do not plant any crop (flowers or vegetables) to treated areas for 18 months since plant injury may result.

DO NOT USE ON ANY VARIETIES OF BLUEGRASS, FESCUE, BENTGRASS, BAHIAGRASS, RYEGRASS, BERMUDAGRASS, OR DICHONDRA LAWNS.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good weed control.

For pre-emergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for arepeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

HOW MUCH TO APPLY

This product may be applied with a broadcast rotary type spreader. Apply 3.27 lbs to 4.37 lbs uniformly over 1,000 sq ft of lawn area. (equivalent to the required 1.5 lb to 2.0 lbs Atrazine active ingredient per acre). This is also equal to 16 1/2 pounds of this product to treat 3,775 sq ft up to 5,000 sq ft. Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommended rates. Apply to dry grass. For best results, rainfall or watering is necessary within 7 to 10 days after treatment to activate the herbicide and fertilizer by washing the product off the grass blades down to the soil surface.

USE PRECAUTIONS

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

22-2-11

MINI FERTILIZER

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 22.00%

4.40% Ammoniacal Nitrogen

17.60% Urea Nitrogen*

AVAILABLE PHOSPHATE (P₂O₅) 2.00%

SOLUBLE POTASH (K₂O) 11.00%

SULFUR (S) Total 8.00%

4.50% Free Sulfur (S)

3.50% Combined Sulfur (S)

IRON (Fe) Total 1.00%

0.01% Water Soluble Iron (Fe)

MANGANESE (Mn) Total 0.50%

0.07% Water Soluble Manganese (Mn)

DERIVED FROM: Polymer Coated Sulfur Coated Urea, Urea, Ammonium sulfate, Ammonium Phosphate, Muriate of Potash, Iron Sulfate, Manganese Sulfate.

CHLORINE (Cl) Max. 8.25%

*4.40% Slowly Available Nitrogen from LESCO Poly Plus® Sulfur Coated Urea.

Made in U.S.A.

F526

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY

NOTICE: LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. LESCO, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall LESCO, Inc. be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept, 15885 Sprague Rd, Strongsville, OH 44136-1772, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a1.com>

LESCO and Poly Plus are registered trademarks of LESCO Technologies, LLC.

G:\REGUL\WP\STRONGSV\Aprvd Lbl - Word\Control Products\Atrazine\069344FL.doc Rev. 4/15/04 WBW (090701)

Net Wt. 50 lb (22.7 kg)



EPA REG. NO. 10404-94

EPA EST. NO. 10404-FL-01(S)

First letter of lot code indicates manufacturing site.

#069344

F526

Manufactured by: **LESCO, Inc.**
722 Webster Turn Drive
Sebring, FL 33870