



# Corral<sup>®</sup>

## 2.68G



## Ornamental Weedgrass Control

- Preemergent control of goosegrass, crabgrass, *Poa annua* and other annual grassy weeds
- Preemergent control of certain broadleaf weeds
- Contains 2.68% pendimethalin

### PACKAGING

#### Unit Size:

40 lb. bag (18.14 kg)

**STOCK #95669**

(See next page  
for label information)

Distributed by:

Scotts-Sierra Crop Protection Company  
14111 Scottslawn Road • Marysville, OH 43041

If you have any questions regarding the use of this product, call TOLL FREE 1-800-492-8255

For Sale To, Use and Storage By Commercial Nursery,  
Cut Flower, Foliage And Landscape Personnel Only.



**Ornamental  
Weedgrass Control**

<b>ACTIVE INGREDIENT:</b>	
Pendimethalin† . . . . .	2.68%
<b>OTHER INGREDIENTS . . . . .</b>	<b>97.32%</b>
	<b>TOTAL 100.00%</b>

†CAS # 40487-42-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:** CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. May be harmful if swallowed. Avoid inhalation.

EPA Reg. No. 538-188

EPA Est. No. 538-OH-1 (S)

EPA Est. No. 8378-IN-1 (K)

Product of U.S.A.

**NET WEIGHT: 40 LBS. (18.14 kg)**

Subscript used is first letter of run code on end of container



Scotts-Sierra Crop Protection Company  
14111 Scottslawn Road  
Marysville, OH 43041  
1-800-492-8255

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eye-wear and dust/mist filtering respirator (MSHA/NIOSH, approval number prefix TC-21) or a NIOSH approved respirator with any R, P, or HE filter.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

**Environmental Hazards:** This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Personal protective equipment required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks, protective eyewear.

**Controls these grassy weeds as they sprout:**

barnyardgrass	foxtail	jungerice	signalgrass
crabgrass	goosegrass	lovegrass	sprangletop
crowfootgrass	itchweed	Poa annua	witchgrass
fall panicum	johnsongrass	sandbur, field	woolly cupgrass
	(from seed)		

**Controls these broadleaf (dicot) weeds as they sprout:**

carpetweed	henbit	pearlwort	rocket, London
chickweed	hop clover	pigweed	shepherdspurse
cudweed	kochia	puncturevine	smartweed
eveningprimrose	knotweed	purslane	velvetleaf
fiddleneck	lambsquarters	pusley, Florida	
filaree	oxalis	prostrate spurge	

## ORNAMENTALS

**For use on these container and field grown ornamentals**

Acer	Correa	Magnolia	Quercus
Achillea	Cortaderia	Malus	Rhododendron
Acorus	Euonymus	Miscanthus	Solanum
Agapanthus	Franklinia	Moraea	Spirea
Ajuga	Fraxinus	Ophiopogon	Stokesia
Aloysia	Hedera	Pachysandra	Syringa
Arbutus	Hemerocallis	Penstemon	Tilia
Arundo	Hosta	Photinia	Trachycarpus
Asparagus fern	Ilex	Pinus	Tsuga
Butia	Juniperus	Pittosporum	Washingtonia
Ceanothus	Lagerstroemia	Platanus	Wisteria
Chamaerops	Ligustrum	Potentilla	
Convolvulus	Liriope	Pseudotsuga	
Cordyline	Lobelia	Pyrus	

**Ground Covers**

Aptenia	Fragaria	Myoporum	Verbena
Arctotheca	Gazania	Osteospermum	Vinca
Baccharis	Helix	Pelargonium	
Carpobrotus	Hypericum	Rosemarnus	

**Perennials**

Aster	Asclepias	Lythrum
-------	-----------	---------

**Ornamental Grasses**

Ammophila	Deschampsia	Festuca	Pennisetum	Phalaris
-----------	-------------	---------	------------	----------

### Ornamentals - Application Recommendations

For goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, cudweed, hop clover, oxalis, eveningprimrose, chickweed, purslane, knotweed, henbit, prostrate spurge and pearlwort, apply in the spring or fall before they sprout or after established weeds are removed.

Apply this product to dry foliage only. Apply either prior to weed seed germination or following complete weed removal.

Water immediately (1/2 to 1 inch) to remove granules from plant foliage and activate the herbicide.

For container grown ornamentals delay first application of this product to bare-root liners 2-4 weeks.

Remove established weeds from potted liners prior to transplanting into containers. After planting irrigate with approximately 1/2 inch of water to firm potting media around root system. After two regular waterings and foliage has dried, broadcast product evenly over container growing area. Water in with 1/2 to 1 inch.

First application to field grown ornamentals can be made following planting or prior to weed seed germination. Soil should be broken by light cultivation and be free of all established weeds.

### PRECAUTIONS - ORNAMENTALS

Does not control mature established weeds.

Do not apply to moist foliage.

Do not apply product in enclosed greenhouse or polyhouse structures as ornamental injury may result.

Following applications, do not disturb the soil surface as the herbicide barrier will be broken.

Do not incorporate product into soil or potting media nor apply product directly to bare roots of woody ornamentals.

Do not apply in wind conditions that affect distribution of the product.

Not for use on vegetables. Do not use on fruit trees that will bear fruit within one year of application.

Users who wish to use Corral® 2.68G on ornamental species or non-bearing fruit trees not listed on this label may determine suitability for such uses by making trial application of this product at a recommended rate to a small number of plants. Prior to using this product on a large number of plants, the treatment should be observed for signs of herbicidal injury for 3 to 6 months to determine if the treatment is safe to the target plant species.

Non-bearing fruit trees such as citrus (oranges, grapefruit, etc.), pome fruits (apples, pears, etc.) and stone fruits (cherries, peaches, etc.) are plants which will not bear fruit for at least one year following application.

The users assume responsibility for any crop damage or other liability from the use of this product on ornamental plant species not recommended on this label.

WEEDS	RATES OF CORRAL 2.68G	COMMENTS
Corral 2.68G alone.	barnyardgrass crabgrass crowfootgrass carpetgrass chickweed cudweed eveningprimrose fall panicum fiddleneck filaree foxtail goosegrass henbit hog clover itchgrass johnsongrass (from seed) junglerice knotweed kochia lambsquarters lovegrass oxalis pearlwort pigw/Oeed Poa annua prostrate spurge puncturevine purslane pusley, Florida rocket, London sandbur, field shepherdspurse signalgrass smartweed sprangleweed velvetleaf witchgrass woolly cupgrass	Apply a repeat application of 76 lbs./acre after 3-5 months to extend control or for weeds that germinate in the fall.
	cudweed hop clover eveningprimrose henbit	114 lbs./acre prior to weed germination.
ROUT® or OH2® plus Corral 2.68G.	Corral® 2.68G can be used to control weeds while plants are producing a new flush of spring growth, or to extend annual weed control after an application of ROUT or OH2. Allow at least 6-8 weeks between an application of ROUT, OH2 or Corral 2.68G	

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT NOTICE: LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its use on ornamentals. However, it is understood that tests may not have been carried out on all varieties and under all growing conditions. The user should always follow label directions and exercise judgment and caution when using this product on a given variety until familiar with the results under their growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.