



Truban[®]

25 EC



Fungicide
25% Emulsifiable Concentrate
For control of damping-off, root and stem rot diseases caused by Pythium, Phytophthora, in ornamental and nursery crops.

- Fungicidal, not fungistatic. Kills, doesn't just suppress.
- Easy to use. Requires little or no agitation after initial mixing.
- Works well through chemigation systems.
- No known resistance after 20 years of use.
- Non-phytotoxic when used according to label directions.
- Long lasting, reducing number of applications needed.

PACKAGING

Unit Size:

1 qt.

Units Per Case: 4

STOCK #95044

(See next page
for label information)

Scotts-Sierra Crop Protection Company
14111 Scottslawn Road • Marysville, OH 43041

If you have any questions regarding the use of this product, call TOLL FREE 1-800-492-8255



Fungicide 25% Emulsifiable Concentrate

ACTIVE INGREDIENTS:	
Etridiazole†	25%
INERT INGREDIENTS.	75%
TOTAL	100%

†CAS #2593-15-9

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. HARMFUL IF SWALLOWED.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Flush eyes immediately with plenty of water for at least 15 minutes. **Get medical attention.**

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. **Call a physician immediately.**

IF INHALED: Induce coughing immediately.

IF ON SKIN: Wash thoroughly with soap and water. If irritation persists, **get medical attention.**

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

EPA Reg. No. 58185-8
Product of U.S.A.

EPA Est. No. 228-IL-1
NET WEIGHT: 1 GAL. (3.78 L)
(Four x 1 Qt. [946 ml] Containers)

© 2001, The Scotts Company. World rights reserved.

TRUBAN® is a registered trademark of Scotts-Sierra Crop Protection Company for its brand of Fungicide products.



Scotts-Sierra Crop Protection Company
14111 Scottslawn Road
Marysville, OH 43041
1-800-492-8255

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER! Corrosive! Causes eye damage and skin irritation. IF IN EYES, immediately flush with plenty of water for at least 15 minutes. **GET MEDICAL ATTENTION.** Do not get in eyes, on skin or clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of spray mist. In case of contact, immediately flush eyes or skin with plenty of water. If in eyes, see a physician. If skin irritation develops, get medical attention. Do not contaminate feed or food.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant gloves, such as barrier laminate or viton.
- Chemical-resistant footwear plus socks.
- Protective eyewear.
- Chemical-resistant headgear for overhead exposure.
- Chemical-resistant apron when cleaning equipment, mixing or loading.
- For exposures outdoors, a NIOSH approved respirator with any R, P, or HE filter.
- For exposures in enclosed areas, a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, P, or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

THIS PESTICIDE IS TOXIC TO FISH. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or Viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

CHEMIGATION

Refer to supplemental labeling entitled "Chemigation Information" for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

FOR PROFESSIONAL USE ONLY

TRUBAN is a soil fungicide for control of damping off, root and stem rot diseases caused by Pythium and Phytophthora. It may be used at time of seeding and transplanting as a drench for control of these problem fungi in greenhouses and outdoors.

TRUBAN suspensions in water should be occasionally agitated to assure uniform application of material. TRUBAN suspensions need not be protected from light. Use at recommended rates. Overdosage may result in toxicity to sensitive plants. If plants are suspect of being sensitive, apply fungicide to a small number to ensure safety. TRUBAN should not be mixed with other pesticides or nutrients.

Thoroughly rinse TRUBAN residues out of all equipment after use.

PLANT TYPE	REMARKS
BEDDING PLANTS Ajuga Larkspur Alyssum Marigold Calendula Pansy Celosia Petunia Cineraria Salvia Chrysanthemum Snapdragon Coleus Verbena Dahlia Vinca Dianthus Zinnia Dusty Miller	BEDDING PLANTS (soil 2" to 3" deep). Mix 3-4 oz. of TRUBAN 25% E.C. with 100 gals. of water apply to 800 sq. ft. of bed area (1 gal. per 8 sq. ft.) and use at time of seeding and transplanting. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 4 to 8 week intervals, if additional applications are necessary.
FOLIAGE PLANTS Aglaonema Hedera Anthurium Hoya Areca Palm Maranta Brassaia Nephrolepis Cactus Nephthytis Caladium Peperomia Calathea Pilea Chamaedorea Philodendron Chinese Evergreen Pothos Crassula Sansevieria Dieffenbachia Stromanthe Gynura Zygocactus	BED AND BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6" deep). Mix 4-8 oz. of TRUBAN 25% E.C. with 100 gals. of water and apply to 400 sq. ft. of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 2 to 3 month intervals to maintain disease control.
CONTAINER & BEDGROWN HERBACEOUS African Violet Gerbera Chrysanthemum Kalanchoe Easter Lily Poinsettia Geranium Sinningia WOODY Azalea Maple Boxwood Pieris Euonymus Podocarpus Ficus Red Pine Juniper Rhododendron Leucothoe	CONTAINER & BEDGROWN PLANTS Mix 4-8 oz. of TRUBAN 25% E.C. with 100 gals. of water per 400 sq. ft. or apply in sufficient amount to saturate the soil mixture, such as 1/2 pint per 6" pot. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 4 to 12 week intervals if necessary.

STORAGE AND DISPOSAL

STORAGE: Not sensitive to light. Stable at normal temperatures and humidities.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, State or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by incineration if allowed by State and local authorities.

**IMPORTANT NOTICE:
DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use on ornamentals. However, it is understood that tests may not have been carried out on all varieties and under all growing conditions. The user should always follow the label directions and exercise judgment and caution when using this product under their growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.

SPECIMEN LABEL