

**RESTRICTED USE PESTICIDE**  
**Toxic to Fish and Aquatic Organisms**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

# DISCIPLINE® CA

For use on beans, brassicas, caneberrries, canola, citrus, corn, cotton, crambe, curcubits, head lettuce, peas, peppers and rapeseed to control a range of insects and mites.

**ACTIVE INGREDIENT:**

**Bifenthrin**

(2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropanecarboxylate\* ..... 25.1%

**INERT INGREDIENTS\*\*** ..... 74.9%

**TOTAL** ..... 100.0%

Contains petroleum distillate

\*Cis isomers 97% minimum, trans isomers 3% maximum.

\*\*Contains xylene range aromatic solvents.

This product contains 2 pounds active ingredient per gallon.

**ALL WEATHER FORMULA**

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING - AVISO**

The product label must be in the possession of the user at the time of application.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID**

If swallowed:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Do not give any liquid to the person.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If inhaled:	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
If in eyes:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If on skin or clothing:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**EMERGENCY INFORMATION**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  
FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:  
Transportation: CHEMTREC.....1-800-424-9300  
Other: AMVAC.....1-323-264-3910

**NOTE TO PHYSICIAN**

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

EPA Reg. No. 5481-517

EPA Est. No. 5481-AL-1

Made in U.S.A.



4100 E. Washington Blvd.

Los Angeles, CA 90023, USA

1-323-264-3910 • www.amvac-chemical.com

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING:** May be fatal if swallowed. Harmful if inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

**Handlers who may be exposed to the dilute through application or other tasks must wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistance gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton.
- Shoes plus socks.

**Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistance gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton.
- Shoes plus socks, and
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Pest Resistance:** Pests have demonstrated the ability to develop resistance to continual use of the same product or to products with a similar mode of action. In order to avoid conditions which favor development of pest resistance, follow established resistance management guidelines in your geography as recommended by state agricultural extension personnel and crop consultants. If product performance diminishes and pest resistance is suspected to have developed in your geography, and specifically on your crop, repeated use of this product may not adequately control the target pest. As soon as practical, contact your local crop consultant, agricultural advisor, or company representative to obtain information on alternative pest management strategies in your geography.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Chemical-resistant gloves, such as Barrier Laminar or Nitrile Rubber or Neoprene Rubber or Viton, and
- Shoes plus socks.

### CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For Low Energy Precision Application (LEPA) irrigation a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment. Product should be applied continuously for the duration of the water application. Product should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

### Special Equipment

The registration of this product in California requires that the product be used in closed systems that meet the criteria for closed systems as established by the California Department of Food and Agriculture. The criteria and a list of the closed systems meeting the criteria are available through the California Department of Food and Agriculture.

### Rotational Crops

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of bifenthrin.

### Tank-Mixture

Product may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

### Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

In **New York State** this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

### Spray Drift Precautions

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made in cotton. Use of ultra low volume (ULV) application on corn and hops is prohibited.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid application when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Do not make aerial or ground applications to corn if heavy rainfall is imminent. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

COTTON		
PEST	DOSAGE	
	LB A/A	FL OZ/A
European Corn Borer Soybean (Banded) Thrips Tobacco Thrips	0.02-0.10	1.3-6.4
Boll Weevil Bollworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworm Fall Armyworm Lygus spp. Plant Bug Saltmarsh Caterpillar Southern Garden Leafhopper Stink Bug Tobacco Budworm Whitefly Yellow Striped Armyworm	0.04-0.10	2.6-6.4
Beet Armyworm Carmine Spider Mite Pink Bollworm Two-spotted Spider Mite	0.06-0.10	3.8-6.4

(CONTINUED)

REMARKS-Cotton		
Product may be applied in water or refined vegetable oil (soybean/cottonseed).		
<b>Application in Water:</b> Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.		
<b>To Control Boll Weevil:</b> Apply product at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels.		
<b>To Control Mites and Aphids:</b> Apply when pests first appear. Repeat as necessary to maintain control. Higher rates will be required once a damaging threshold is established.		
RESTRICTIONS-Cotton		
Do not apply more than 0.3 pound active per acre per season.		
Do not make more than 3 applications per season.		
Do not apply within 14 days of harvest.		
Do not graze livestock in treated areas or cut treated crops for feed.		
Do not make more than 10 synthetic pyrethroid applications (one of product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythron®, Capture®, Danitol®, Karate®, Mustang®, Pounce® and Scout X-TRA®.		

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (AT PLANT USE)**

PEST	DOSAGE			
	LB A/A	FL OZ/A		
Western Corn Rootworm Larvae	0.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row		
Cutworm spp. Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm or Armyworm spp. Wireworm	0.0023 to 0.0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row		
Row Spacings (inches)	40	38	36	30
Product (pounds ai per acre)	0.060	0.064	0.069	0.080
Product (formulated ounces per acre)	3.9	4.1	4.4	5.12

**REMARKS**

Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the product needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. Mix product with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of product, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Applications of this product alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of product and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.

**RESTRICTIONS**

Do not apply to soil where there is greater than 30% cover of crop residue remaining.

Do not apply within 30 days of harvest.

Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.

Do not apply more than 0.1 pound active per acre per season as an at plant application.

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR USE)**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids Beet Armyworm Chinch Bug Corn Earworm Corn Rootworm Adult Cucumber Beetle Adult Cutworm Species Fall Armyworm Flea Beetle Grasshoppers Greenbug Leafhoppers Stinkbugs Thrips True Armyworm and Armyworm spp. Webworms Yellow striped Armyworm	0.033-0.10	2.1-6.4
Banks Grass Mite Carmine Mite Two-spotted Spider Mite	0.08-0.10	5.12-6.4
Western Corn Rootworm Larvae	0.08-0.10 (Chemigation Only)	5.12-6.4 (Chemigation Only)

**REMARKS—Corn**

**General:** Apply in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**To control ear-attacking pests:** Apply this product just before silking and repeat as necessary to maintain control.

**For control of other insect pests:** Apply when pests first appear and repeat as necessary.

**For control of Mites:** Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.

For Two-spotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.

**To control Western Corn Rootworm Larvae by chemigation use only:** Apply at or closely following corn rootworm egg hatch in a minimum of 1 inch irrigation water per acre.

**RESTRICTIONS—Corn**

- Do not apply more than 0.2 pound active per acre per season.
- Do not apply within 30 days of harvest.
- Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
- Use of ultra low volume (ULV) application on corn is prohibited.
- Do not make aerial or ground applications to corn if heavy rainfall is imminent.
- Use of this product on corn is prohibited in all coastal counties.

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (PRE & PPI)**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Black Cutworm White Grub Wireworm Seedcorn Maggot Armyworm spp Common Stalk Borer	0.047-0.062 Pre-Plant incorporated (PPI)	3-4 Pre-Plant Incorporated (PPI)
Black Cutworm Armyworm spp Common Stalk Borer	0.040 Pre-Emergence (PRE)	2.56 Pre-Emergence (PRE)

**REMARKS**

The 3-4 oz/A rate must be pre-plant incorporated (PPI) and can be tank-mixed and applied with PPI herbicides. Incorporation of this product into the soil should be close to the intended seed planting depth and no deeper than 3 inches.

The 2.56 oz/A rate may be applied PRE and can be tank-mixed and applied with PRE herbicides.

**SWEET CORN  
SWEET CORN GROWN FOR SEED (AT PLANT USE)**

PEST	DOSAGE			
	LB A/A	FL OZ/A		
Western Corn Rootworm Larvae	0.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row		
Cutworm spp. Crubs Seed Corn Maggot True Armyworm and Armyworm spp. Wireworm	0.0023 to 0.0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row		
Row Spacings (inches)	40	38	36	30
Product (pounds ai per acre)	0.060	0.064	0.069	0.080
Product (formulated ounces per acre)	3.9	4.1	4.4	5.12

**REMARKS**

Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the product needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. Mix this product with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of product, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Applications of this product alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of product and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.

**RESTRICTIONS**

- Do not apply to soil where there is greater than 30% cover of crop residue remaining.
- Do not apply within 30 days of harvest.
- Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.
- Do not apply more than 0.1 pound active per acre per season as an at plant application.

SWEET CORN SWEET CORN GROWN FOR SEED (FOLIAR USE)		
PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids Army Cutworm Beet Armyworm Chinch Bug Corn Earworm Corn Rootworm Adult Cucumber Beetle Adult Cutworm Species Fall Armyworm Flea Beetle Grasshoppers Greenbug Leafhoppers Stinkbugs Thrips True Armyworm and Armyworm spp. Webworms Yellow striped Armyworm	0.033-0.10	2.1-6.4
Banks Grass Mite Carmine Mite Pacific Spider Mite Two-spotted Spider Mite	0.08-0.10	5.12-6.4
Western Corn Rootworm Larvae	0.08-0.10 (Chemigation Only)	5.12-6.4 (Chemigation Only)
<p><b>REMARKS—Sweet Corn</b></p> <p><b>General:</b> Apply in a minimum of 2 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p><b>To control ear-attacking pests:</b> Apply this product when silking begins and repeat as necessary to maintain control.</p> <p><b>For control of other insect pests:</b> Apply when pests first appear and repeat as necessary.</p> <p><b>For control of Mites:</b> Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.</p> <p>For Two-spotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.</p> <p>Higher rates will be necessary for heavier initial populations and corn under heat or drought stress.</p> <p><b>To control Western Corn Rootworm Larvae by chemigation use only:</b> Apply at or closely following corn rootworm egg hatch in a minimum of 1 inch irrigation water per acre.</p> <p><b>RESTRICTIONS—Sweet Corn</b></p> <p>Do not apply more than 0.2 pound active ingredient (12.8 ounces formulated) per acre per season.</p> <p>Do not apply within one day of harvest.</p> <p>Do not apply within 18 days of harvest for sweet corn that is hand harvested.</p> <p>Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application.</p> <p>Use of ultra low volume (ULV) application on corn is prohibited.</p> <p>Do not make aerial or ground applications to corn if heavy rainfall is imminent.</p> <p>Use of this product on corn is prohibited in all coastal counties.</p>		

CANOLA, CRAMBE, RAPESEED			
CROP	PEST	DOSAGE	
		LB A/A	FL OZ/A
Canola Crambe Rapeseed	Aphids Cutworms Diamondback Moth Loopers Other Lepidopterous Larvae Flea Beetle Flea Hopper Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly Armyworms	0.033-0.04	2.1-2.6
<p><b>REMARKS</b></p> <p>Apply this product in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p><b>RESTRICTIONS</b></p> <p>Do not apply more than 0.08 lb. active ingredient (5.12 ounces) per acre per season.</p> <p>Do not make applications less than 14 days apart.</p> <p>Do not apply within 35 days of harvest.</p>			

CUCURBITS			
CROP	PEST	DOSAGE	
		LB A/A	FL OZ/A
Chayote (fruit) Chinese waxgourd (Chinese preserving melon) Citrin melon Cucumber Gherkin Gourd, edible (includes hyotan, cucuzza) ( <i>Luffa spp.</i> ) (includes hechima, Chinese okra) ( <i>Momordica spp.</i> ) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber) Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i> ) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkin ( <i>Cucurbita spp.</i> ) Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini) Squash, winter (includes butternut squash, calabaza, hubard squash ( <i>C. mixta</i> ; <i>C. pepo</i> ) includes acorn squash, spaghetti squash) Watermelon (includes hybrids and/or varieties of <i>Citrullus spp.</i> )	Aphids Cutworm Cabbage Looper Leafhoppers Cucumber Beetle Squash Bug Melonworm Pickleworm Plant Bug Stink Bug Rindworm Squash Vine Borer Armyworm Corn Earworm Tobacco Budworm Grasshopper	0.04-0.10	2.6-6.4
	Whitefly Banks Grass Mite Two-spotted Spider Mite Carmine Mite Lygus spp.	0.08-0.10	5.12-6.4

(CONTINUED)

**REMARKS**

Apply this product in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**RESTRICTIONS**

Do not apply more than 0.3 lb. active ingredient (19.2 ounces formulated) per acre per season.  
Do not make more than two applications after bloom.  
Do not make applications less than 7 days apart.  
Do not apply within 3 days of harvest.

**LETTUCE, HEAD**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids Armyworm Cabbage Maggot Corn Earworm Cucumber Beetle Cutworm Diamondback Moth Flea Beetle Grasshoppers Imported Cabbageworm Leafhopper Looper Saltmarsh Caterpillar Stink Bug spp. Thrips Tobacco Budworm Whitefly	0.033-0.10	2.1-6.4
Lygus spp. Carmine Mite Two-spotted Spider Mite	0.08-0.10	5.12-6.4

**REMARKS**

Apply this product in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**RESTRICTIONS**

Do not make applications less than 7 days apart.  
A maximum of 0.5 lb. active ingredient may be applied per acre per season.  
Do not apply within 7 days of harvest.

**LIMA BEAN (GREEN)**

CROP	PEST	DOSAGE	
		LB A/A	FL OZ/A
Lima Bean (green)	Flea Beetle Leafhoppers	0.025-0.10	1.6-6.4
	Plant Bug Stink Bugs Corn Earworm Loopers Whitefly	0.033-0.10	2.1-6.4
	Banks Grass Mite Two-spotted Spider Mite Carmine Mite Lygus spp.	0.08-0.10	5.12-6.4

**REMARKS**

Apply this product in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**RESTRICTIONS**

Do not apply more than 0.2 lb. active ingredient (12.8 ounces) per acre per season.  
Do not apply within 3 day of harvest  
Do not apply within 9 days of harvest for peas/beans that are hand harvested.

BRASSICAS			
CROP	PEST	DOSAGE	
		LB A/A	FL OZ/A
Head and Stem Brassica Vegetables including: Broccoli Chinese Broccoli (gai lan, white flowering broccoli) Brussels Sprouts Cauliflower Cavalo broccolo Kohlrabi Cabbage Chinese Cabbage (napa)	Cutworms	0.033-0.10	2.1-6.4
	Corn Earworm		
	Saltmarsh Caterpillar		
	Flea Beetles		
	Imported Cabbageworm		
	Aphids		
	Whitefly		
	Armyworms		
	Loopers		
	Crickets		
	Diamondback Moth	0.08-0.10	5.12-6.4
	Lygus Spp.		

**REMARKS**

Apply this product in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**RESTRICTIONS**

Do not apply more than 0.5 lb. active ingredient (1 quart) per acre per season.  
Do not make more than 5 applications after bloom.  
Do not make applications less than 7 days apart.  
Do not apply within 7 days of harvest.

**CANEBERRIES**

CROP	PEST	DOSAGE	
		LB A/A	FL OZ/A
Caneberries including: Blackberries Bingberries Dewberries Loweberries Marionberries Ollalieberries Youngberries Loganberries Raspberries	Leafroller	0.05-0.10	3.2-6.4
	Orange Tortrix		
	Root Weevil	0.10	6.4
	Spider Mites		

**REMARKS**

Apply this product by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons per acre by air and 50 gallons per acre by ground). One application may be made pre-bloom and a second application may be made post-bloom.

**RESTRICTIONS**

Do not apply within 3 days of harvest.  
Do not apply within 4 days of harvest for cranberries that are hand-harvested

**PEPPERS, BELL AND NON-BELL**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids Armyworm Corn Earworm Cucumber Beetle Cutworm Flea Beetle Leafhopper Leafminer Looper Pepper Weevil Stinkbug spp. Thrips Tobacco Budworm Whitefly	0.033-0.10	2.1-6.4
Lygus spp. Carmine Mite Two-spotted Spider Mite		

(CONTINUED)

**REMARKS**

Apply this product using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.

**RESTRICTIONS**

Do not make applications less than 7 days apart.  
Do not apply more than 0.2 lb active ingredient per acre per season.  
Do not apply within 7 days of harvest.

**STORAGE AND DISPOSAL**

**Pesticide Storage**

Do not freeze or store below 40°F. If crystals are observed, gradually warm material to above 60°F by placing container in a warmer location. Gently shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Chemtrec 1-800-424-9300

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**Container Disposal**

Metal or Plastic Container: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

Returnable Product Container: Do not attempt to rinse returnable container, break, tamper with container seals or empty any remaining product contents from remaining product contents from container. Return container intact to point of purchase.

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

**THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSSES OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**

**BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT ALLOWED BY LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

Ammo, Mustang and Pounce are registered trademarks of FMC Corporation.

Karate is a registered trademark of Syngenta Group Company.

Asana is a registered trademark of E.I duPont de Nemours & Company.

Baythroid and Scout X-TR are registered trademarks of Bayer Aktiengesellschaft.

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Scout Xtra is a registered trademark of Bayer Aktiengesellschaft.

Discipline and Ambush are registered trademarks of AMVAC Chemical Corporation.





4100 E. Washington Boulevard  
Los Angeles, CA 90023, U.S.A.

1-323-264-3910 • [www.amvac-chemical.com](http://www.amvac-chemical.com)