

# Specimen Label



## Herbicide

®Trademark of Dow AgroSciences LLC

**For postemergent control of annual broadleaf weeds in wheat (including durum), barley, oats, rye and triticale.**

Group	2	4	HERBICIDES
-------	---	---	------------

Active Ingredient:

florasulam: N-(2,6-difluorophenyl)-8-fluoro-5-methoxy (1,2,4)triazolo(1,5-c)pyrimidine-2-sulfonamide.....	0.50%
fluroxypyr 1-methylheptyl ester: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic acid, 1-methylheptyl ester .....	14.45%
Other Ingredients .....	85.05%
Total .....	100.00%

EPA Reg. No. 62719-604

Acid equivalent – fluroxypyr – 10.03% - 0.833 lb/gal  
Contains 0.042 lb of florasulam per gallon.  
Contains petroleum distillates

## Keep Out of Reach of Children WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**Shake Well Before Use -- Avoid Freezing**

### Precautionary Statements

**Hazards to Humans and Domestic Animals**

## WARNING

**Causes Substantial But Temporary Eye Injury • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals**

**Do not get in eyes or on clothing. Wear protective eyewear.**

## Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves  $\geq$ 14 mils (such as barrier laminate, butyl rubber, nitrile rubber and viton)
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**Note to Physician:** Contains petroleum distillate – vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

## Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

## Physical or Chemical Hazards

**Combustible.** Do not use or store near heat or open flame.

## Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical-resistant gloves  $\geq$ 14 mils (such as barrier laminate, butyl rubber, nitrile rubber and viton)
- Shoes plus socks
- Protective eyewear

## Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Storage below 14°F will cause the product to freeze. If product freezes, bring to room temperature and agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Refillable containers 5 gallons or larger:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Nonrefillable containers 5 gallons or larger:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## General Information

Use Starane® Flex herbicide as a postemergence herbicide for the control of annual broadleaf weeds such as kochia, chickweed, wild buckwheat, mustards and cleavers in wheat (including spring, winter and durum), barley, oats, rye and triticale not underseeded with legumes.

Starane Flex rapidly stops growth of susceptible weeds. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 2 weeks after application depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent upon weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

## Use Precautions and Restrictions

When applying this product in tank mix combination, follow all applicable use directions, precautions, and limitations on each manufacturer's label.

**Chemigation:** Do not apply this product through any type of irrigation system.

Do not apply Starane Flex directly to, or otherwise permit it to come into direct contact with, susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, or tobacco. Do not permit spray mists containing Starane Flex to drift onto such plants.

For best resistance management stewardship, do not use more than once per season or use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Do not apply to crops underseeded with legumes.

## Crop Rotation Intervals

The following rotational crops may be planted at the indicated interval following application of Starane Flex.

Numbers in parentheses (-) refer to Specific Crop Rotation Information.

Crop	Rotation Interval (1) (Months)
barley, oats, wheat, grasses	0
field corn, popcorn, seed corn, sweet corn, sorghum	3
alfalfa, canola, chickpea, soybean, dry bean, peas (dry and succulent), flax, lentil, potato, safflower, sugar beet, sunflower	9
other crops not listed	12

### Specific Crop Rotation Information:

1. Minimum number of months that must pass before planting other crops after application of Starane Flex.

## Avoid Injurious Spray Drift

This product can affect broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Do not apply Starane Flex directly to, or allow spray drift to come into contact with, broadleaf crops including alfalfa, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See Crop Rotation Intervals section.)

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure crops, whether dormant or actively growing. When applying Starane Flex, use low pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets. Under adverse weather conditions, fine spray droplets that do not settle rapidly onto target vegetation may be carried a considerable distance from the treatment area. A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use directions and precautions on the product label.

**Ground Applications:** To minimize spray drift, apply Starane Flex in a total spray volume of 8 gallons or more per acre using spray equipment designed to produce large droplet, low pressure sprays. Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Spot treatments should be applied only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine droplet spray.

**Aerial Application:** To minimize spray drift, apply Starane Flex in a total spray volume of 3 gallons or more per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high potential for temperature inversion. Spray drift from aerial application can be minimized by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom that does not exceed 75% of wingspan or 90% of rotor diameter. Spray pattern and droplet size distribution can be evaluated by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Do not apply under conditions of a low level air temperature inversion. A temperature inversion is characterized by little or no wind and air temperature that is lower near the ground than at higher levels. The behavior of smoke generated by an aircraft-mounted device or continuous

smoke column released at or near site of application will indicate the direction and velocity of air movement. A temperature inversion is indicated by layering of smoke at some level above the ground and little or no lateral movement.

### Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

- The distance of the outer most operating nozzles on the boom must not exceed 75% of wingspan or 90% of rotor diameter.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory. (This information is advisory in nature and does not supersede mandatory label requirements.)

### Aerial Drift Reduction Advisory

**Information on Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

### Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Boom Length:** For some use patterns, reducing the effective boom length to less than 75% of the wingspan or 90% of rotor length may further reduce drift without reducing swath width.

**Application Height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke

generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

### Mixing Directions

#### Starane Flex - Alone

1. Fill the tank with 1/2 of the total amount of water.
2. Start agitation.
3. Add the required amount of Starane Flex.
4. Continue agitation while filling the spray tank to the required volume.
5. To ensure a uniform spray mixture, continuous agitation is required during application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

#### Starane Flex - Tank Mix

If a broader spectrum of weed control is needed, Starane Flex may be tank mixed with labeled rates of other herbicides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

#### Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed specified application rates for respective products or maximum allowable application rates for any active ingredient in the tank mix.
- Do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment have been adequately cleaned. (See Equipment Clean-Out Procedures.)
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** A jar test is recommended prior to tank mixing to ensure compatibility of Starane Flex and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

#### Mixing Order for Tank Mixes:

1. Fill the spray tank to 3/4 of the total spray volume required with water.
2. Start agitation.
3. Add Starane Flex and agitate for 2 to 3 minutes
4. After adding Starane Flex, add different formulation types in the following order: (1) dry flowables; (2) wettable powders; (3) aqueous suspensions, flowables and liquids. Maintain agitation and add: (4) emulsifiable concentrates; (5) solutions; and (6) adjuvants. Allow time for complete mixing and dispersion after each addition.
5. Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

#### Clean-Out Procedures for Spray Equipment

1. Drain any remaining spray mixture from the application equipment.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add household ammonia at a rate of 1 gallon per 100 gallons of water. Recirculate for 5 minutes and spray out part of this mixture for 5 minutes through the boom. Drain tank.
4. Remove all spray nozzles and screens and clean separately.
5. If spray equipment will be used for pesticide application to crops sensitive to Starane Flex, steps 1 through 3 should be repeated. Exterior surfaces of spray equipment should also be thoroughly cleaned.

**Note:** Rinsate may be disposed of on site according to label use directions or at an approved waste disposal facility.

## Weeds Controlled or Suppressed

### Weeds Controlled

buckwheat, wild	mustard, black
canola, volunteer <sup>1</sup>	mustard, blue (purple)
chamomile, corn	mustard, tansy
chamomile, false (scentless)	mustard, treacle (bushy wallflower)
chamomile, mayweed (dogfennel)	mustard, tumble (Jim Hill)
chamomile, wild	mustard, wild
chickweed, common	pennycress, field
cleavers	pineappleweed
flax, volunteer	prickly lettuce
flixweed	puncturevine
knotweed	purslane, common
kochia <sup>2</sup>	ragweed, common
London rocket	shepherdspurse
mallow, Venice	smartweed
	(green, ladysthumb, Pennsylvania)
morningglory	sunflower

### Weeds Suppressed

bindweed, field	marshelder
canola, volunteer <sup>3</sup>	nightshade species
dandelion	pigweed, redroot
hempsnettle	sowthistle, annual
horseweed (marestail)	sowthistle, perennial
mallow, common	thistle, Russian

<sup>1</sup>Including herbicide-tolerant canola varieties except Clearfield (imidazolinone-tolerant) canola.

<sup>2</sup>Includes herbicide tolerant or resistant biotypes.

<sup>3</sup>Clearfield (imidazolinone-tolerant) canola.

## Resistance Management

### Management of Kochia Biotypes

Research has suggested that many biotypes of kochia can occur within a single field. Application of Starane Flex at rates below 13.5 fl oz per acre can result in a shift to more tolerant biotypes within a field.

### Best Resistance Management Practice

Extensive populations of dicamba tolerant kochia have been identified in certain small grain and corn production regions (such as Chouteau, Fergus, Liberty, Toole, and Treasure counties in the state of Montana). In these areas, Starane Flex should be rotated with products **that do not contain dicamba** to minimize selection pressure. Unless recommended by Dow AgroSciences, do not use less than 13.5 fl oz of Starane Flex per acre for control of ALS-resistant biotypes of kochia.

## Application Directions

### Application Timing

Apply Starane Flex early postemergence to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds that have emerged at the time of application will be controlled.** If foliage is wet at the time of application, control may be decreased. Applications of Starane Flex are rainfast within 4 hours after application.

### Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Do not broadcast apply in less than 3 gallons of total spray volume per acre. For best results and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As vegetative canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under Avoid Injurious Spray Drift.

### Adjuvants

Generally, this product does not require the use of an adjuvant to achieve satisfactory weed control. An adjuvant may be added to optimize herbicidal activity when applications are made at lower use rates or lower carrier volumes; under conditions of cool temperature, low relative humidity or drought; to small, heavily pubescent kochia; or when a tank mix partner recommends the use of an adjuvant. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

## Wheat (Including Durum), Barley, Oats, Rye, Triticale

Apply 13.5 fl oz of Starane Flex per acre to actively growing wheat (including spring, winter and durum), barley, oats, rye and triticale from the 3-leaf crop growth stage up to flag leaf emergence (Zadoks scale 39). Apply when susceptible broadleaf weed seedlings and kochia are actively growing and less than 4 inches tall. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from application made to seedling weeds.

Warm, moist growing conditions promote active weed growth and enhance the activity of Starane Flex by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

**Tank Mixes for Wheat (Including Durum):** Starane Flex may be applied in tank mix combination with labeled rates of other products registered for postemergence application in wheat. See Tank Mixing Precautions under Mixing Directions. When tank mixing, do not exceed specified application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

### Restrictions:

- **Preharvest Interval:** Do not apply within 60 days of crop harvest.
- Do not apply more than 13.5 fl oz of Starane Flex per acre per growing season.
- Livestock may be grazed on treated crops 7 days following application.

---

## Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

## Warranty Disclaimer

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

## Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

## Limitation of Remedies

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

Label Code: D02-390-001  
Initial printing  
LOES Number: 010-02240  
EPA accepted 07/06/09