

- Water-based fogging concentrate
- Effective and fast acting flushing action, knockdown and kill of insects

ACTIVE INGREDIENTS:

Pyrethrins.....	1.00%
* Piperonyl butoxide, Technical	2.00%
** N-Octyl bicycloheptene dicarboximide.....	3.00%
OTHER INGREDIENTS	94.00%
	100.00%

* Equivalent to 1.6% (butylcarbityl) (6-propylpiperonyl) ether and 0.4% related compounds

** MGK® 264, Insecticide Synergist



KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE

GENERAL: This concentrate is designed for use full strength or diluted as desired, according to the dilution table listed below. Achieve optimum performance using a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply as a space, area or contact spray in residential and commercial structures and animal housing.

Do not apply directly to food. Do not allow spray to contact stored food or food contact surfaces. Do not apply as a space spray while food processing is underway. Cover or remove foods before treatment. Cover or wash all food processing surfaces and equipment with an effective cleaning compound followed by a potable water rinse. In the home cover all food processing surfaces, dishes, and utensils before treatment or thorough wash before use following treatment.

Avoid contamination of food or feedstuffs. Remove pets and birds and cover fish aquariums and turn off aquarium air pump motor before spraying.

In health care facilities: Remove patients from area to be treated prior to treatment. Ventilate treated area for two hours after spraying. Do not return patients to treated areas until ventilation is complete. When used as a coarse wet spray, the sprayed surfaces must be dried before patients are allowed to reoccupy treated areas.

NOTE: Some wood and plastic finishes can be marred or damaged by insecticide applications. To avoid damage, cover those items and do not treat directly. If in doubt, test by applying this material to an inconspicuous area or surface and evaluate before application.

NOTE: If there is a need to treat such areas as conduits, motor housings, junction and switch boxes or other electrical equipment, turn off electrical equipment AND shut off power to the area prior to application. After application, allow time for insecticide to dry before restoring power to the area, starting motors or placing switches in the "ON" position.

AREAS for use in and around:

Animal Areas and Quarters	Food Storage Facilities	Peanut Storage Warehouses
Animal Shelters	Freight Containers	Pool Halls
Apartments	Fruit Packing Sheds	Poultry Houses
Arcades	Funeral Parlors	Poultry Processing Plants
Athletic Fields	Garages	Prisons
Attics	Garbage Bins	Public Buildings
Auditoriums	Garbage Cans	Public Storages
Bakeries	Garbage Compactors	Rabbit Housing
Bars	Garbage Trucks	Rabbit Processing Plants
Basements	Gas Stations	Railroad Cars
Bathrooms	Goat Houses	Recreational Vehicles
Beverage Plants	Grain Elevators	Research Animal Quarters
Boats	Grain Harvesting and Handling Equipment	Restaurants
Bottling Plants	Grain Mills	Restrooms
Brandy Storage Warehouses	Granaries	Retail Stores
Breweries	Grocery Stores	Rice Mills
Brewery Warehouses	Grocery Warehouses	Schools
Buildings	Healthcare Facilities	Shade Houses
Buses	Homes	Ship Holds
Cabinets	Horse Stables and Barns	Ships
Cabins	Horse Trailers	Smoke Shops
Cafeterias	Hospices	Sports Facilities
Campers	Hospitals	Sports Stadiums
Canneries	Hotels	Stable Bins
Cargo Holds	Indoor Eating Establishments	Stables
Cattle Barns	Industrial Installations	Storage Bins (areas)
Churches	Institutional Dining Areas	Stored Food in Multi-walled Paper or Cloth Bags
Circus Tents	Institutions	Stored Seed Warehouses
City Missions	Jails	Stores
Closets	Kennels	Supermarkets
Club Houses	Laboratories	Swine Housing
Commercial Buildings	Laundromats	Taverns
Communication Centers	Lavatories	Textile Mills and Warehouses
Concession Stands	Libraries	Theaters
Condominiums	Liquor Stores	Tobacco Processing Facilities
Correctional Facilities	Livestock Barns	Tobacco Warehouses
Court Houses	Loafing Sheds	Tool Sheds
Crawl Spaces	Locker Rooms	Townhouses
Crematoriums	Marinas	Trains
Dairy and Livestock Barns	Mausoleums	Transportation Equipments
Day Care Centers	Meat Packing Plants	Truck Trailers
Delicatessens	Meat Rendering Plants	Trucks
Departments Stores	Mess Halls	Tunnels, Utility
Diners	Milk Rooms	Utility Rooms
Doghouses	Mills	Vegetable Packing Facilities
Drains and Sewers	Mini-storages	Vending Machines
Dried Fruit Processing Plants	Missions	Veterinary Hospitals
Dried Fruit Warehouses	Mobile Canteens	Wall and Equipment Voids (attics, walls, ceilings)
Drugstores	Morgues	Warehouses
Dumpsters	Mortuaries	Washrooms
Dwellings	Motels	Wholesale Stores
Egg Processing Plants	Movie Theaters	Wine Cellars
Factories	Museums	Wine Storage Warehouses
False Ceilings	Mushroom Processing Plants	Wineries
Federally Inspected Meat and Poultry Establishments	Nursing Homes	Zoos
Flour Mills	Office Buildings	
Food Handling Establishments	Offices	
Food Processing Plants	Other Public Buildings	
	Parking Garages	
	Passenger Rail Cars	
	Peanut Processing Plants	

FOR THE CONTROL OF INSECTS including:

Almond Moths	Drugstore Beetles	Mites
American Black Flour Beetles	Earwigs	Mosquitoes
Angoumois Grain Moths	Elm Leaf Beetles	Mushroom Flies
Ants	European Grain Moths	Palmetto Bugs
Barn Flies	Face Flies	Pillbugs
Bean Weevils	Firebrats	Pink Scavenger Caterpillars
Bed Bugs	Flat Grain Beetles	Red Flour Beetles
Bees	Fleas	Red Horned Grain Beetles
Beetles	Flies	Beetles
Black Carpet Beetles	Flying Moths	Rice Flour Beetles
Black Fungus Beetles	Foreign Grain Beetles	Rice Moths
Blow Flies	Fruit Flies	Rusty Grain Beetles
Book Lice	Fungus Beetles	Saw-toothed Grain Beetles
Bottle Flies	Fungus Gnats	Beetles
Boxelder Bugs	Gnats	Scorpions
Broad-horned Flour Beetles	Grain Mites	Sheep Ticks
Brown Dog Ticks	Granary Weevils	Silverfish
Brown Spider Beetles	Grasshoppers	Skipper Flies
Cadelles	Ground Beetles	Slender-horned Flour Beetles
Carpet Beetles	Hairy Fungus Beetles	Small Flying Moths
Centipedes	Hairy Spider Beetles	Small-eyed Flour Beetles
Cereal Beetles	Hide Beetles	Sowbugs
Cheese Mites	Horn Flies	Spider Beetles
Cheese Skippers	Hornets	Spiders
Chocolate Moths	House Flies	Spinous Ear Ticks
Cigarette Beetles	Indian Meal Moths	Square-necked Grain Beetles
Clothes Moths	Khapra Beetles	Stable Flies
Clover Mites	Large Black Flour Beetles	Tobacco Moths
Cluster Flies	Larger Grain Borers	Trogoderma Beetles
Cockroaches	Lesser Grain Borers	Two Banded Beetles
Cocoa Bean Moths	Lesser Houseflies	Vinegar Flies
Coffee Bean Weevils	Lesser Mealworms	Warehouse Beetles
Confused Flour Beetles	Longheaded Flour Beetles	Wasps
Corn Sap Beetles	Beetles	Waterbugs
Crickets	Maize Weevils	Webbing Clothes Moths
Dark Mealworms	Meal Worms	Weevils
Darkling Beetles	Mediterranean Flour Moths	White Marked Spider Beetles
Depressed Flour Beetles	Merchant Grain Beetles	Yellow Jackets
Dermestid Beetles	Mexican Grain Beetles	Yellow Mealworms
Drain Flies	Midges	
Dried Fruit Beetles	Miller Moths	
	Millipedes	

APPLICATION RATES

Type of pest	Method of application	Recommended rate
Flying insects and related pests	As a ULV, fog or space spray	1/2 to 1 fluid ounce of concentrate per 1,000 cubic feet of space
	As a crack and crevice injection	1/2 to 1 fluid ounce of concentrate per 1,000 cubic feet of space
Crawling insects and related pests	As a ULV, fog or space spray	1 fluid ounce per 1,000 cubic feet of space
	As a contact, surface or crack and crevice spray	6 to 32 fluid ounces in one gallon of water per 1,000 square feet of surface area
	As a crack and crevice injection	1 fluid ounce per 1,000 cubic feet of space

DILUTION TABLE

Application method	% pyrethrins in solution	Mixture	Application rate	Lbs. pyrethrins
ULV, fog or space spray	1.00%	Use full strength/undiluted	1 fluid ounce per 1,000 cubic feet of space	0.0006 lbs. per 1,000 cubic feet
	0.50%	1 part concentrate to 1 part water		0.00032 lbs. per 1,000 cubic feet
Contact, surface or crack and crevice spray	0.25%	32 fluid ounces of concentrate in one gallon of water	1 gallon of solution per 1,000 square feet of surface area	0.02 lbs. per 1,000 square feet
	0.15%	18 fluid ounces of concentrate in one gallon of water		0.012 lbs. per 1,000 square feet
	0.10%	12 fluid ounces of concentrate in one gallon of water		0.008 lbs. per 1,000 square feet
	0.05%	6 fluid ounces of concentrate in one gallon of water		0.004 lbs. per 1,000 square feet
Crack and crevice injection	1.00%	Use full strength/undiluted	1 fluid ounce per 1,000 cubic feet of space	0.0006 lbs. per 1,000 cubic feet
	0.50%	1 part concentrate to 1 part water		0.00032 lbs. per 1,000 cubic feet

SPACE SPRAY APPLICATIONS

General instructions for use as a space application in listed areas: Close doors, windows and shut off all ventilating equipment. Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply at the rate of 1/2 to 1 fluid ounce of undiluted concentrate per 1,000 cubic feet of space, or dilute with water according to the dilution table. Direct spray toward upper corners of room and ceiling. Do not remain in treated areas after application. Leave room closed for at least 1/2 hour. Ventilate thoroughly before occupants are allowed to re-enter. Repeat treatment when reinfestation occurs. Do not use equipment designed for deep void or crack and crevice injection for space spray applications in accessible areas.

Do not apply directly to food. Do not allow spray to contact stored food. Do not apply as a space spray while food processing is underway. Foods must be removed or covered during treatment. All food processing surfaces and equipment must be covered or washed with an effective cleaning compound followed by a potable water rinse. In the home all food processing surfaces, dishes and utensils must be covered during treatment or thoroughly washed before use. Food processing operations may continue when this product is applied as a wet spray as a spot, or crack and crevice treatment, with care and in accordance with the directions and precautions listed on this label.

When applying as a space spray to kill flying insects: Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Following the general and general space spray instructions on the label, apply undiluted at a rate of 1/2 to 1 fluid ounce per 1,000 cubic feet. If diluted solution is preferred, mix 1 part concentrate to 1 part water and apply at the rate of 1 fluid ounce of diluted solution per 1,000 cubic feet of space.

When applying as a space spray to kill crawling insects: Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Following the general use and general space spray instructions on the label, treat as a space spray. First, treat cracks and crevices in woodwork, walls, floors, behind and beneath all equipment, cupboards, obstructions or dark harborage areas, underneath sinks, behind pipes, under pallets and in all places that harbor insects. Contact as many insects as possible with the application. Then, apply undiluted concentrate at the rate of 1 fluid ounce per 1,000 cubic feet of space.

Dried fruit products in storage or processing: Following the general use and general space spray instructions, do not apply to fruit directly. Apply one fluid ounce of concentrate, or diluted concentrate, per 1,000 cubic feet of space. Disperse this product as a fine mist in the air above the trays and shelves. Repeat treatment when reinfestation occurs.

For crack and crevice injection: Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Always follow equipment manufacturer's instructions for proper equipment set up and operating pressures. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic areas and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids.

CONTACT OR SURFACE SPRAY APPLICATIONS

General instructions for use as a contact or surface spray: To kill accessible stages of listed insects, mix 6 to 32 fluid ounces of concentrate with water to make one gallon of spray solution. Using a compressed air sprayer or similar type of sprayer, apply as a coarse, wet spray. Apply 1 gallon of spray solution mixture per 1,000 square feet of surface area. To kill listed insects, spray thoroughly into all cracks, molding, crevices, walls, floors and other surfaces, storage and handling areas. Treat areas where insects hide, travel or rest, such as around drains, sinks, under pallets, wall voids, nests, ant trails, water pipes, around and behind heavy machinery and equipment. Contact as many insects as possible. Leave room ventilation or air conditioning equipment on during application. Food processing operations may continue when this mixture is applied as a spot or crack and crevice treatment, with care and in accordance with the directions and precautions listed on this label.

For use in transit protection: Do not use in aircraft. To kill accessible stages of listed insects, mix 6 to 32 fluid ounces of concentrate with water to make one gallon of spray solution. Using a compressed air sprayer or similar type of sprayer, apply as a coarse, wet spray. Apply 1 gallon of spray solution mixture per 1,000 square feet of surface area. Clean and spray truck beds, boxcars, ships holds and other listed transit equipment before loading. Thoroughly treat walls and floors. Spray each layer of cartons or containers lightly as they are stacked for transit.

For use on stored grain products: To assist in the kill of insects in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of insects. In case of severe infestations of flying moths, break up webbing before spraying and make a second application in two weeks.

ANIMAL HOUSING APPLICATIONS

General instructions for use in and around animal housing and related facilities: For use in and around listed livestock areas to kill accessible stages of listed insects. Do not apply spray directly to animals that will be used for milk or meat production. If facilities are to be used for milk and/or meat animals, remove animals before application. Milk and/or meat animals may re-occupy the facility once the insecticide has dried. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying.

As a space spray in animal housing: Following the general and general animal housing instructions, use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply at the rate of 1/2 to 1 fluid ounce of undiluted concentrate per 1,000 cubic feet of space above the animals, or dilute with water according to the dilution table. Direct application above the animals, towards upper portions of the enclosure, filling the room with mist. Repeat treatment when reinfestation occurs.

As a contact or surface spray in animal housing: To kill accessible stages of listed insects, mix 6 to 18 fluid ounces of concentrate with water to make one gallon of spray solution. Following the general and general animal housing instructions use a compressed air sprayer, or similar type of sprayer, to apply solution as a coarse, wetting spray. Apply approximately 1 gallon of spray solution mixture per 1,000 square feet of surface area. Spray walls, cracks and crevices, foundation, rafters, floor and any area where insects hide or rest. Animals may re-occupy the facility once the insecticide has dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, feed, seed, fertilizers, and veterinary supplies. Place formulations on lower shelves and dry formulations above.

PESTICIDE DISPOSAL: To avoid wastes, use all materials in the containers by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows (for containers of 5 Gallons [18.9 L] or less): Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your state water board or regional office of the EPA.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or a doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.



EPA Reg. No. 1021-1851
EPA Est. No. 1021-MN-2
0311-0911



All other marks are property of their respective holders. Manufactured by MGK®.
©2011 MGK. All rights reserved.
8810 Tenth Avenue North, Minneapolis, MN 55427

2.5M-1111