


STOLLER® MICRO-PLUS Manganese

Specimen Label



GUARANTEED ANALYSIS

Manganese (Mn):.....

6% Chelated Manganese

Derived from manganese chloride chelated with citric acid Contains EDTA chelating agent

GENERAL INFORMATION

MICRO- PLUS Liquid Chelates are for soil and foliar* application. They are readily available and rapidly absorbed by the root system or foliage.

MICRO-PLUS Liquid Chelates are formulated to be used in Stoller liquid fertilizers (NITRO- PLUS, URA- CAL, FOLIZYME) and Nitrogen solutions. When used with polyphosphate containing fertilizers, dilute with nitrogen solution before addition. Broadcast or row applications are acceptable, although greater efficiency may be expected from row applications. Micronutrient deficiencies usually occur when plants are small; therefore, micronutrients should be added to the fertilizer at or before planting.

STORAGE INSTRUCTIONS

MICRO- PLUS Liquid Chelates are chelates with medium corrositivity. It is recommended that MICRO- PLUS Liquid Chelates (10% Zn, 5% Fe, 5% Mn, 5% Cu) be stored in fiberglass, polyethylene or polyolefin. We recommend storing pails a maximum of three pails high. Do not stack pallets.

APPLICATION INSTRUCTIONS

Soil application of MICRO-PLUS MANGANESE can be incorporated in liquid fertilizer or applied separately. MICRO-PLUS MANGANESE can be coated on dry blend fertilizer to control dust in the blend plant and supply needed micronutrients.

For best results follow soil test and plant analysis recommendations. In absence of a soil test use the following as a guide:

Soil applications:

Maintenance: 1-2 quart per acre Moderate Deficiency: 2-4 quart per acre **Heavy Deficiency:** 4-6 quart per acre

Foliar applications

Row crops 1 qt per acre every 2-3 weeks with a

maximum of 4 applications including but not limited to corn, wheat, soybean

1 qt per acre every 10-14 days with a Vegetable crops maximum of 4 applications including

but not limited to: beans, broccoli, cauliflower, brussels sprouts, cabbage, carrot, spinach, lettuce, potato, onion

and tomato.

Fruit and Nut trees 1 qt per acre beginning when fully

leafed out, repeat 3-4 times. Apply 1 qt per acre as needed. Do not

Blueberries, Caneberries,

Cranberries and Strawberries:

apply after fruit set.

Forage Crops 1 qt per acre as needed.

CAUTION: KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat and/ or skin. Avoid contact with eyes, skin and clothing. **IF SWALLOWED**: Induce vomiting by sticking finger down throat. Never give anything to an unconscious person. Get medical attention. IF IN EYES: Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officers of Seller, is authorized to make any warranty, quarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product limited to replacement of product or refund of purchase price.

*Do not exceed one quart per acre for foliar sprays per application.