

This is a specimen label, intended for use only as a guide in providing general information regarding use of this product. As labels are subject to revision, always carefully read and follow the label on the product container.

Chlorsulf E-Pro 75 WDG

Herbicide

Chlorsulf E-Pro 75 WDG
Herbicide contains chlorsulfuron,
the active ingredient used in Telar®.

DRY FLOWABLE

| ACTIVE INGREDIENT: | By Weight |
|--|-----------|
| Chlorsulfuron: 2-Chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl] benzenesulfonamide..... | 75.0% |
| OTHER INGREDIENTS: | 25.0% |
| TOTAL: | 100.0% |

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 79676-72
EPA Est No. 082694-DEU-001

Manufactured for:
Etigra
501 Cascade Pointe Lane, Suite 103
Cary, NC 27513
REV 0409



| FIRST AID | |
|---|--|
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information. | |

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and other handlers" and have such PPE immediately available for use in an emergency such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or wastes.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid over-filling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Only use Chlorsulf E-Pro 75 WDG Herbicide in accordance with directions on this label or in separately published Etigra supplemental labeling.

Etigra will not be responsible for losses or damages resulting from the use of this product in any manner not specifically approved by Etigra.

Do not apply this product through any type of irrigation system.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Use on non-crop sites is not within the scope of the Worker Protection Standard.

Do not enter or allow entry into treated areas until sprays have dried.

GENERAL INFORMATION

Chlorsulf E-Pro 75 WDG Herbicide contains the active ingredient chlorsulf which is a herbicide used for control of many broadleaf weeds found in non-crop industrial sites (including industrial (unimproved) turf and for growth suppression and seedhead inhibition of established desirable grasses). Some non-crop industrial sites include airports, fence rows, government and private lands, military installations, petroleum tank farms, pipeline and utility rights-of-way, plant sites, pumping installations, railroads, roadsides and associated rights-of-way, and storage areas.

Some of these sites may contain temporary pools of surface water as a result of site management. Chlorsulf E-Pro 75 WDG Herbicide may be used to treat intermittent drainage, intermittently flooded low lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. In addition, Chlorsulf E-Pro 75 WDG Herbicide may be applied to bogs, marshes, and swamps after water has receded and to seasonally dry flood deltas. DO NOT make applications to natural or man-made bodies of water such as canals, lakes, ponds, reservoirs, and streams.

Both preemergent and postemergent applications of Chlorsulf E-Pro 75 WDG Herbicide will control weeds although several factors (including use rate, weed growth stage at the time of application, and post-application weather conditions) will affect the range of weeds controlled and the length of residual activity. **Annual weeds** are best controlled from application of Chlorsulf E-Pro 75 WDG Herbicide in the early stages of weed development. **Perennial weeds** are best controlled from application of Chlorsulf E-Pro 75 WDG Herbicide when weeds are in the bud to bloom or fall rosette stage.

ENVIRONMENTAL CONDITIONS AND ACTIVATION OF Chlorsulf E-Pro 75 WDG Herbicide

Chlorsulf E-Pro 75 WDG Herbicide moves into plants by absorption through the roots and foliage and rapidly inhibits the growth of susceptible weeds. Within two to three weeks after application, the weed growth slows and the new growth changes to a red-purple color. By four to six weeks after application, discoloration of the leaf veins and leaves is apparent, and growing points subsequently die.

For optimum control of target weeds, Chlorsulf E-Pro 75 WDG Herbicide needs to reach the weed roots. Rainfall or irrigation after an application moves the Chlorsulf E-Pro 75 WDG Herbicide into the soil and the weed root zone. Under cold, dry conditions movement of Chlorsulf E-Pro 75 WDG Herbicide into the root zone will be delayed. Chlorsulf E-Pro 75 WDG Herbicide is less effective to weeds hardened off by cold weather or under stress from lack of water.

Under most normal conditions, Chlorsulf E-Pro 75 WDG Herbicide will not harm labeled desirable grasses. Injury may result from application of Chlorsulf E-Pro 75 WDG Herbicide to grasses that are growing under stress (due to extreme temperatures or moisture, abnormal soil conditions, or cultural practices) or to certain sensitive species of grass.

RESISTANCE MANAGEMENT

Any weed population may contain or develop plants naturally resistant to herbicides with the same mode of action. The resistant biotypes may dominate the weed population if herbicides with the same mode of action are used repeatedly in the same field, and adequate control of these resistant weeds cannot be expected. Should an application not control the target weeds, retreat the area using an herbicide with a different mode of action.

To delay herbicide resistance, follow resistance management strategies such as:

- Rotation of Chlorsulf E-Pro 75 WDG Herbicide with herbicides having different modes of action to treat the same weeds.
- Application of tank mixes of herbicides with different modes of action, when such use is permitted.
- Use of herbicides as part of an IPM program.
- Prevention of resistant weed seed movement to other fields by cleaning harvesting and tillage equipment, and planting clean seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension specialist for specific alternative cultural practices or herbicide alternatives available in your area.

INTEGRATED PEST MANAGEMENT

Chlorsulf E-Pro 75 WDG Herbicide may be used as part of an Integrated Pest Management (IPM) program. This program relies on tillage (or other mechanical), biological, cultural, and chemical control practices to prevent economic pest damage. IPM principles and practices include field monitoring, historical information related to herbicide use and crop rotation, correct identification of target pests, population monitoring, and treatment when target pest populations reach a locally-determined action threshold. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine the action treatment threshold levels for treating specific pest/crop systems in your area.

IMPORTANT PRECAUTIONS AND RESTRICTIONS

Read the following restrictions and precautions to avoid injury to or loss of desirable trees or other desirable plants or vegetation.

- To avoid severe injury or death, do not drain or flush equipment rinses on or near desirable trees or other plants, on areas where their roots may extend, or in areas where the product may be washed or moved into contact with desirable plant roots.
- To minimize off-site movement of product on treated soils which can lead to damage of susceptible crops, do not apply if soils are powdery, dry or light, or sandy and if rainfall is not expected soon after treatment. Treated soil particles may move off-site to non-target crop sites through wind or water. Low levels of Chlorsulf E-Pro 75 WDG Herbicide may injure or kill most crops (except small grains), especially when crops are irrigated.
- The following conditions should be avoided during application to prevent runoff and movement of Chlorsulf E-Pro 75 WDG Herbicide residues: periods of intense rainfall, soils already saturated with water, asphalt or concrete paved surfaces, frozen soils or soils through which rain water will not readily penetrate. Do not disturb treated soils to minimize the potential for Chlorsulf E-Pro 75 WDG Herbicide movement by soil erosion from wind or water.
- Before using Chlorsulf E-Pro 75 WDG Herbicide, consult your state experimental station, university, or extension agent as to sensitivity of grass species or varieties to various herbicides. If the sensitivity of grass species or varieties is unknown, test Chlorsulf E-Pro 75 WDG Herbicide on a small area of the grass species. Tolerance to Chlorsulf E-Pro 75 WDG Herbicide of components in a grass seed mixture will vary and the final stand may not reflect the seed ratio.
- To avoid injury, do not apply Chlorsulf E-Pro 75 WDG Herbicide to grasses growing under conditions of stress (severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage). Injury to grasses is also possible if grasses are under stress before or after an application of Chlorsulf E-Pro 75 WDG Herbicide. Other weather conditions (such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures, drought, low fertility, water-saturated soils, disease, or insect damage) in effect before or after Chlorsulf E-Pro 75 WDG Herbicide applications may result in temporary discolorations and/or grass injury.
- Do not use this product on lawns, walks, driveways, tennis courts, or similar areas.
- Do not apply this product in or on irrigation or drainage ditches or canals including their outer banks.
- Do not allow Chlorsulf E-Pro 75 WDG Herbicide to drift or move into irrigation or drainage ditches.
- Do not apply through any type of irrigation system.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla, and Conejos.

There are no hay harvest restrictions or grazing restrictions for livestock (including lactating animals) when Chlorsulf E-Pro 75 WDG Herbicide is applied at rates up to 1-1/3 ounces per Acre.

INSTRUCTIONS FOR MIXING

1. Using clean fresh water, fill the spray tank 1/4 to 1/3 full.
2. Begin agitation and then add the required amount of Chlorsulf E-Pro 75 WDG Herbicide.
3. Allow the solution to agitate for 5 minutes to completely disperse the dry flowable Chlorsulf E-Pro 75 WDG Herbicide formulation.
4. Continue agitation and fill the spray tank with the remaining water. Do not add any other material until the Chlorsulf E-Pro 75 WDG Herbicide is thoroughly mixed with the water.
5. As the tank is filling with the remaining water, add any tank mix partners followed by the necessary volume of spray adjuvants. Always add the spray adjuvants last.
6. NOTE: Continuous agitation is required or settling will occur. Before spraying, reagitrate the solution to ensure a uniform solution is sprayed.
7. Make only a sufficient amount of Chlorsulf E-Pro 75 WDG Herbicide spray mixture that can be used within 24 hours of mixing. The product may degrade if allowed to sit unused.
8. For application of multiple loads of Chlorsulf E-Pro 75 WDG Herbicide and a tank mix partner, make a pre-slurry of Chlorsulf E-Pro 75 WDG Herbicide in clean water and then add to the spray tank. This pre-mix helps to prevent the tank mix partner from interfering with the dissolution of the Chlorsulf E-Pro 75 WDG Herbicide.

Do not mix Chlorsulf E-Pro 75 WDG Herbicide with spray additives that reduce the pH of the spray solution below 5.0. Additional information is found in the section, **SPRAY ADJUVANTS**.

HOW TO CLEAN SPRAYER EQUIPMENT

Clean all spray equipment before making an application of Chlorsulf E-Pro 75 WDG Herbicide.

Immediately after an application or multiple applications of Chlorsulf E-Pro 75 WDG Herbicide, clean all spray equipment using the cleanup procedures described on the labels of previously applied products. If there are no cleanup directions, use the following cleanup procedures.

Note: The directions for sprayer cleanup presented below are only effective for Chlorsulf E-Pro 75 WDG Herbicide and for general uses specified under "Directions for Use". Do not use the sprayer equipment on food crops (except wheat, barley and oats), feed crops (except range land, CRP and pasture), fine turf, ornamentals, and other desirable plants.

After spraying is completed at the end of the day, rinse the interior of the tank with fresh water. Partially refill the tank with fresh water and flush the boom and hoses. These rinses will prevent deposits of dried pesticide residues that can remain in the application equipment. Use the following steps to clean all mixing and spray equipment immediately following applications of Chlorsulf E-Pro 75 WDG Herbicide:

1. Drain the spray tank and then use fresh water to rinse the interior surfaces of the tank. Then flush the tank, boom, and hoses with water for at least 5 minutes.
2. Use fresh clean water to fill the tank and add a cleaning solution[†]. Flush the boom, hoses, and nozzles with the cleaning solution. Allow the equipment to sit for 15 minutes with agitation running, and then drain the tank.
3. Repeat Step 2.
4. Repeat Step 1.
5. Remove and clean the nozzles and screens separately. Traces of the cleaning solution can be removed by rinsing the tank thoroughly with clean water and flushing the water through the hoses and boom.

[†] Cleaning Solutions approved for spray equipment cleanup:

1. One gallon of 3% ammonia per 100 gallons of water.
2. "Nutra-sol" (carefully follow the directions for use on the "Nutra-sol" label).
3. Loveland Spray Tank Cleaner (carefully follow the directions for use on the Loveland Spray Tank Cleaner label).
4. "Tank-Cleaner" (carefully follow the directions for use on the "Tank-Cleaner" label).

TANK MIXTURES

Other herbicides which are registered for the same uses as Chlorsulf E-Pro 75 WDG Herbicide may be tank mixed with Chlorsulf E-Pro 75 WDG Herbicide. Use whichever label has the most restrictive directions for the tank mix. Do not tank mix Chlorsulf E-Pro 75 WDG Herbicide with DuPont™ HYVAR® X-L herbicide.

Before preparing large amounts of the tank mix, perform a jar test to insure the tank-mix partners are compatible with Chlorsulf E-Pro 75 WDG Herbicide. A clear jar with a lid can be used to mix the tank mix ingredients in their relative proportions. After adding the ingredients, invert the jar several times and then allow the jar to stand for 30 minutes. The mixture is compatible if, after 30 minutes, the solution remains mixed, or, if separation occurs, if the solution readily mixes again after agitation. Signs of incompatibility include separation into layers which do not readily remix when agitated, the presence of flakes, precipitates, gels, or heavy oily film on the jar.

SPRAY EQUIPMENT

Non-crop sites: Make applications of Chlorsulf E-Pro 75 WDG Herbicide using ground equipment only unless otherwise specified in Supplemental or Special Local Need Labeling.

Refer to the manufacturer's documentation for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Use calibrated air or ground equipment, and apply in a spray volume and delivery system to ensure a thorough, uniform spray coverage of weed pests. Higher spray volumes will produce better coverage to dense canopies of weeds. Do not overlap sprays. To avoid injury to desirable species, turn off spray booms while starting, turning, slowing, or stopping.

Severe injury or death of crops (excluding pasture, range and small grains) may occur if the same equipment used to apply Chlorsulf E-Pro 75 WDG Herbicide to pasture, range, CRP or non-crop sites is used to apply other products to crops. Traces of Chlorsulf E-Pro 75 WDG Herbicide in the spray equipment may injure or kill the crops (except pasture, range, and small grains).

Do not make applications using equipment and/or spray volumes or under weather conditions that might cause spray drift onto nontarget sites. Additional information is provided in the section, **SPRAY DRIFT MANAGEMENT**.

Use application equipment that will ensure constant agitation of Chlorsulf E-Pro 75 WDG Herbicide spray solutions.

GROUND APPLICATION

BROADCAST APPLICATION

Apply Chlorsulf E-Pro 75 WDG Herbicide at 20 to 40 GPA using calibrated ground broadcast application equipment. Optimum control is obtained when weeds are treated in a sufficient volume to receive a thorough, uniform coverage.

Industrial turf: Do not overlap sprays. To avoid injury to desirable species, turn off spray booms while starting, turning, slowing, or stopping.

HIGH VOLUME HANDGUN APPLICATION

Apply Chlorsulf E-Pro 75 WDG Herbicide at 100 to 300 GPA using calibrated hand-gun broadcast application equipment. Mix 1 ounce Chlorsulf E-Pro 75 WDG Herbicide in 100 gals. of water. Do not apply more than 300 gals. of spray mix per acre.

INVERT SPRAY APPLICATION

Apply the high viscosity invert solution of Chlorsulf E-Pro 75 WDG Herbicide at 10 to 40 GPA. Mix 1/4 to 3 ounces of Chlorsulf E-Pro 75 WDG Herbicide in the water phase of the invert solution for application to 1 Acre. The labeled use rate for target weeds is found in the section, **WEEDS CONTROLLED BY Chlorsulf E-Pro 75 WDG Herbicide**. Follow all use directions and precautionary statements appearing on the labels of the inverting oils and additives or in the operator's manual of the inverting equipment.

SPOT APPLICATIONS

NON-CROP SITES

Mix 1 to 3 ounces of Chlorsulf E-Pro 75 WDG Herbicide with 100 gallons of water. Do not apply more than 300 gallons of the Chlorsulf E-Pro 75 WDG Herbicide at the 1 ounce spray mix rate per Acre, and no more than 100 gallons of the Chlorsulf E-Pro 75 WDG Herbicide of the 3 ounce spray mix rate per Acre.

SPRAY ADJUVANTS

Include a high quality spray adjuvant (but not LI-700 or other acidifying adjuvants) with the Chlorsulf E-Pro 75 WDG Herbicide to improve postemergence weed control. Follow the manufacturer's labeled rate for the adjuvant.

SPRAY DRIFT CONTROL AGENTS

Include a spray drift control agent with the Chlorsulf E-Pro 75 WDG Herbicide tank mix to reduce the chance of drift. Follow the manufacturer's labeled rate for the drift control agent.

FIELD BIOASSAY

When crop or grass species/varieties which are not listed on this label are to be planted to areas previously treated with Chlorsulf E-Pro 75 WDG Herbicide, a field bioassay test must be carried out to determine if this species can be replanted without injury. Test the crop or grass intended to be planted the year following a treatment with Chlorsulf E-Pro 75 WDG Herbicide by growing the crop or grass in small plots which received the Chlorsulf E-Pro 75 WDG Herbicide treatment. The crop or grass response will determine the feasibility of rotating this crop or grass to large areas which had been treated with Chlorsulf E-Pro 75 WDG Herbicide. Additional information on the procedures for carrying out field bioassays can be obtained from your local dealer or Etigra representative.

GRAZING/HAYING

No hay harvest restrictions or grazing restrictions for livestock (including lactating animals) apply when Chlorsulf E-Pro 75 WDG Herbicide is applied at up to 1-1/3 ounces per Acre. Animals do not need to be enclosed.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

Apply Chlorsulf E-Pro 75 WDG Herbicide in a minimum of 3 GPA.

Solid stream nozzles which are oriented straight back must be used when applying Chlorsulf E-Pro 75 WDG Herbicide by air in areas adjacent to sensitive crops. Avoid spray drift damage to sensitive crops downwind by adjusting the swath. To minimize spray drift, apply Chlorsulf E-Pro 75 WDG Herbicide using ground equipment to treat border edges of fields. See additional information in the **SPRAY DRIFT MANAGEMENT** section, below.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce the drift potential is to apply large droplets (>150-200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **WIND, TEMPERATURE AND HUMIDITY**, and **SURFACE TEMPERATURE INVERSIONS** sections of this label.

CONTROLLING DROPLET SIZE – GENERAL TECHNIQUES

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

CONTROLLING DROPLET SIZE – AIRCRAFT

- Number of Nozzles – Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation – Orientating nozzles so that the spray is emitted backwards, parallel to the airstream, will produce larger droplets than other orientations.
- Nozzle Type – Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

BOOM LENGTH AND HEIGHT

- Boom length (aircraft) – The boom length should not exceed 3/4 of the wing length; using shorter booms decreases drift potential. For helicopters, use a boom length and position that prevents droplets from entering the rotor vortices.
- Boom Height (aircraft) – Application more than 10 ft. above the canopy increases the potential for spray drift.
- Boom Height (ground) – Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

NON-AGRICULTURAL USES

NON-CROP SITES - INDUSTRIAL AREAS

To control annual, biennial and perennial broadleaf weeds found in non-crop, industrial areas, apply Chlorsulf E-Pro 75 WDG Herbicide at the rates listed in the sections below and follow all directions for use on this label. Apply by ground equipment unless directed otherwise by Special Local Need or Supplemental labeling. Make preemergent or early postemergent spray applications of Chlorsulf E-Pro 75 WDG Herbicide to actively germinating or growing **annual** weeds. **Perennial weeds** are best controlled from application of Chlorsulf E-Pro 75 WDG Herbicide when weeds are in the bud to bloom or fall rosette stage.

INDUSTRIAL TURF (UNIMPROVED ONLY)

Directions for Application: To control weeds found in industrial turf (unimproved), on roadside or other non-crop sites, apply Chlorsulf E-Pro 75 WDG Herbicide at the rates listed in the table below and follow all directions for use on this label. The higher Chlorsulf E-Pro 75 WDG Herbicide rates will control weeds for longer periods of time compared with the lower Chlorsulf E-Pro 75 WDG Herbicide rates. Temporary chlorosis of desirable grasses may occur when Chlorsulf E-Pro 75 WDG Herbicide is applied at the higher rate or in combination with a surfactant.

Timing: Make applications of Chlorsulf E-Pro 75 WDG Herbicide when desirable grasses have become well-established to avoid any top kill or stand reduction. Optimum results are seen if turf is treated at green-up.

Weeds: Refer to the section **WEEDS CONTROLLED BY Chlorsulf E-Pro 75 WDG Herbicide**.

RATES FOR CONTROL OR SUPPRESSION OF WEEDS IN INDUSTRIAL TURF

| 1/4 to 1/2 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
|--|--|
| Fescue (<i>Festuca spp.</i>) | Smooth brome (<i>Bromus inermis</i>) |
| 1/2 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
| Bentgrass (<i>Agrostis spp.</i>) | Kleingrass (<i>Panicum coloratum</i>) |
| Bluestems, big, little, plains, sand, ww spar (<i>Andropogon spp.</i>) | Lovegrasses, sand, weeping (<i>Eragrostis spp.</i>) |
| Buffalograss (<i>Buchloe dactyloides</i>) | Prairie sandreed (<i>Calamovilfa longifolia</i>) |
| Galleta (<i>Hilaria jamesii</i>) | Sheep fescue (<i>Festuca ovina</i>) |
| Needlegrass, green (<i>Stipa viridula</i>) | Sideoats gramma (<i>Bouteloua curtipendula</i>) |
| Green sprangletop (<i>Leptochloa dubia</i>) | Switchgrass (<i>Panicum virgatum</i>) |
| Indiangrass (<i>Sorghastrum nutans</i>) | Wildrye grasses, beardless, Russian (<i>Elymus spp.</i>) |
| Indian ricegrass (<i>Oryzopsis hymenoides</i>) | |
| 1/4 to 1 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
| Bahiagrass (<i>Paspalum notatum</i>) | Bromegrass, meadow, smooth (<i>Bromus spp.</i>) |
| Bermudagrass (<i>Cynodon dactylon</i>) | Orchardgrass (<i>Dactylis glomerata</i>) |
| Blue gramma (<i>Bouteloua gracilis</i>) | Wheatgrasses, crested, intermediate, pubescent, slender, streambank, tall, thick, spike, western (<i>Agropyron spp.</i>) |
| Bluegrass (<i>Poa spp.</i>) | |

GROWTH SUPPRESSION AND SEEDHEAD INHIBITION

Directions for Application: To suppress grass growth (chemical mowing) and inhibit seedhead formation, apply Chlorsulf E-Pro 75 WDG Herbicide as a tank mix with other herbicides registered for this use and at the rates listed in the table below. Follow all directions for use on this label. The higher Chlorsulf E-Pro 75 WDG Herbicide rates will control weeds for longer periods of time compared with the lower Chlorsulf E-Pro 75 WDG Herbicide rates.

Timing: Make applications of Chlorsulf E-Pro 75 WDG Herbicide when desirable grasses have become well-established to avoid any top kill or stand reduction. Time application to occur at green-up and before seed-heads emerge (boot stage).

Weeds: Refer to the section **WEEDS CONTROLLED BY Chlorsulf E-Pro 75 WDG Herbicide**.

Precautions:

- To avoid injury, do not use Chlorsulf E-Pro 75 WDG Herbicide or Chlorsulf E-Pro 75 WDG Herbicide in a tank mix with Embark® 2S on bahiagrass turf or turf that is under stress (due to drought, insects, disease, cold temperature, or poor fertility).
- To avoid injury, apply Chlorsulf E-Pro 75 WDG Herbicide only to turf that has been established for at least 1 year.
- Wait 6 months after an application of Chlorsulf E-Pro 75 WDG Herbicide before planting grass seed in treated areas. Cultivate the area before planting.
- To avoid turf injury, only make spot applications to control those weeds listed under the 1 to 3 oz. rate in the **NON-CROP, INDUSTRIAL SITES** section of this label. Broadcast applications to turf at this 1 to 3 oz. rate may cause excessive turf injury.

Restrictions:

- Broadcast applications: do not apply more than 1/2 ounce Chlorsulf E-Pro 75 WDG Herbicide per Acre per year (12 months).

RATES FOR GROWTH SUPPRESSION AND SEEDHEAD INHIBITION

| 1/4 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide PLUS 1/4 to 1/2 pt. Embark® 2S | |
|---|--|
| Fescue (<i>Festuca spp.</i>) | Bluegrass (<i>Poa spp.</i>) |
| 1/2 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide PLUS 1/2 to 1 pt. Embark® 2S † | |
| Fescue (<i>Festuca spp.</i>) | Smooth brome (<i>Bromus inermis</i>) |
| Annual bluegrass (<i>Poa annua</i>) | Orchardgrass (<i>Dactylis glomerata</i>) |
| Perennial ryegrass (<i>Lolium perenne</i>) | Reed canarygrass (<i>Phalaris arundinacea</i>) |
| † For use in the Pacific Northwest Only | |

WEEDS CONTROLLED BY Chlorsulf E-Pro 75 WDG Herbicide

NOTE: The higher Chlorsulf E-Pro 75 WDG Herbicide rates will control weeds for longer periods of time compared with the lower Chlorsulf E-Pro 75 WDG Herbicide rates.

TABLE OF WEEDS CONTROLLED AT DIFFERENT Chlorsulf E-Pro 75 WDG Herbicide USE RATES

| 1/4 to 1/2 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
|---|---|
| Annual sowthistle (<i>Sonchus oleraceus</i>) | London rocket (<i>Sisymbrium irio</i>) † |
| Blue mustard (<i>Chorispora tenella</i>) | Mayweed (<i>Anthemis cotula</i>) † |
| Common chickweed (<i>Stellaria media</i>) | Miner's lettuce (<i>Montia perfoliata</i>) † |
| Common speedwell (<i>Veronica officinalis</i>) | Pineapple-weed (<i>Matricaria matricarioides</i>) † |
| Common spikeweed (<i>Hemizonia pungens</i>) † | Prostrate pigweed (<i>Amaranthus blitoides</i>) † |
| Conical catchfly (<i>Silene conoidea</i>) † | Redroot pigweed (<i>Amaranthus retroflexus</i>) |
| Cutleaf eveningprimrose (<i>Oenothera laciniata</i>) † | Shepherd's purse (<i>Capsella bursa-pastoris</i>) † |
| Fiddleneck, tarweed (<i>Amsinckia lycopsoides</i>) † | Smooth pigweed (<i>Amaranthus chlorostachys</i>) † |
| Field pennycress (<i>Thlaspi arvense</i>) | Treacle mustard (<i>Erysimum spp.</i>) †† |
| Flixweed (<i>Descurainia Sophia</i>) | Tumble mustard, Jim Hill (<i>Sisymbrium altissimum</i>) |
| Hempnettle (<i>Galeopsis spp.</i>) † | Wild mustard (<i>Sinapis arvensis</i>) |
| Henbit (<i>Lamium amplexicaule</i>) | |
| 1/2 to 1 ounce per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
| Bouncingbet (<i>Saponaria officinalis</i>) | Groundsel, common (<i>Senecio vulgaris</i>) † |
| Bur beakchervil (<i>Anthriscus caucalis</i>) † | Musk thistle (<i>Carduus nutans</i>) |
| Buttercup (<i>Ranunculus spp.</i>) | Smallseed falseflax (<i>Camelina microcarpa</i>) † |
| Carolina geranium (<i>Geranium carolinianum</i>) † | Sweet clover (<i>Melilotus spp.</i>) †† |
| Common lambsquarter (<i>Chenopodium album</i>) | Tumble pigweed (<i>Amaranthus albus</i>) † |
| Common sunflower (<i>Helianthus annuus</i>) | Turkey mullein (<i>Eremocarpus setigerus</i>) †† |
| Dandelion, common (<i>Taraxacum officinale</i>) †† | Whitetop, hoar cress (<i>Cardaria draba</i>) ††† |
| Erect knotweed (<i>Polygonum erectum</i>) † | Wild buckwheat (<i>Polygonum convolvulus</i>) † |
| Goldenrod (<i>Solidago spp.</i>) | Wild parsnip (<i>Pastinaca sativa</i>) |
| 1 to 3 ounces per Acre of Chlorsulf E-Pro 75 WDG Herbicide | |
| Asters (<i>Aster spp.</i>) | Horsetail (<i>Equisetum spp.</i>) |
| Bedstraw (<i>Galium spp.</i>) | Italian ryegrass (<i>Lolium multiflorum</i>) †† |
| Black mustard (<i>Brassica nigra</i>) | Marestail/horseweed (<i>Conyza canadensis</i>) |
| Bull thistle (<i>Cirsium vulgare</i>) | Pepperweed (<i>Lepidium spp.</i>) † |
| Burclover (<i>Medicago spp.</i>) | Pepperweed (perennial) (<i>Lepidium latifolium</i>) |
| Canada thistle (<i>Cirsium arvense</i>) | Poison-hemlock (<i>Conium maculatum</i>) |
| Common cinquefoil (<i>Potentilla canadensis</i>) | Prostrate knotweed (<i>Polygonum aviculare</i>) |
| Common mallow (<i>Malva neglecta</i>) | Puncturevine (<i>Tribulus terrestris</i>) |
| Common mullein (<i>Verbascum thapsus</i>) | Red clover (<i>Trifolium pretense</i>) † |
| Common ragweed (<i>Ambrosia elatior</i>) †† | Russian knapweed (<i>Acroptilon repens</i>) ††† |
| Common tansy (<i>Tanacetum vulgare</i>) | Scotch thistle (<i>Onopordum acanthium</i>) |
| Common teasel (<i>Dipsacus fullonum</i>) | Scouringrush (<i>Equisetum hyemale</i>) |
| Common yarrow (<i>Achillea millefolium</i>) | Spreading orach (<i>Atriplex patula</i>) |
| Corn spurry (<i>Spergula arvensis</i>) | Tansymustard (<i>Descurainia pinnata</i>) |
| Cow cockle (<i>Vaccaria pyramidata</i>) | Tansy ragwort (<i>Senecio jacobaea</i>) † |
| Curly dock (<i>Rumex crispus</i>) | White clover (<i>Trifolium repens</i>) |
| Dyer's woad (<i>Isatis tinctoria</i>) | Wild carrot (<i>Daucus carota</i>) |
| False chamomile (<i>Matricaria maritima</i>) † | Wild garlic/wild onion (<i>Allium vineale</i>) |
| Foxtails (<i>Setaria spp.</i>) †† | Yellow starthistle (<i>Centaurea solstitialis</i>) †† |
| <p>† Do not use Chlorsulf E-Pro 75 WDG Herbicide on this weed in California. †† Chlorsulf E-Pro 75 WDG Herbicide provides only partial control of this weed. ††† Time application to occur at prebloom to bloom and fall rosette stage.</p> | |

ADDITIONAL DIRECTIONS FOR SPECIFIC WEEDS

Dalmation Toadflax (*Linaria genistifolia*): For optimum control, apply in the fall at a rate of 2 to 3 ounces of Chlorsulf E-Pro 75 WDG Herbicide per acre as a high volume foliar spray (minimum of 24 gallons of water per acre) plus a surfactant (refer to **SPRAY ADJUVANTS** section, above).

Kochia, Russian Thistle, and Prickly Lettuce: For optimum results, apply after the weeds have emerged but before mature seeds form. Use a tank mix of Chlorsulf E-Pro 75 WDG Herbicide with herbicides with different modes of action (such as 2,4-D plus dicamba).

Yellow Toadflax (*Linaria vulgaris*): For optimum control, use a minimum of 1.5 ounces of Chlorsulf E-Pro 75 WDG Herbicide per acre.

Yellow Starthistle (*Centaurea solstitialis*): Use a tank mix of Chlorsulf E-Pro 75 WDG Herbicide at 1/2 to 3 oz. per Acre with other herbicides registered for this use (such as Transline®, Tordon® 22K or 2,4-D) at the tank-mix partner label rates. Refer to the **TANK MIX** section above. Add a surfactant to improve control of emerged weeds (refer to **SPRAY ADJUVANTS** section above). For preemergence control of this weed (early emergence to bolting stage of growth), apply when rainfall is expected so that residues of Chlorsulf E-Pro 75 WDG Herbicide reach the root zone. Note: the higher Chlorsulf E-Pro 75 WDG Herbicide rates will control weeds for longer periods of time compared with the lower Chlorsulf E-Pro 75 WDG Herbicide rates.

TIMING FOR REPLANTING OF GRASSES

Non-crop areas that were treated in the spring or early summer with Chlorsulf E-Pro 75 WDG Herbicide may be replanted with grasses after the minimum time periods noted in the tables below have elapsed. If an application of Chlorsulf E-Pro 75 WDG Herbicide is made in late summer or early fall, replanting may be carried out after the minimum time periods noted in the tables below have elapsed **starting from the spring after the application took place**. When replanting grasses, the tolerance of different grass varieties and species to soils treated with Chlorsulf E-Pro 75 WDG Herbicide may differ. If a grass species to be seeded is not listed in the tables below, carry out a field bioassay test (see section **FIELD BIOASSAY**, above) to determine if this species can be replanted without injury.

| Species | Chlorsulf E-Pro 75 WDG Herbicide Rate (oz. per Acre) | Replant Interval, Days |
|---|--|------------------------|
| For soils with a pH of 7.5 or less: | | |
| Brome, meadow (<i>Bromus erectus</i>) | 1/2 to 1 | 30 |
| | 1 to 2 | 60 |
| Brome, smooth (<i>Bromus inermis</i>) | 1/2 to 1 | 60 |
| | 1 to 2 | 120 |
| Fescue, alta/tall (<i>Festuca arundinacea</i>) | 1/2 | 60 |
| | 1 | 90 |
| | 2 | 150 |
| Fescue, sheep (<i>Festuca ovina</i>) | 1/2 to 1 | 60 |
| | 1 to 2 | 120 |
| Foxtail, meadow (<i>Alopecurus pratensis</i>) | 1/2 | 90 |
| | 1 | 120 |
| | 2 | 180 |
| Needlegrass, green (<i>Stip viridula</i>) | 1/2 to 2 | 30 |
| Orchardgrass (<i>Dactylis glomerata</i>) | 1/2 | 60 |
| | 1 to 2 | 90 |
| Russian wildrye (<i>Elymus spp.</i>) | 1/2 to 2 | 30 |
| Switchgrass (<i>Panicum virgatum</i>) | 1/2 to 2 | 90 |
| Timothy (<i>Phleum pratense</i>) | 1/2 | 60 |
| | 1 | 120 |
| | 2 | 180 |
| Wheatgrass, western (<i>Agropyron smithii</i>) | 1/2 | 30 |
| | 1 | 60 |
| | 2 | 120 |
| For soils having a pH of 7.5 and greater: | | |
| Alkali sacaton (<i>Sporobolus airoides</i>) | 1/2 | 30 |
| | 1 | 90 |
| | 2 | >90 |
| Bluestem, Big (<i>Andropogon gerardii</i>) | 1/2 | 90 |
| Brome, Mountain (<i>Bromus marginatus</i>) | 1/2 | 30 |
| | 1 | 60 |
| | 2 | >90 |
| Gamma, Blue (<i>Bouteloua gracilis</i>) | 1/2 | 30 |
| | 1 | 60 |
| | 2 | >90 |
| Gamma, Sideoats (<i>Bouteloua curtipendula</i>) | 1 to 2 | >90 |
| Switchgrass (<i>Panicum virgatum</i>) | 1 to 2 | >90 |
| Wheatgrass, Bluebunch (<i>Agropyron spicatum</i>) | 1-1/3 | 30 |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Etigra nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Etigra.

Chlorsulf E-Pro 75 WDG Herbicide is not manufactured or distributed by DuPont, seller of Telar®.

Tordon® and Transline® are trademarks of Dow AgroSciences LLC.
Hyvar® and Telar® are trademarks of E.I. DuPont de Nemours and Company.
Etigra® is a trademark of Etigra.
Embark® is a trademark of PBI-Gordon Corp.

EPA 20071210